

HAWKES PROCESSES ON LARGE NETWORKS

SYLVAIN DELATTRE, NICOLAS FOURNIER AND MARC HOFFMANN

ABSTRACT. We generalise the construction of multivariate Hawkes processes to a possibly infinite network of counting processes on a directed graph \mathbb{G} . The process is constructed as the solution to a system of Poisson driven stochastic differential equations, for which we prove pathwise existence and uniqueness under some reasonable conditions.

We next investigate how to approximate a standard N -dimensional Hawkes process by a simple inhomogeneous Poisson process in the mean-field framework where each pair of individuals interact in the same way, in the limit $N \rightarrow \infty$. In the so-called linear case for the interaction, we further investigate the large time behaviour of the process. We study in particular the stability of the central limit theorem when exchanging the limits $N, T \rightarrow \infty$ and exhibit different possible behaviours.

We finally consider the case $\mathbb{G} = \mathbb{Z}^d$ with nearest neighbour interactions. In the linear case, we prove some (large time) laws of large numbers and exhibit different behaviours, reminiscent of the infinite setting. Finally we study the propagation of a *single impulsion* started at a given point of \mathbb{Z}^d at time 0. We compute the probability of extinction of such an impulsion and, in some particular cases, we can accurately describe how it propagates to the whole space.

Mathematics Subject Classification (2010): 60F05, 60G55, 60G57.

Keywords: Point processes. Multivariate Hawkes processes. Stochastic differential equations. Limit theorems. Mean-field approximations. Interacting particle systems.

1. INTRODUCTION

1.1. Motivation. In several apparently different applied fields, a growing interest has been observed recently for a better understanding of stochastic interactions between multiple entities evolving through time. These include: seismology for modelling earthquake replicas (Helmstetter-Sornette [19], Kagan [25], Ogata [31], Bacry-Muzy [5]), neuroscience for modelling spike trains in brain activity (Grün *et al.* [14], Okatan *et al.* [32], Pillow *et al.* [33], Reynaud *et al.* [35, 36]), genome analysis (Reynaud-Schbath [34]), financial contagion (Ait-Sahalia *et al.* [1]), high-frequency finance (order arrivals, see Bauwens-Hautsch [6], Hewlett [20], market micro-structure see Bacry *et al.* [2] and market impact see Bacry-Muzy [4, 5]), financial price modelling across scales (Bacry *et al.* [3], Jaisson-Rosenbaum [22]), social networks interactions (Blundell *et al.* [7], Simma-Jordan [38], Zhou *et al.* [40]) and epidemiology like for instance viral diffusion on a network (Hang-Zha [39]), to name but a few. In all these contexts, observations are often represented as events (like spikes or features) associated to agents or nodes on a given network, and that arrive randomly through time but that are not stochastically independent.

In practice, we observe a multivariate counting process $(Z_t^1, \dots, Z_t^N)_{t \geq 0}$, each component Z_t^i recording the number of events of the i -th component of the system during $[0, t]$, or equivalently the time stamps of the observed events. Under relatively weak general assumptions, a multivariate counting process $(Z_t^1, \dots, Z_t^N)_{t \geq 0}$ is characterised by its intensity process $(\lambda_t^1, \dots, \lambda_t^N)_{t \geq 0}$, informally defined by

$$\Pr(Z^i \text{ has a jump in } [t, t + dt] \mid \mathcal{F}_t) = \lambda_t^i dt, \quad i = 1, \dots, N,$$

where \mathcal{F}_t denotes the sigma-field generated by $(Z^i)_{1 \leq i \leq N}$ up to time t . For modelling the interactions, a particularly attractive family of multivariate point processes is given by the class of (mutually exciting) Hawkes processes (Hawkes [16], Hawkes-Oakes [17]), with intensity process given by

$$\lambda_t^i = h_i \left(\sum_{j=1}^N \int_0^{t-} \varphi_{ji}(t-s) dZ_s^j \right),$$

where the *causal* functions $\varphi_{ji} : [0, \infty) \rightarrow \mathbb{R}$ model how Z^j acts on Z^i by affecting its intensity process λ^i . The nonnegative functions h_i account for some non-linearity, but if we set $h_i(x) = \mu_i + x$ with $\mu_i \geq 0$, we obtain *linear Hawkes processes* where μ_i can be interpreted as a baseline Poisson intensity. In the degenerate case $\varphi_{ji} = 0$, we actually retrieve standard Poisson processes.

Multivariate Hawkes processes have long been studied in probability theory (see for instance the comprehensive textbook of Daley-Vere-Jones [11] and the references therein, Brémaud-Massoulié [8] or the recent results of Zhu [41, 42]). Their statistical inference is relatively well understood too, from a classical parametric angle (Ogata [29]) together with recent significant advances in nonparametrics (Reynaud-Bouret-Schbath [34], Hansen *et al.* [18]). However, the frontier is progressively moving to understanding the case of large N , when the number of components may become increasingly large or possibly infinite, see Massoulié [28] and Galvez-Löcherbach [15] for some constructions in that direction. This context is potentially of major importance for future developments in the aforementioned applied fields. This is the topic of the present paper.

1.2. Setting. We work on a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \Pr)$. We say that $(X_t)_{t \geq 0}$ is a counting process if it is non-decreasing, càdlàg, integer-valued (and finite for all times), with all its jumps of height 1. For $(X_t)_{t \geq 0}$ a $(\mathcal{F}_t)_{t \geq 0}$ -adapted counting process, there is a unique non-decreasing predictable process $(\Lambda_t)_{t \geq 0}$, called *compensator* of $(X_t)_{t \geq 0}$, such that $(X_t - \Lambda_t)_{t \geq 0}$ is a $(\mathcal{F}_t)_{t \geq 0}$ -local martingale, see Jacod-Shiryaev [23, Chapter I].

We consider a countable directed graph $\mathbb{G} = (\mathcal{S}, \mathcal{E})$ with vertices (or nodes) $i \in \mathcal{S}$ and (directed) edges $e \in \mathcal{E}$. We write $e = (j, i) \in \mathcal{E}$ for the oriented edge. We also need to specify the following parameters: a kernel $\varphi = (\varphi_{ji}, (j, i) \in \mathcal{E})$ with $\varphi_{ji} : [0, \infty) \mapsto \mathbb{R}$, and a nonlinear intensity component $\mathbf{h} = (h_i, i \in \mathcal{S})$ with $h_i : \mathbb{R} \mapsto [0, \infty)$. The natural generalisation of finite-dimensional Hawkes processes is the following.

Definition 1. A Hawkes process with parameters $(\mathbb{G}, \varphi, \mathbf{h})$ is a family of $(\mathcal{F}_t)_{t \geq 0}$ -adapted counting processes $(Z_t^i)_{i \in \mathcal{S}, t \geq 0}$ such that

- (i) almost surely, for all $i \neq j$, $(Z_t^i)_{t \geq 0}$ and $(Z_t^j)_{t \geq 0}$ never jump simultaneously,
- (ii) for every $i \in \mathcal{S}$, the compensator $(\Lambda_t^i)_{t \geq 0}$ of $(Z_t^i)_{t \geq 0}$ has the form $\Lambda_t^i = \int_0^t \lambda_s^i ds$, where the intensity process $(\lambda_t^i)_{t \geq 0}$ is given by

$$\lambda_t^i = h_i \left(\sum_{j \rightarrow i} \int_0^{t-} \varphi_{ji}(t-s) dZ_s^j \right),$$

with the notation $\sum_{j \rightarrow i}$ for summation over $\{j : (j, i) \in \mathcal{E}\}$.

This model can be seen as a particular case of that introduced by Massoulié [28] who considers more general sets of sites (possibly \mathbb{R}^d) and a larger class of intensities. We say that a Hawkes process is *linear* when $h_i(x) = \mu_i + x$ for every $x \in \mathbb{R}$, $i \in \mathcal{S}$, with $\mu_i \geq 0$ and when $\varphi_{ji} \geq 0$.

We will give some general existence, uniqueness and approximation results for *nonlinear* Hawkes processes, but all the precise large-time estimates we will prove concern the linear case.

A Hawkes process $(Z_t^i)_{i \in \mathcal{S}, t \geq 0}$ with parameters $(\mathbb{G}, \varphi, \mathbf{h})$ behaves as follows. For each $i \in \mathcal{S}$, the rate of jump of Z^i is, at time t , $\lambda_i(t) = h_i(\sum_{j \rightarrow i} \sum_{k \geq 1} \varphi_{ji}(t - T_k^j) \mathbf{1}_{\{T_k^j < t\}})$, where $(T_k^j)_{k \geq 1}$ are the jump times of Z^j . In other words, each time one of the Z^j 's has a jump, it *excites* its *neighbours* in that it increases their rate of jump (in the natural situation where h is increasing and φ is positive). If φ is positive and decreases to 0, the case of almost all applications we have in mind, the influence of a jump decreases and tends to 0 as time evolves.

1.3. Main results. In the case where \mathbb{G} is a finite graph, under some appropriate assumptions on the parameters, the construction of $(Z_t^i)_{i \in \mathcal{S}, t \geq 0}$ is standard. However, for an infinite graph, the situation is more delicate: we have to check, in some sense, that the interaction does not come from infinity.

The first part of this paper (Section 2) consists of writing a Hawkes process as the solution to a system of Poisson-driven S.D.E.s and of finding a set of assumptions on \mathbb{G} and on the parameters (φ, \mathbf{h}) under which we can prove the pathwise existence and uniqueness for this system of S.D.E.s. Representing counting processes as solutions to S.D.E.s is classical, see Lewis-Shedler [27], Ogata, [30], Brémaud-Massoulié [8], Chevallier [10]. However, the well-posedness of such S.D.E.s is not obvious when \mathbb{G} is an infinite graph.

In a second part (Section 3), we study the *mean-field* situation: we assume that we have a finite (large) number N of *particles* behaving similarly, with no geometry. In other words, $\mathcal{S} = \{1, \dots, N\}$ is endowed with the set of all possible edges $\mathcal{E} = \{(i, j) : i, j \in \mathcal{S}\}$, and there are two functions h and φ such that $h_i = h$ and $\varphi_{ij} = N^{-1}\varphi$ for all $i, j \in \mathcal{S}$. We show that, as $N \rightarrow \infty$, Hawkes processes can be approximated by an i.i.d. family of inhomogeneous Poisson processes. Concerning the large-time behaviour, we discuss, in the linear case, the possible law of large numbers and central limit theorems as $(t, N) \rightarrow (\infty, \infty)$ and we observe some different situations according to the position of $\int_0^\infty \varphi(t) dt$ with respect to 1 (the so-called critical case).

Finally, we consider in Section 4 the case where \mathbb{G} is \mathbb{Z}^d , endowed with the set of edges $\mathcal{E} = \{(i, j) : |i - j| = 0 \text{ or } 1\}$, where $|\cdot|$ denotes the Euclidean distance. We study the large time behaviour, in the linear case where $h_i(x) = \mu_i + x$ and when $\varphi_{ij} = (2d + 1)^{-1}\varphi$ does not depend on i, j . We first assume that μ_i does not depend too much on i (consider *e.g.* the case where the μ_i are random, i.i.d. and bounded) and show that (i) if $\int_0^\infty \varphi(t) dt > 1$, then there is a law of large numbers and the interaction makes everything *flat*, in the sense that for all $i \neq j$, $Z_t^i \sim Z_t^j$ as $t \rightarrow \infty$; (ii) if $\int_0^\infty \varphi(t) dt < 1$, then there is again a law of large numbers, but the limiting value depends on i . We also explain why these results are reminiscent of the infinite setting and of the interaction. Finally, we study the case where $\mu_i = 0$ for all i but where there is an *impulsion* at time 0 at $i = 0$. We compute the probability of extinction of such an impulsion and, in some particular cases, we study how it propagates to the whole space (when it does not blow out).

1.4. Notation. The Laplace transform of $\varphi : [0, \infty) \mapsto \mathbb{R}$ is defined, when it exists, by

$$\mathcal{L}_\varphi(\alpha) = \int_0^\infty e^{-\alpha t} \varphi(t) dt.$$

We also introduce the convolution of $h, g : [0, \infty) \mapsto \mathbb{R}$ as (if it exists) $(g \star h)_t = \int_0^t g_s h_{t-s} ds = \int_0^t g_{t-s} h_s ds$. As is well-known, when everything makes sense, $\mathcal{L}_{g \star h}(\alpha) = \mathcal{L}_g(\alpha) \times \mathcal{L}_h(\alpha)$.

2. WELL-POSEDNESS USING A POISSON S.D.E.

We will study Hawkes processes through a system of Poisson-driven stochastic differential equations. This will allow us to speak of pathwise existence and uniqueness and to prove some propagation of chaos using some simple *coupling* arguments.

Consider, on a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \Pr)$, a family $(\pi^i(ds dz), i \in \mathcal{S})$ of i.i.d. $(\mathcal{F}_t)_{t \geq 0}$ -Poisson measures with intensity measure $ds dz$ on $[0, \infty) \times [0, \infty)$.

Definition 2. A family $(Z_t^i)_{i \in \mathcal{S}, t \geq 0}$ of càdlàg $(\mathcal{F}_t)_{t \geq 0}$ -adapted processes is called a Hawkes process with parameters $(\mathbb{G}, \boldsymbol{\varphi}, \mathbf{h})$ if a.s., for all $i \in \mathcal{S}$, all $t \geq 0$

$$(1) \quad Z_t^i = \int_0^t \int_0^\infty \mathbf{1}_{\{z \leq h_i(\sum_{j \rightarrow i} \int_0^{s-} \varphi_{ji}(s-u) dZ_u^j)\}} \pi^i(ds dz).$$

This formulation is consistent with Definition 1.

Proposition 3. (a) A Hawkes process in the sense of Definition 2 is also a Hawkes process in the sense of Definition 1.

(b) Consider a Hawkes process in the sense of Definition 1 (on some filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \Pr)$). Then we can build, on a possibly enlarged probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{F}}_t)_{t \geq 0}, \tilde{\Pr})$, a family $(\tilde{\pi}^i(ds dz), i \in \mathcal{S})$ of i.i.d. $(\tilde{\mathcal{F}}_t)_{t \geq 0}$ -Poisson measures with intensity measure $ds dz$ on $[0, \infty) \times [0, \infty)$ such that $(Z_t^i)_{i \in \mathcal{S}, t \geq 0}$ is a Hawkes process in the sense of Definition 2.

Point (a) is very easy: for a Hawkes process $(Z_t^i)_{i \in \mathcal{S}, t \geq 0}$ in the sense of Definition 2, it is clear that for every $i \in \mathcal{S}$, the compensator of Z^i is $\int_0^t \int_0^\infty \mathbf{1}_{\{z \leq h_i(\sum_{j \rightarrow i} \int_0^{s-} \varphi_{ji}(s-u) dZ_u^j)\}} dz ds$, which is equal to $\int_0^t h_i(\sum_{j \rightarrow i} \int_0^{s-} \varphi_{ji}(s-u) dZ_u^j) ds$. Furthermore, the independence of the Poisson random measures $(\pi^i(ds dz), i \in \mathcal{S})$ guarantees that for all $i \neq j$, $(Z_t^i)_{t \geq 0}$ and $(Z_t^j)_{t \geq 0}$ a.s. never jump simultaneously.

Point (b) is more delicate but standard and a very similar result was given in Brémaud-Massoulié [8]. Their proof is based on results found in the book [21] of Jacod, of which one of the main goals is exactly this topic: prove the equivalence between martingale problems and S.D.E.s. We also refer to Chevallier [10, Section IV] where a very complete proof is given as well as a historical survey. Let us mention that the idea to integrate an indicator function with respect to a Poisson measure in order to produce an inhomogeneous Poisson process with given intensity was first introduced by Lewis-Shedler [27], and later extended by Ogata [30] in the case of a stochastic intensity.

The following set of assumptions will guarantee the well-posedness of (1).

Assumption 4. There are some nonnegative constants $(c_i)_{i \in \mathcal{S}}$, some positive weights $(p_i)_{i \in \mathcal{S}}$ and a locally integrable function $\phi : [0, \infty) \mapsto [0, \infty)$ such that

- (a) for every $i \in \mathcal{S}$, every $x, y \in \mathbb{R}$, $|h_i(x) - h_i(y)| \leq c_i |x - y|$,
- (b) $\sum_{i \in \mathcal{S}} h_i(0) p_i < \infty$,
- (c) for every $s \in [0, \infty)$, every $j \in \mathcal{S}$, $\sum_{i, (j,i) \in \mathcal{E}} c_i p_i |\varphi_{ji}(s)| \leq p_j \phi(s)$.

Let us give a few examples of parameters $(\mathbb{G}, \boldsymbol{\varphi}, \mathbf{h})$ satisfying Assumption 4.

Remark 5. (i) If \mathcal{S} is finite, then Assumption 4 holds true, with the choice $p_i = 1$, as soon as h_i is Lipschitz continuous for all $i \in \mathcal{S}$ and φ_{ji} is locally integrable for all $(j, i) \in \mathcal{E}$.

(ii) If $\mathcal{S} = \mathbb{Z}^d$ is endowed with $\mathcal{E} = \{(i, j) : |i - j| = 0 \text{ or } 1\}$, then Assumption 4 holds, with the choice $p_i = 2^{-|i|}$, if $\sum_{i \in \mathbb{Z}^d} 2^{-|i|} |h_i(0)| < \infty$ and if there are $c > 0$ and $\varphi \in L^1_{loc}([0, \infty))$ such that $|h_i(x) - h_i(y)| \leq c|x - y|$ and $|\varphi_{jk}(t)| \leq \varphi(t)$ for all $i \in \mathcal{S}$, $x, y \in \mathbb{R}$, $(j, k) \in \mathcal{E}$ and $t \geq 0$.

(iii) Consider next $\mathcal{S} = \mathbb{Z}^d$ endowed with the set of all possible edges $\mathcal{E} = \{(i, j) : i, j \in \mathbb{Z}^d\}$ and assume that there is $c > 0$ such that $|h_i(0)| \leq c$ and $|h_i(x) - h_i(y)| \leq c|x - y|$ for all $i \in \mathcal{S}$, $x, y \in \mathbb{R}$. Assume that there are $\varphi \in L^1_{loc}([0, \infty))$ and a nonincreasing $a : [0, \infty) \mapsto [0, \infty)$ such that $|\varphi_{ji}(t)| \leq a(|i - j|)\varphi(t)$ for all $(i, j) \in \mathcal{E}$ and $t \geq 0$. Then if $\sum_{i \in \mathbb{Z}^d} a(|i|) < \infty$, Assumption 4 holds true.

(iv) Consider the (strongly oriented) graph \mathbb{Z}_+ endowed with the set of edges $\mathcal{E} = \{(i, i + 1) : i \in \mathbb{Z}_+\}$. Then Assumption 4 holds true as soon as there is $\varphi \in L^1_{loc}([0, \infty))$ such that for every $i \in \mathbb{Z}_+$, there are $c_i > 0$ and $a_i > 0$ such that $|h_i(x) - h_i(y)| \leq c_i|x - y|$ and $|\varphi_{i(i+1)}| \leq a_i\varphi$.

Points (ii) and (iii) of course extend to other graphs. In (iv), there is no growth condition on $|h_i(0)|$, c_i and a_i . This comes from the fact that the interaction is directed: Z^0 is actually a Poisson process with rate $h_0(0)$, the intensity of Z^1 is entirely determined by that of Z^0 , and so on. Hence this example is not very interesting. But we can mix e.g. points (ii) and (iv): informally, coefficients corresponding to edges directed to the origin have to be well-controlled, while coefficients corresponding to edges directed to infinity require less assumptions.

Proof. Point (i) is obvious. To check (ii), simply note that for all $j \in \mathcal{S}$, $\sum_{i, (j,i) \in \mathcal{E}} c 2^{-|i|} |\varphi_{ji}| \leq c\varphi 2^{-|j|} \sum_{i, (j,i) \in \mathcal{E}} 2^{|j|-|i|} \leq c 2(2d + 1)\varphi 2^{-|j|}$ and define $\phi = c 2(2d + 1)\varphi$. Point (iv) holds with $(p_i)_{i \in \mathbb{Z}_+}$ defined by $p_0 = 1$ and, by induction, $p_{i+1} = \min\{2^{-i}/(1 + h_{i+1}(0)), p_i/(1 + a_i c_{i+1})\}$. This of course implies that $\sum_{i \in \mathbb{Z}_+} p_i |h_i(0)| < \infty$ and that for all $j \geq 1$, $\sum_{i, (j,i) \in \mathcal{E}} c_i p_i |\varphi_{ji}| = c_{j+1} p_{j+1} |\varphi_{j(j+1)}| \leq c_{j+1} p_{j+1} a_j \varphi \leq p_j \varphi$ as desired.

To prove (iii), we work with the sup norm $|i| = |(i_1, \dots, i_d)| = \max\{|i_1|, \dots, |i_d|\}$. The delicate part consists in showing that there is $b : \mathbb{N} \mapsto [0, \infty)$ and a constant $C > 0$ such that $\sum_{i \in \mathbb{Z}^d} b(|i|) < \infty$ and, for all $j \in \mathbb{Z}^d$, $\sum_{i \in \mathbb{Z}^d} b(|i|) a(|i - j|) \leq C b(|j|)$. Then the result will easily follow, with the choices $p_i = b(|i|)$ and $\phi = C c \varphi$. We define b recursively, by $b(0) = a(0)$ and $b(k + 1) = \max\{a(k + 1), [(k + 1)/(k + 2)]^{2d} b(k)\}$. Using that a is nonincreasing, we easily check that b is nonincreasing. We next check that $\sum_{i \in \mathbb{Z}^d} b(|i|) < \infty$, i.e. that $\sum_{k \geq 0} k^{d-1} b(k) < \infty$, knowing by assumption that $\sum_{k \geq 0} k^{d-1} a(k) < \infty$. We have, for $k \geq 0$,

$$\begin{aligned} b(k + 1) - a(k + 1) &= \left(\left[\frac{k + 1}{k + 2} \right]^{2d} b(k) - a(k + 1) \right)_+ \\ &\leq \left[\frac{k + 1}{k + 2} \right]^{2d} (b(k) - a(k))_+ + \left(\left[\frac{k + 1}{k + 2} \right]^{2d} a(k) - a(k + 1) \right)_+ \\ &\leq \left[\frac{k + 1}{k + 2} \right]^{2d} (b(k) - a(k)) + (a(k) - a(k + 1)). \end{aligned}$$

Recalling that $b(0) = a(0)$, one gets $b(k) - a(k) \leq \sum_{\ell=1}^k (a(\ell - 1) - a(\ell)) \left[\frac{\ell + 1}{k + 1} \right]^{2d}$ by iteration. Hence

$$\begin{aligned} \sum_{k \geq 1} k^{d-1} (b(k) - a(k)) &\leq \sum_{k \geq 1} k^{d-1} \sum_{\ell=1}^k (a(\ell - 1) - a(\ell)) \left[\frac{\ell + 1}{k + 1} \right]^{2d} \\ &= \sum_{\ell \geq 1} (a(\ell - 1) - a(\ell)) (\ell + 1)^{2d} \sum_{k \geq \ell} k^{d-1} (k + 1)^{-2d} \\ &\leq C \sum_{\ell \geq 1} (a(\ell - 1) - a(\ell)) \ell^d. \end{aligned}$$

This last quantity is nothing but $C \sum_{\ell \geq 1} a(\ell)[(\ell + 1)^d - \ell^d] \leq C \sum_{\ell \geq 1} a(\ell)\ell^{d-1} < \infty$. We have thus checked that $\sum_{k \geq 0} k^{d-1}b(k) = \sum_{k \geq 1} a(k)k^{d-1} + \sum_{k \geq 1} k^{d-1}(b(k) - a(k)) < \infty$.

We finally prove that for all $j \in \mathbb{Z}^d$, $\sum_{i \in \mathbb{Z}^d} b(|i|)a(|i - j|) \leq Cb(|j|)$. First, we claim that there is C such that $b(k) \leq Cb(2k)$ for all $k \geq 0$. This is easily checked, iterating the inequality $b(k) \leq [(k + 2)/(k + 1)]^{2d}b(k + 1)$. Next we write, using that a and b are nonincreasing,

$$\begin{aligned} \sum_{i \in \mathbb{Z}^d} b(|i|)a(|i - j|) &\leq \sum_{|i| < |j|/2} b(|i|)a(|i - j|) + \sum_{|i| \geq |j|/2} b(|i|)a(|i - j|) \\ &\leq a(|j|/2) \sum_{i \in \mathbb{Z}^d} b(|i|) + b(|j|/2) \sum_{i \in \mathbb{Z}^d} a(|i - j|) \\ &\leq Ca(|j|/2) + Cb(|j|/2). \end{aligned}$$

By definition of b , we have $a(|j|/2) \leq b(|j|/2)$. And we have just seen that $b(|j|/2) \leq Cb(|j|)$. We finally have checked that $\sum_{i \in \mathbb{Z}^d} b(|i|)a(|i - j|) \leq Cb(|j|)$ as desired. \square

Our well-posedness result is the following.

Theorem 6. *Under Assumption 4, there exists a pathwise unique Hawkes process $(Z_t^i)_{i \in \mathcal{S}, t \geq 0}$ such that $\sum_{i \in \mathcal{S}} p_i \mathbb{E}[Z_t^i] < \infty$ for all $t \geq 0$.*

Observe that this result is not completely obvious in the case of an infinite graph. In some sense, we have to check that the interaction does not come from infinity. Let us insist on the fact that, even in simple situations, a *graphical construction* is not possible: consider *e.g.* the case of \mathbb{Z} endowed with the set of edges $\mathcal{E} = \{(i, j) : |i - j| = 0 \text{ or } 1\}$, assume that $h_i(x) = 1 + x$ for all $i \in \mathcal{S}$ and that $\varphi_{ij} = 1$ for all $(i, j) \in \mathcal{E}$. Then one easily gets convinced that we cannot determine the values of $(Z_t^0)_{t \in [0, T]}$ by observing the Poisson measures π^i in a (random) finite box.

As a second comment, let us mention that we believe it is not possible, or at least quite difficult, to obtain the full uniqueness, *i.e.* uniqueness outside the class of processes satisfying $\sum_{i \in \mathcal{S}} p_i \mathbb{E}[Z_t^i] < \infty$ (or something similar). Indeed, consider again the case of \mathbb{Z} endowed with $\mathcal{E} = \{(i, j) : |i - j| = 0 \text{ or } 1\}$, assume that $h_i(x) = 1 + x$ for all $i \in \mathcal{S}$ and that $\varphi_{ji} = 1$ for all $(i, j) \in \mathcal{E}$. One easily checks that for $(Z_t^i)_{i \in \mathcal{S}, t \geq 0}$ a Hawkes process, for $m_t^i = \mathbb{E}[Z_t^i]$, it holds that $m_t^i = t + \int_0^t (m_s^{i-1} + m_s^i + m_s^{i+1}) ds$ for every i . This infinite system of equations is of course closely related to the heat equation $\partial_t u(t, x) = 1 + \partial_{xx} u(t, x)$ on $[0, \infty) \times \mathbb{R}$ and with initial condition $u(0, x) = 0$. As is well-known, uniqueness for this equation fails to hold true without imposing some growth conditions as $|x| \rightarrow \infty$. See *e.g.* Tychonov's counterexample of uniqueness, which can be found in John [24, Chapter 7].

To compare Theorem 6 with the results of Massoulié [28], let us consider a very simple situation. Let $\mathcal{S} = \mathbb{Z}^d$ be endowed with $\mathcal{E} = \{(i, j) : |i - j| = 0 \text{ or } 1\}$, let $h_i(x) = 1 + x$ for all $i \in \mathcal{S}$ and $\varphi_{ij}(t) = \varphi(t) > 0$ for all $(i, j) \in \mathcal{E}$. Then Theorem 6 applies as soon as $\varphi \in L_{loc}^1([0, \infty))$, see Remark 5-(ii). Theorem 1 of [28] does not apply for two reasons: $\sup_{i \in \mathcal{S}} h_i$ is not bounded and, more important, it does not hold true that $\sum_{(i, j) \in \mathcal{E}} \varphi_{ij} \in L_{loc}^1([0, \infty))$ since $\sum_{(i, j) \in \mathcal{E}} \varphi_{ij}(t) = +\infty$ for all $t \geq 0$. On the contrary, [28, Theorem 2] applies, but only in the subcritical case where $(2d + 1) \int_0^\infty \varphi(t) dt < 1$.

Proof. We first prove uniqueness. Let thus $(Z_t^i)_{i \in \mathcal{S}, t \geq 0}$ and $(\tilde{Z}_t^i)_{i \in \mathcal{S}, t \geq 0}$ be two solutions to (1) satisfying the required condition. Set

$$\Delta_t^i = \int_0^t |d(Z_s^i - \tilde{Z}_s^i)| \quad \text{for } i \in \mathcal{S}, t \geq 0.$$

In other words, Δ_t^i is the total variation norm of the signed measure $d(Z_s^i - \tilde{Z}_s^i)$ on $[0, t]$. We also put $\delta_t^i = \mathbb{E}[\Delta_t^i]$ and first prove that

$$(2) \quad \delta_t^i \leq c_i \int_0^t \sum_{j \rightarrow i} |\varphi_{ji}(t-s)| \delta_s^j ds.$$

We have

$$\Delta_t^i = \int_0^t \int_0^\infty \left| \mathbf{1}_{\{z \leq h_i(\sum_{j \rightarrow i} \int_0^{s-} \varphi_{ji}(s-u) dZ_u^j)\}} - \mathbf{1}_{\{z \leq h_i(\sum_{j \rightarrow i} \int_0^{s-} \varphi_{ji}(s-u) d\tilde{Z}_u^j)\}} \right| \pi^i(ds dz).$$

Taking expectations, we deduce that

$$(3) \quad \begin{aligned} \delta_t^i &= \int_0^t \mathbb{E} \left[\left| h_i \left(\sum_{j \rightarrow i} \int_0^{s-} \varphi_{ji}(s-u) dZ_u^j \right) - h_i \left(\sum_{j \rightarrow i} \int_0^{s-} \varphi_{ji}(s-u) d\tilde{Z}_u^j \right) \right| \right] ds \\ &\leq c_i \sum_{j \rightarrow i} \mathbb{E} \left[\int_0^t \int_0^{s-} |\varphi_{ji}(s-u)| d\Delta_u^j ds \right] \end{aligned}$$

by Assumption 4-(a). Using Lemma 22, we see that

$$\int_0^t ds \int_0^{s-} |\varphi_{ji}(s-u)| d\Delta_u^j = \int_0^t |\varphi_{ji}(t-u)| \Delta_u^j du$$

which, plugged into (3), yields (2).

Set $\delta_t = \sum_{i \in \mathcal{S}} p_i \delta_t^i$, where the weights p_i were introduced in Assumption 4. By assumption, δ_t is well-defined and finite. We infer by (2) that

$$\delta_t \leq \int_0^t \sum_{i \in \mathcal{S}} p_i c_i \sum_{j \rightarrow i} |\varphi_{ji}(t-s)| \delta_s^j ds.$$

By Assumption 4-(c),

$$\delta_t \leq \int_0^t \sum_{j \in \mathcal{S}} \delta_s^j \sum_{i, (j,i) \in \mathcal{E}} c_i p_i |\varphi_{ji}(t-s)| ds \leq \int_0^t \sum_{j \in \mathcal{S}} p_j \delta_s^j \phi(t-s) ds = \int_0^t \phi(t-s) \delta_s ds.$$

Lemma 23-(i) thus implies that $\delta_t = 0$ identically, from which uniqueness follows.

We now quickly prove existence by a Picard iteration. Let $Z_t^{i,0} = 0$ and, for $n \geq 0$,

$$(4) \quad Z_t^{i,n+1} = \int_0^t \int_0^\infty \mathbf{1}_{\{z \leq h_i(\sum_{j \rightarrow i} \int_0^{s-} \varphi_{ji}(s-u) dZ_u^{j,n})\}} \pi^i(ds dz).$$

We define $\delta_t^{i,n} = \mathbb{E}[\int_0^t |dZ_s^{i,n+1} - dZ_s^{i,n}|]$ and $\delta_t^n = \sum_{i \in \mathcal{S}} p_i \delta_t^{i,n}$. As in the proof of uniqueness, we obtain, for $n \geq 0$,

$$(5) \quad \delta_t^{n+1} \leq \int_0^t \phi(t-s) \delta_s^n ds.$$

Next, we put $m_t^{i,n} = \mathbb{E}[Z_t^{i,n}]$. By Assumption 4-(a), $h_i(x) \leq h_i(0) + c_i|x|$, whence

$$m_t^{i,n+1} \leq \mathbb{E} \left[\int_0^t \left(h_i(0) + c_i \sum_{j \rightarrow i} \int_0^{s-} |\varphi_{ji}(s-u)| dZ_u^{j,n} \right) ds \right] \leq \int_0^t \left(h_i(0) + c_i \sum_{j \rightarrow i} |\varphi_{ji}(t-s)| m_s^{j,n} \right) ds,$$

where we used that, by Lemma 22, $\int_0^t \int_0^{s-} |\varphi_{ji}(s-u)| dZ_u^{j,n} ds = \int_0^t |\varphi_{ji}(t-u)| Z_u^{j,n} du$. Setting $u_t^n = \sum_{i \in \mathcal{S}} p_i m_t^{i,n}$ and using Assumption 4-(b)-(c),

$$(6) \quad u_t^{n+1} \leq t \sum_{i \in \mathcal{S}} h_i(0) p_i + \int_0^t \sum_{i \in \mathcal{S}} p_i c_i \sum_{j \rightarrow i} |\varphi_{ji}(s-u)| m_s^{j,n} ds \leq Ct + \int_0^t \phi(t-s) u_s^n ds.$$

Since $u_t^0 = 0$ and ϕ is locally integrable, we easily check by induction that u^n is locally bounded for all $n \geq 0$. Consequently, δ^n is also locally bounded for all $n \geq 0$. Lemma 23-(ii) implies that for all $T \geq 0$, $\sum_{n \geq 1} \delta_T^n < \infty$. This classically implies that the Picard sequence is Cauchy and thus converges: there exists a family $(Z_t^i)_{i \in \mathcal{S}, t \geq 0}$ of càdlàg nonnegative adapted processes such that for all $T \geq 0$, $\lim_n \sum_{i \in \mathcal{S}} p_i \mathbb{E}[\int_0^T |dZ_s^i - dZ_s^{i,n}|] = 0$. It is then not hard to pass to the limit in (4) to deduce that $(Z_t^i)_{i \in \mathcal{S}, t \geq 0}$ solves (1). Finally, Lemma 23-(iii) implies that $\sup_n u_t^n < \infty$ for all $t \geq 0$, from which $\sum_{i \in \mathcal{S}} p_i \mathbb{E}[Z_t^i] < \infty$ as desired. \square

3. MEAN-FIELD LIMIT

In this section, we work in the following setting.

Assumption 7. Let $h : \mathbb{R} \mapsto [0, \infty)$ be such that $|h|_{lip} = \sup_{x \neq y} |x - y|^{-1} |h(x) - h(y)| < \infty$ and let $\varphi = [0, \infty) \mapsto \mathbb{R}$ be a locally square integrable function.

For each $N \geq 1$, we consider the complete graph \mathbb{G}_N with vertices $\mathcal{S}_N = \{1, \dots, N\}$ and edges $\mathcal{E}_N = \{(i, j) : i, j \in \mathcal{S}_N\}$, i.e. all pairs of points in \mathcal{S}_N are connected. We put $h_i^N = h$ for all $i \in \mathcal{S}_N$ and $\varphi_{ji}^N = N^{-1} \varphi$ for $(i, j) \in \mathcal{E}_N$.

Under Assumption 7, the triplet $(\mathbb{G}_N, \varphi^N, \mathbf{h}^N)$ satisfies Assumption 4 (the graph \mathbb{G}_N is finite) for each $N \geq 1$. Therefore, a Hawkes process $(Z_t^{N,1}, \dots, Z_t^{N,N})_{t \geq 0}$ with parameters $(\mathbb{G}_N, \varphi^N, \mathbf{h}^N)$ is uniquely defined by Theorem 6.

Introduce the *limit* equation

$$(7) \quad \bar{Z}_t = \int_0^t \int_0^\infty \mathbf{1}_{\{z \leq h(\int_0^s \varphi(s-u) d\mathbb{E}[\bar{Z}_u])\}} \pi(ds dz), \quad \text{for every } t \geq 0,$$

where $\pi(ds dz)$ is a Poisson measure on $[0, \infty) \times [0, \infty)$ with intensity measure $dsdz$. Also, $d\mathbb{E}[\bar{Z}_u]$ is the measure on $[0, \infty)$ associated to the (necessarily) non-decreasing function $u \mapsto \mathbb{E}[\bar{Z}_u]$. Note that a solution $\bar{Z} = (\bar{Z}_t)_{t \geq 0}$, if it exists, is an inhomogeneous Poisson process on $[0, \infty)$ with intensity $\lambda_t = h(\int_0^t \varphi(t-u) d\mathbb{E}[\bar{Z}_u])$.

3.1. Propagation of chaos. The main result of this section reads as follows. We denote by $\mathbb{D}([0, \infty), \mathbb{R})$ the set of càdlàg \mathbb{R} -valued functions on $[0, \infty)$ and by $\mathcal{P}(\mathbb{D}([0, \infty), \mathbb{R}))$ the set of probability measures on $\mathbb{D}([0, \infty), \mathbb{R})$.

Theorem 8. *Work under Assumption 7.*

(i) *There is a pathwise unique solution $(\bar{Z}_t)_{t \geq 0}$ to (7) such that $(\mathbb{E}[\bar{Z}_t])_{t \geq 0}$ is locally bounded.*

(ii) It is possible to build simultaneously the Hawkes process $(Z_t^{N,1}, \dots, Z_t^{N,N})_{t \geq 0}$ with parameters $(\mathbb{G}_N, \boldsymbol{\varphi}^N, \mathbf{h}^N)$ and an i.i.d. family $(\bar{Z}_t^i)_{t \geq 0, i=1, \dots, N}$ of solutions to (7) in such a way that for all $T > 0$, all $i = 1, \dots, N$,

$$\mathbb{E} \left[\sup_{[0, T]} |Z_t^{N,i} - \bar{Z}_t^i| \right] \leq C_T N^{-1/2},$$

the constant C_T depending only on h , φ and T (see Remark 9 below for some bounds of C_T in a few situations).

(iii) Consequently, we have the mean-field approximation

$$\frac{1}{N} \sum_{i=1}^N \delta_{(Z_t^{N,i})_{t \geq 0}} \longrightarrow \mathcal{L}((\bar{Z}_t)_{t \geq 0}) \text{ in probability, as } N \rightarrow \infty,$$

where $\mathcal{P}(\mathbb{D}([0, \infty), \mathbb{R}))$ is endowed with the weak convergence topology associated with the topology (on $\mathbb{D}([0, \infty), \mathbb{R}))$ of the uniform convergence on compact time intervals.

Proof. For $(\bar{Z}_t)_{t \geq 0}$ a solution to (7), the equation satisfied by $m_t = \mathbb{E}[\bar{Z}_t]$ writes

$$(8) \quad m_t = \int_0^t h \left(\int_0^s \varphi(s-u) dm_u \right) ds \text{ for every } t \geq 0.$$

By Lemma 24, we know that this equation has a unique non-decreasing locally bounded solution, which furthermore is of class C^1 on $[0, \infty)$. We now split the proof in several steps.

Step 1. Here we prove the well-posedness of (7). For $(\bar{Z}_t)_{t \geq 0}$ a solution to (7), its expectation $m_t = \mathbb{E}[\bar{Z}_t]$ solves (8) and is thus uniquely defined. Thus the right hand side of (7) is uniquely determined, which proves uniqueness. For the existence, consider m the unique solution to (8) and put $\bar{Z}_t = \int_0^t \int_0^\infty \mathbf{1}_{\{z \leq h(\int_0^s \varphi(s-u) dm_u)\}} \pi(ds dz)$. We thus only have to prove that $\mathbb{E}[\bar{Z}_t] = m_t$. But $\mathbb{E}[\bar{Z}_t] = \int_0^t h(\int_0^s \varphi(s-u) dm_u) ds$, which is nothing but m_t since m solves (8).

Step 2. We next introduce a suitable coupling. Let $(\pi^i(ds dz))_{i \geq 1}$ be an i.i.d. family of Poisson measures with common intensity measure $dsdz$ on $[0, \infty) \times [0, \infty)$. For each $N \geq 1$, we consider the Hawkes process $(Z_t^{N,1}, \dots, Z_t^{N,N})_{t \geq 0}$

$$Z_t^{N,i} = \int_0^t \int_0^\infty \mathbf{1}_{\{z \leq h(N^{-1} \sum_{j=1}^N \int_0^{s-} \varphi(s-u) dZ_u^{N,j})\}} \pi^i(ds dz).$$

Next, still denoting by m the unique solution to (8), we put, for every $i \geq 1$,

$$\bar{Z}_t^i = \int_0^t \int_0^\infty \mathbf{1}_{\{z \leq h(\int_0^{s-} \varphi(s-u) dm_u)\}} \pi^i(ds dz).$$

Clearly, $((\bar{Z}_t^i)_{t \geq 0})_{i \geq 1}$ is an i.i.d. family of solutions to (7).

Step 3. Here we introduce $\Delta_N^i(t) = \int_0^t |d(\bar{Z}_u^i - Z_u^{N,i})|$ and $\delta_N(t) = \mathbb{E}[\Delta_N^i(t)]$, which obviously does not depend on i (by exchangeability). Observe that

$$(9) \quad \sup_{[0, t]} |\bar{Z}_u^i - Z_u^{N,i}| \leq \Delta_N^i(t), \quad \text{whence} \quad \mathbb{E} \left[\sup_{[0, t]} |\bar{Z}_u^i - Z_u^{N,i}| \right] \leq \delta_N(t).$$

The first inequality follows from the fact that $|\bar{Z}_u^i - Z_u^{N,i}| \leq |\int_0^u d(\bar{Z}_r^i - Z_r^{N,i})| \leq \int_0^u |d(\bar{Z}_r^i - Z_r^{N,i})|$. We show in this step that for all $t > 0$,

$$(10) \quad \delta_N(t) \leq |h|_{lip} N^{-1/2} \int_0^t \left(\int_0^s \varphi^2(s-u) dm_u \right)^{1/2} ds + |h|_{lip} \int_0^t |\varphi(t-s)| \delta_N(s) ds.$$

First, $\Delta_N^1(t)$ equals

$$\int_0^t \int_0^\infty \left| \mathbf{1}_{\{z \leq h(N^{-1} \sum_{j=1}^N \int_0^{s-} \varphi(s-u) dZ_u^{N,j})\}} - \mathbf{1}_{\{z \leq h(\int_0^{s-} \varphi(s-u) dm_u)\}} \right| \pi^i(ds dz).$$

Taking expectations, we find

$$\delta_N(t) = \int_0^t \mathbb{E} \left[\left| h \left(\int_0^s \varphi(s-u) dm_u \right) - h \left(N^{-1} \sum_{j=1}^N \int_0^s \varphi(s-u) dZ_u^{N,j} \right) \right| \right] ds,$$

whence

$$(11) \quad \begin{aligned} \delta_N(t) &\leq |h|_{lip} \int_0^t \mathbb{E} \left[\left| \int_0^s \varphi(s-u) dm_u - N^{-1} \sum_{j=1}^N \int_0^s \varphi(s-u) d\bar{Z}_u^j \right| \right] ds \\ &\quad + |h|_{lip} \int_0^t \mathbb{E} \left[\left| N^{-1} \sum_{j=1}^N \int_0^s \varphi(s-u) d[\bar{Z}_u^j - Z_u^{N,j}] \right| \right] ds \\ &= |h|_{lip} (A + B). \end{aligned}$$

Using exchangeability and Lemma 22,

$$(12) \quad B \leq \int_0^t \mathbb{E} \left[\int_0^s |\varphi(s-u)| d\Delta_N^1(u) \right] ds = \int_0^t |\varphi(t-u)| \delta_N(u) du.$$

Next, we use that $X_s^j = \int_0^s \varphi(s-u) d\bar{Z}_u^j$ are i.i.d. with mean $\int_0^s \varphi(s-u) dm_u$, whence

$$(13) \quad A \leq N^{-1/2} \int_0^t (\text{Var } X_s^1)^{1/2} ds.$$

But it holds that

$$X_s^1 = \int_0^s \int_0^\infty \mathbf{1}_{\{z \leq h(\int_0^u \varphi(u-r) dm_r)\}} \varphi(s-u) \pi^1(du dz).$$

Since the integrand is deterministic, denoting by $\tilde{\pi}^1$ the compensated Poisson measure,

$$X_s^1 - \mathbb{E}[X_s^1] = \int_0^s \int_0^\infty \mathbf{1}_{\{z \leq h(\int_0^u \varphi(u-r) dm_r)\}} \varphi(s-u) \tilde{\pi}^1(du dz).$$

Recalling Assumption 7, we find

$$(14) \quad \text{Var } X_s^1 = \int_0^s \varphi^2(s-u) h \left(\int_0^u \varphi(u-r) dm_r \right) du = \int_0^s \varphi^2(s-u) dm_u.$$

We used (8) for the last equality. Gathering (11), (12), (13) and (14) completes the step.

Step 4. Here we conclude that for all $T \geq 0$, $\sup_{[0,T]} \delta_N(t) \leq C_T N^{-1/2}$. This will end the proof of (ii) by (9). This is not hard: it suffices to start from (10), to apply Lemma 23-(i) and to observe that $\int_0^t \left(\int_0^s \varphi^2(s-u) dm_u \right)^{1/2} ds$ is locally bounded (which follows from the assumption that φ is locally square integrable and the fact that m is C^1 on $[0, \infty)$).

Step 5. Finally, (iii) follows from (ii): by Sznitman [37, Proposition 2.2], it suffices to check that for each fixed $\ell \geq 1$, $((Z_t^{N,1})_{t \geq 0}, \dots, (Z_t^{N,\ell})_{t \geq 0})$ goes in law, as $N \rightarrow \infty$, to ℓ independent copies of $(\bar{Z}_t)_{t \geq 0}$ (for the uniform topology on compact time intervals). This clearly follows from (ii). \square

We now want to show that the constant C_T we get can be quite satisfactory.

Remark 9. *Work under Assumption 7.*

(a) Assume that $|h|_{\text{lip}} \int_0^\infty |\varphi(s)| ds < 1$ (subcritical case) and that $\int_0^\infty \varphi^2(s) ds < \infty$. Then (ii) of Theorem 8 holds with $C_T = CT$, for some constant $C > 0$. This is a satisfactory slow growth.

(b) Assume that $h(x) = \mu + x$ for some $\mu > 0$ and that $\varphi(t) = ae^{-bt}$ for some $a > b > 0$ (if $a < b$, then point (a) applies). Then $m_t = \mathbb{E}[\bar{Z}_t] \sim \mu a(a-b)^{-2} e^{(a-b)t}$ as $t \rightarrow \infty$ and (ii) of Theorem 8 holds with $C_T = Ce^{(a-b)T}$, for some constant $C > 0$. This is again quite satisfactory: the error is of order $N^{-1/2} m_T$.

Proof. We start with (a). Using the notation of the previous proof, it suffices (see (9)) to show that $\delta_N(T) \leq CTN^{-1/2}$. Setting $\Lambda = |h|_{\text{lip}} \int_0^\infty |\varphi(s)| ds < 1$, starting from (10) and observing that δ_N is non-decreasing, we find $\delta_N(t) \leq |h|_{\text{lip}} N^{-1/2} \int_0^t (\int_0^s \varphi^2(s-u) dm_u)^{1/2} ds + \Lambda \delta_N(t)$, whence $\delta_N(t) \leq CN^{-1/2} \int_0^t (\int_0^s \varphi^2(s-u) dm_u)^{1/2} ds$. We thus only have to check that $\int_0^s \varphi^2(s-u) dm_u$ is bounded on $[0, \infty)$. Since $\int_0^\infty \varphi^2(s) ds < \infty$, it suffices to prove that m' is bounded on $[0, \infty)$. But $m'_t = h(\int_0^t \varphi(t-u) m'_u du) \leq h(0) + |h|_{\text{lip}} \int_0^t |\varphi(t-u)| m'_u du$, whence $\sup_{[0,T]} m'_t \leq h(0) + \Lambda \sup_{[0,T]} m'_t$ and thus $\sup_{[0,T]} m'_t \leq h(0)/(1-\Lambda)$ for any $T > 0$.

We next check (b). First, (8) rewrites $m_t = \mu t + a \int_0^t \int_0^s e^{-b(s-u)} dm_u ds$, with unique solution

$$m_t = \frac{-\mu b t}{a-b} + \frac{\mu a (e^{(a-b)t} - 1)}{(a-b)^2} \sim \frac{\mu a}{(a-b)^2} e^{(a-b)t}.$$

Next, using (10) and the explicit expressions of h , φ and m , we find

$$\begin{aligned} \delta_N(t) &\leq N^{-1/2} \int_0^t \left(\int_0^s \varphi^2(s-u) dm_u \right)^{1/2} ds + \int_0^t \varphi(t-s) \delta_N(s) ds \\ &\leq CN^{-1/2} e^{(a-b)t/2} + a \int_0^t e^{-b(t-s)} \delta_N(s) ds. \end{aligned}$$

Setting $u_N(t) = \delta_N(t) e^{bt}$, we get $u_N(t) \leq CN^{-1/2} e^{(a+b)t/2} + a \int_0^t u_N(s) ds$. By Grönwall's lemma, $u_N(t) \leq CN^{-1/2} e^{(a+b)t/2} + a \int_0^t CN^{-1/2} e^{(a+b)s/2} e^{a(t-s)} ds$. One easily deduces, since $a > b$, that $u_N(t) \leq CN^{-1/2} e^{at}$ so that $\delta_N(t) \leq CN^{-1/2} e^{(a-b)t}$. The use of (9) ends the proof. \square

3.2. Large time behaviour. We now address the important problem of the large time behaviour. Since the solution $(\bar{Z}_t)_{t \geq 0}$ to (7) is nothing but an inhomogeneous Poisson process, its large-time behaviour is easily and precisely described, provided we have sufficiently information on the solution to (8). The question is thus: can we use the large time estimates of the mean-field limit to describe the large-time behaviour of the true Hawkes process with a large number of particles? To fix the ideas, we consider the linear case. For A a symmetric nonnegative matrix, we denote by $\mathcal{N}(0, A)$ the centered Gaussian distribution with covariance matrix A .

We treat separately the subcritical and supercritical cases.

Theorem 10. *Work under Assumption 7 with φ nonnegative and $h(x) = \mu + x$ for some $\mu > 0$. Assume also that $\Lambda = \int_0^\infty \varphi(s) ds < 1$. For each $N \geq 1$, consider the Hawkes process*

$(Z_t^{N,1}, \dots, Z_t^{N,N})_{t \geq 0}$ with parameters $(\mathbb{G}_N, \boldsymbol{\varphi}^N, \mathbf{h}^N)$. Consider also the unique solution $(m_t)_{t \geq 0}$ to (8).

1. We have $m_t \sim a_0 t$ as $t \rightarrow \infty$, where $a_0 = \mu/(1 - \Lambda)$.

2. For any fixed $i \geq 1$, $Z_t^{N,i}/m_t$ tends to 1 in probability as $t \rightarrow \infty$, uniformly in N . More precisely, $\mathbb{E}[|Z_t^{N,i}/m_t - 1|] \leq C m_t^{-1/2}$ for some constant C .

3. For any fixed $\ell \geq 1$ $(m_t^{1/2}(Z_t^{N,i}/m_t - 1))_{i=1, \dots, \ell}$ goes in law to $\mathcal{N}(0, I_\ell)$ as $(t, N) \rightarrow (\infty, \infty)$ (without condition on the regime).

Point 2. is of course related to the classical law of large numbers for multivariate Hawkes processes, see e.g. Brémaud-Massoulié [8] or [3]. See also Zhu [43, 42] for some large deviations results. What we prove here is that this law of large numbers is uniform in N .

Theorem 11. *Work under Assumption 7 with φ nonnegative and $h(x) = \mu + x$ for some $\mu > 0$. Assume also that $\Lambda = \int_0^\infty \varphi(s) ds \in (1, \infty]$. Assume finally that $t \mapsto \int_0^t |d\varphi(s)|$ has at most polynomial growth. For each $N \geq 1$, consider the Hawkes process $(Z_t^{N,1}, \dots, Z_t^{N,N})_{t \geq 0}$ with parameters $(\mathbb{G}_N, \boldsymbol{\varphi}^N, \mathbf{h}^N)$. Consider also the unique solution $(m_t)_{t \geq 0}$ to (8).*

1. We have $m_t \sim a_0 e^{\alpha_0 t}$ as $t \rightarrow \infty$, where $\alpha_0 > 0$ is determined by $\mathcal{L}_\varphi(\alpha_0) = 1$ and where $a_0 = \mu \alpha_0^{-2} (\int_0^\infty t \varphi(t) e^{-\alpha_0 t} dt)^{-1}$.

2. For any fixed $i \geq 1$, $Z_t^{N,i}/m_t$ tends to 1 in probability as $(t, N) \rightarrow (\infty, \infty)$. More precisely, there is a constant C such that $\mathbb{E}[|Z_t^{N,i}/m_t - 1|] \leq m_t^{-1/2} + C N^{-1/2} (1 + m_t^{-1})$.

3. For any fixed $\ell \geq 1$,

(i) $(m_t^{1/2}(Z_t^{N,i}/m_t - 1))_{i=1, \dots, \ell}$ goes in law to $\mathcal{N}(0, I_\ell)$ if $t \rightarrow \infty$ and $N \rightarrow \infty$ with $m_t/N \rightarrow 0$;

(ii) $(N^{1/2}(Z_t^{N,i}/m_t - 1))_{i=1, \dots, \ell}$ goes in law to (X, \dots, X) , if $t \rightarrow \infty$ and $N \rightarrow \infty$ with $m_t/N \rightarrow \infty$. Here X is a $\mathcal{N}(0, \sigma^2)$ -distributed random variable, where $\sigma^2 = \alpha_0^2 \mu^{-2} \int_0^\infty e^{-2\alpha_0 s} m'_s ds$.

Let us summarize. At first order (law of large numbers), the mean-field approximation is always good for large times. At second order (central limit theorem), the mean field approximation is always good for large times in the subcritical case, but fails to be relevant for too large times (depending on N) in the supercritical case: the independence property breaks down.

In the supercritical case, we have the technical condition that $t \mapsto \int_0^t |d\varphi(s)|$ has at most polynomial growth. This is useful to have some precise estimates of the solution m to (8). This is, e.g. always satisfied when φ is bounded and non-increasing, as is often the case in applications. It is slightly restrictive however, since it forces $\varphi(0)$ to be finite.

It should be possible to study also the *critical* case, but then the situation is more intricate: many regimes might arise. With a little more work, we could also study, in the supercritical case, the regime where $m_t/N \rightarrow x \in (0, \infty)$.

In order to prove Theorems 10 and 11, we will use the following central limit theorem for martingales. For two càdlàg martingales M and N , we denote by $[M, N]$ the quadratic covariation defined by $[M, N]_t = M_t N_t - \int_0^t M_{s-} dN_s - \int_0^t N_{s-} dM_s$. When M and N are purely discontinuous, it holds that $[M, N]_t = \sum_{s \leq t} \Delta M_s \Delta N_s$, see Jacod-Shiryaev [23, Chapter I, §4e].

Lemma 12. *Let $\ell \geq 1$ be fixed. For $N \geq 1$, consider a family $(M_t^{N,1}, \dots, M_t^{N,\ell})_{t \geq 0}$ of ℓ -dimensional local martingales satisfying $M_0^{N,i} = 0$. Assume that all their jumps are uniformly*

bounded and that $[M^{N,i}, M^{N,j}]_t = 0$ for every $N \geq 1$, $i \neq j$ and $t \geq 0$. Assume also that there is a continuous increasing function $(v_t)_{t \geq 0} : [0, \infty) \mapsto [0, \infty)$ such that for all $i = 1, \dots, \ell$, $\lim_{(t,N) \rightarrow (\infty, \infty)} v_t^{-2} [M^{N,i}, M^{N,i}]_t = 1$ in probability. In the case where $v_\infty = \lim_{t \rightarrow \infty} v_t < \infty$, assume moreover that for all $i = 1, \dots, \ell$, all $t_0 > 0$, uniformly in $t \geq t_0$, $\lim_{N \rightarrow \infty} [M^{N,i}, M^{N,i}]_t = v_t^2$ in probability.

Then $v_t^{-1}(M_t^{N,1}, \dots, M_t^{N,\ell})$ converges in law to the Gaussian distribution $\mathcal{N}(0, I_\ell)$ as $(t, N) \rightarrow (\infty, \infty)$, where I_ℓ is the $\ell \times \ell$ identity matrix.

Proof. Let $(t_N)_{N \geq 1}$ be a sequence of positive numbers such that $t_N \rightarrow \infty$. We want to prove that $v_{t_N}^{-1}(M_{t_N}^{N,1}, \dots, M_{t_N}^{N,\ell})$ converges in law to $\mathcal{N}(0, I_\ell)$. For all $u \in [0, 1]$, set

$$\tau_u^N = \inf\{t \geq 0 : v_t^2 \geq u v_{t_N}^2\}.$$

Since v is increasing and continuous, τ^N is also continuous and increasing for each N . We also clearly have $v_{\tau_u^N}^2 = u v_{t_N}^2$ for all $u \in [0, 1]$ and $\tau_1^N = t_N$. Finally, for each $u > 0$ fixed, the sequence τ_u^N is increasing.

For all $u \in (0, 1]$, $\lim_N v_{\tau_u^N}^{-2} [M^{N,i}, M^{N,i}]_{\tau_u^N} = 1$ in probability. Indeed, in the case $v_\infty = \infty$, this follows from the facts that $\lim_N \tau_u^N = \infty$ and $\lim_{(t,N) \rightarrow (\infty, \infty)} v_t^{-2} [M^{N,i}, M^{N,i}]_t = 1$. When $v_\infty < \infty$, the additional assumption (uniformity in $t \geq t_0$ of the convergence as $N \rightarrow \infty$) clearly suffices, since the sequence τ_u^N is increasing and thus bounded from below.

We define the martingales $(L_u^{N,i})_{u \in [0,1]}$ by $L_u^{N,i} = v_{\tau_u^N}^{-1} M_{\tau_u^N}^{N,i}$. All their jumps are uniformly bounded (because those of $M^{N,i}$ are assumed to be uniformly bounded and because $\sup_N v_{t_N}^{-1} < \infty$ since v is increasing). We also have $[L^{N,i}, L^{N,j}]_u = 0$ for all $i \neq j$, all $u \in [0, 1]$. Furthermore, using that $v_{\tau_u^N}^2 = u v_{t_N}^2$,

$$[L^{N,i}, L^{N,i}]_u = \frac{[M^{N,i}, M^{N,i}]_{\tau_u^N}}{v_{t_N}^2} = \frac{[M^{N,i}, M^{N,i}]_{\tau_u^N}}{v_{\tau_u^N}^2} u \rightarrow u$$

in probability. Therefore, according to Jacod-Shiryaev [23] (Theorem VIII-3.11), the process $(L_u^{N,1}, \dots, L_u^{N,\ell})_{u \in [0,1]}$ converges in law to $(B_u^1, \dots, B_u^\ell)_{u \in [0,1]}$ where the B^i are independent standard Brownian motions. In particular, $(L_1^{N,1}, \dots, L_1^{N,\ell})$ goes in law to $\mathcal{N}(0, I_\ell)$. To conclude the proof, it thus suffices to observe that $L_1^{N,i} = v_{t_N}^{-1} M_{\tau_1^N}^{N,i} = v_{t_N}^{-1} M_{t_N}^{N,i}$. \square

We can now give the

Proof of Theorem 10. In the present (linear) case, we can rewrite (8) as $m_t = \int_0^t (\mu + \int_0^s \varphi(s-u) dm_u) ds = \mu t + \int_0^t \varphi(t-s) m_s ds$ by Lemma 22. This equation is studied in details in Lemma 25: recalling that $\Lambda = \int_0^\infty \varphi(s) ds < 1$, we have $m_t' \sim a_0$ and $m_t \sim a_0 t$ as $t \rightarrow \infty$, where $a_0 = \mu/(1-\Lambda)$, which proves point (i). The proof is now divided in several steps. Step 1 will also be used in the supercritical case.

Step 1. Recall that, for some i.i.d. family $(\pi^i(ds dz))_{i \geq 1}$ of Poisson measures on $[0, \infty) \times [0, \infty)$ with intensity measure $ds dz$,

$$Z_t^{N,i} = \int_0^t \int_0^\infty \mathbf{1}_{\{z \leq \mu + N^{-1} \sum_{j=1}^N \int_0^{s-} \varphi(s-u) dZ_u^{N,j}\}} \pi^i(ds dz).$$

We have $\mathbb{E}[Z_t^{N,i}] = m_t$. Indeed, by exchangeability, we see that $\mathbb{E}[Z_t^{N,i}] = \mathbb{E}[Z_t^{N,1}]$ and that

$$\mathbb{E}[Z_t^{N,1}] = \int_0^t \left(\mu + N^{-1} \sum_{j=1}^N \int_0^s \varphi(s-u) d\mathbb{E}[Z_u^{N,j}] \right) ds = \int_0^t \left(\mu + \int_0^s \varphi(s-u) d\mathbb{E}[Z_u^{N,1}] \right) ds,$$

whence $(\mathbb{E}[Z_t^{N,1}])_{t \geq 0}$ solves (8), of which the unique solution is $(m_t)_{t \geq 0}$ by Lemma 24.

We next introduce $U_t^{N,i} = Z_t^{N,i} - m_t$ and the martingales (here $\tilde{\pi}^i(ds dz) = \pi^i(ds dz) - ds dz$)

$$M_t^{N,i} = \int_0^t \int_0^\infty \mathbf{1}_{\{z \leq \mu + N^{-1} \sum_{j=1}^N \int_0^{s-} \varphi(s-u) dZ_u^{N,j}\}} \tilde{\pi}^i(ds dz).$$

We consider the mean processes $\bar{Z}_t^N = N^{-1} \sum_1^N Z_t^{N,i}$, $\bar{U}_t^N = N^{-1} \sum_1^N U_t^{N,i}$ and finally $\bar{M}_t^N = N^{-1} \sum_1^N M_t^{N,i}$. An easy computation using (8) and Lemma 22 shows that

$$\begin{aligned} U_t^{N,i} &= M_t^{N,i} + \int_0^t N^{-1} \sum_{j=1}^N \int_0^s \varphi(s-u) dZ_u^{N,j} ds - m_t \\ &= M_t^{N,i} + \int_0^t \int_0^s \varphi(s-u) \left(N^{-1} \sum_{j=1}^N dZ_u^{N,j} - dm_u \right) ds \\ &= M_t^{N,i} + \int_0^t \varphi(t-s) \left(N^{-1} \sum_{j=1}^N Z_s^{N,j} - m_s \right) ds, \end{aligned}$$

so that

$$(15) \quad U_t^{N,i} = M_t^{N,i} + \int_0^t \varphi(t-s) \bar{U}_s^N ds.$$

This directly implies that

$$(16) \quad \bar{U}_t^N = \bar{M}_t^N + \int_0^t \varphi(t-s) \bar{U}_s^N ds.$$

Next, we observe that $[M^{N,i}, M^{N,j}]_t = 0$ for all $i \neq j$ (because these martingales a.s. never jump simultaneously) and that $[M^{N,i}, M^{N,i}]_t = Z_t^{N,i}$. Hence $[\bar{M}^N, \bar{M}^N]_t = N^{-1} \bar{Z}_t^N$. We thus have $\mathbb{E}[(M_t^{N,i})^2] = \mathbb{E}[Z_t^{N,i}] = m_t$ and $\mathbb{E}[(\bar{M}_t^N)^2] = N^{-1} \mathbb{E}[\bar{Z}_t^N] = N^{-1} m_t$.

Step 2. Recalling (16) and using that $\Lambda = \int_0^\infty \varphi(s) ds < 1$, we observe that $\sup_{[0,t]} |\bar{U}_s^N| \leq \sup_{[0,t]} |\bar{M}_s^N| + \Lambda \sup_{[0,t]} |\bar{U}_s^N|$. Consequently,

$$\mathbb{E} \left[\sup_{[0,t]} |\bar{U}_s^N| \right] \leq (1 - \Lambda)^{-1} \mathbb{E} \left[\sup_{[0,t]} |\bar{M}_s^N| \right] \leq CN^{-1/2} m_t^{1/2}$$

by the Doob and Cauchy-Schwarz inequalities. We easily deduce that

$$\mathbb{E} \left[\int_0^t \varphi(t-s) |\bar{U}_s^N| ds \right] \leq \Lambda \mathbb{E} \left[\sup_{[0,t]} |\bar{U}_s^N| \right] \leq CN^{-1/2} m_t^{1/2},$$

whence finally, recalling (15),

$$m_t^{-1} \mathbb{E}[|U_t^{N,i}|] \leq m_t^{-1} \mathbb{E}[|M_t^{N,i}|] + C m_t^{-1} N^{-1/2} m_t^{1/2} \leq C m_t^{-1/2}.$$

This says that $\mathbb{E}[|Z_t^{N,i}/m_t - 1|] \leq C m_t^{-1/2}$ and thus proves point 2.

Step 3. We then fix $\ell \geq 1$ and use (15) to write, for $i = 1, \dots, \ell$,

$$m_t^{1/2}(Z_t^{N,i}/m_t - 1) = m_t^{-1/2}U_t^{N,i} = m_t^{-1/2}M_t^{N,i} + m_t^{-1/2}\int_0^t \varphi(t-s)\overline{U}_s^N ds.$$

First, $\mathbb{E}[m_t^{-1/2}\int_0^t \varphi(t-s)|\overline{U}_s^N|ds] \leq CN^{-1/2}$, which tends to 0 as $(t, N) \rightarrow (\infty, \infty)$, by the estimate proved in Step 2. To conclude the proof of point 3, we thus only have to prove that $(m_t^{-1/2}M_t^{N,i})_{i=1, \dots, \ell}$ goes in law to $\mathcal{N}(0, I_\ell)$ as $(t, N) \rightarrow (\infty, \infty)$. To this end, we apply Lemma 12. The jumps of the martingales $M^{N,i}$ are uniformly bounded (by 1) and we have seen that $[M^{N,i}, M^{N,j}]_t = 0$ for all $i \neq j$. The function $(m_t)_{t \geq 0}$ is continuous and increases to infinity. It thus suffices to check that, as $(t, N) \rightarrow (\infty, \infty)$, $m_t^{-1}[M^{N,i}, M^{N,i}]_t \rightarrow 1$ in probability. Since $[M^{N,i}, M^{N,i}]_t = Z_t^{N,i}$, this is an immediate consequence of point 2. \square

We now turn to the supercritical case.

Proof of Theorem 11. We rewrite (8) as $m_t = \int_0^t (\mu + \int_0^s \varphi(s-u)dm_u)ds = \mu t + \int_0^t \varphi(t-s)m_s ds$ by Lemma 22. This equation is studied in details in Lemma 26: there is a unique $\alpha_0 > 0$ such that $\mathcal{L}_\varphi(\alpha_0) = 1$ and, defining $a_0 \in (0, \infty)$ as in the statement, we have $m_t \sim a_0 e^{\alpha_0 t}$ and $m'_t \sim a_0 \alpha_0 e^{\alpha_0 t}$ as $t \rightarrow \infty$, which proves point 1. We also know that $\Gamma(t) = \sum_{n \geq 1} \varphi^{*n}(t) \sim (a_0 \alpha_0^2 / \mu) e^{\alpha_0 t}$, that $\Upsilon(t) = \int_0^t \Gamma(s)ds \sim (a_0 \alpha_0 / \mu) e^{\alpha_0 t}$ and further properties of m, m', Γ, Υ are proved in Lemma 26.

Step 1. We adopt the same notation as in the proof of Theorem 10-Step 1, of which all the results remain valid in the present case. Point 1 follows from Lemma 26-(a).

Step 2. First, (16) says exactly that $\overline{U}^N = \overline{M}^N + \varphi \star \overline{U}^N$. Using Lemma 26-(e), we deduce that $\overline{U}^N = \overline{M}^N + \Gamma \star \overline{M}^N$ (the processes \overline{U}^N and \overline{M}^N are clearly a.s. càdlàg and thus locally bounded). Since $\mathbb{E}[(\overline{M}_t^N)^2] = N^{-1}m_t$ by Step 1,

$$\begin{aligned} \mathbb{E}[|\overline{U}_t^N|] &\leq \mathbb{E}[|\overline{M}_t^N|] + \int_0^t \Gamma(t-s)\mathbb{E}[|\overline{M}_s^N|]ds \\ &\leq N^{-1/2}m_t^{1/2} + \int_0^t \Gamma(t-s)N^{-1/2}m_s^{1/2}ds \\ &\leq CN^{-1/2}(1+m_t). \end{aligned}$$

The last inequality easily follows from Lemma 26-(b).

Using (16) again, we see that $\int_0^t \varphi(t-s)\overline{U}_s^N ds = \overline{U}_t^N - \overline{M}_t^N$, whence

$$\mathbb{E}\left[\left|\int_0^t \varphi(t-s)\overline{U}_s^N\right|\right] \leq \mathbb{E}[|\overline{M}_t^N|] + \mathbb{E}[|\overline{U}_t^N|] \leq N^{-1/2}m_t^{1/2} + CN^{-1/2}(1+m_t) \leq CN^{-1/2}(1+m_t).$$

On the other hand, we know from Step 1 that $\mathbb{E}[(M_t^{N,i})^2] = m_t$. Using (15), we conclude that

$$\mathbb{E}[|Z_t^{N,i}/m_t - 1|] = m_t^{-1}\mathbb{E}[|U_t^{N,i}|] \leq m_t^{-1/2} + CN^{-1/2}(1+m_t^{-1}).$$

which ends the proof of 2.

Step 3. We then fix $\ell \geq 1$ and write, for $i = 1, \dots, \ell$, by (15),

$$(Z_t^{N,i}/m_t - 1) = m_t^{-1}U_t^{N,i} = m_t^{-1}M_t^{N,i} + m_t^{-1}\int_0^t \varphi(t-s)\overline{U}_s^N ds.$$

Step 3.1. We first consider the regime $(t, N) \rightarrow (\infty, \infty)$ with $m_t/N \rightarrow 0$ and study

$$m_t^{1/2}(Z_t^{N,i}/m_t - 1) = m_t^{-1/2}M_t^{N,i} + m_t^{-1/2} \int_0^t \varphi(t-s) \overline{U}_s^N ds.$$

The second term tends to 0 in probability, because we can bound, using Step 2, its L^1 -norm by $Cm_t^{-1/2}N^{-1/2}(1+m_t)$, which tends to 0 in the present regime. We thus just have to prove that $(m_t^{-1/2}M_t^{N,i})_{i=1,\dots,\ell}$ goes in law to $\mathcal{N}(0, I_\ell)$. We use Lemma 12: the martingales $M^{N,i}$ have uniformly bounded (by 1) jumps and we have seen that $[M^{N,i}, M^{N,j}]_t = 0$ for $i \neq j$. The function $(m_t)_{t \geq 0}$ is continuous and increases to infinity. It only remains to check that $m_t^{-1}[M^{N,i}, M^{N,i}]_t$ tends to 1 in probability. But $[M^{N,i}, M^{N,i}]_t = Z_t^{N,i}$, so that the conclusion follows from point 2.

Step 3.2. We finally consider the regime $(t, N) \rightarrow (\infty, \infty)$ with $m_t/N \rightarrow \infty$ and study

$$N^{1/2}(Z_t^{N,i}/m_t - 1) = N^{1/2}m_t^{-1}M_t^{N,i} + N^{1/2}m_t^{-1} \int_0^t \varphi(t-s) \overline{U}_s^N ds.$$

First, $N^{1/2}m_t^{-1}M_t^{N,i} \rightarrow 0$ in probability, because its L^1 -norm is bounded by $N^{1/2}m_t^{-1/2}$ (recall that $\mathbb{E}[(M_t^{N,i})^2] = m_t$), which tends to 0 in the present regime. Since $V_t^N := N^{1/2}m_t^{-1} \int_0^t \varphi(t-s) \overline{U}_s^N ds$ does not depend on i , it only remains to prove that V_t^N goes in law to $\mathcal{N}(0, \sigma^2)$. We write, using (16), recalling that $\overline{U}^N = \overline{M}^N + \Gamma \star \overline{M}^N$ (see Step 2) and integrating by parts (recall that $\Upsilon(t) = \int_0^t \Gamma(s) ds$)

$$V_t^N = N^{1/2}m_t^{-1}(\overline{U}_t^N - \overline{M}_t^N) = N^{1/2}m_t^{-1} \int_0^t \Gamma(t-s) \overline{M}_s^N ds = N^{1/2}m_t^{-1} \int_0^t \Upsilon(t-s) d\overline{M}_s^N.$$

Introduce $W_t^N = (\alpha_0/\mu)N^{1/2} \int_0^t e^{-\alpha_0 s} d\overline{M}_s^N$ and observe that, since $\mathbb{E}[[\overline{M}^N, \overline{M}^N]_t] = N^{-1}m_t$,

$$\begin{aligned} \mathbb{E}[(V_t^N - W_t^N)^2] &= \mathbb{E}\left[N \int_0^t (m_t^{-1}\Upsilon(t-s) - (\alpha_0/\mu)e^{-\alpha_0 s})^2 d[\overline{M}^N, \overline{M}^N]_s\right] \\ &= \int_0^t (m_t^{-1}\Upsilon(t-s) - (\alpha_0/\mu)e^{-\alpha_0 s})^2 m'_s ds. \end{aligned}$$

Lemma 26-(c) tells us that this tends to 0 as $t \rightarrow \infty$. We thus only have to prove that W_t^N goes in law to $\mathcal{N}(0, \sigma^2)$ as $(t, N) \rightarrow (\infty, \infty)$.

This follows again from Lemma 12 (with $\ell = 1$): the jumps of the martingale $(W_t^N)_{t \geq 0}$ are bounded by $(\alpha_0/\mu)N^{-1/2}$ (because those of \overline{M}^N are bounded by N^{-1}). The function $v_t = (\alpha_0/\mu)(\int_0^t e^{-2\alpha_0 s} m'_s ds)^{1/2}$ is continuous and increasing to the finite limit $v_\infty = \sigma$ (which was defined in the statement). We thus only have to prove that (a) $v_t^{-2}[W^N, W^N]_t \rightarrow 1$ in probability as $(t, N) \rightarrow (\infty, \infty)$, (b) for all $t_0 > 0$, uniformly in $t \geq t_0$, $v_t^{-2}[W^N, W^N]_t \rightarrow 1$ in probability as $N \rightarrow \infty$. By Lemma 12, we will deduce that $v_t^{-1}W_t^N$ goes in law to $\mathcal{N}(0, 1)$ as $(t, N) \rightarrow (\infty, \infty)$, which of course implies that W_t^N goes in law to $\mathcal{N}(0, \sigma^2)$ as desired.

We have, since $[\overline{M}^N, \overline{M}^N]_t = N^{-1}\overline{Z}_t^N$,

$$[W^N, W^N]_t = (\alpha_0/\mu)^2 N \int_0^t e^{-2\alpha_0 s} d[\overline{M}^N, \overline{M}^N]_s = (\alpha_0/\mu)^2 \int_0^t e^{-2\alpha_0 s} d\overline{Z}_s^N.$$

Using that $\bar{Z}_t^N = \bar{U}_t^N + m_t$ and performing an integration by parts, we see that

$$\begin{aligned} [W^N, W^N]_t &= v_t^2 + (\alpha_0/\mu)^2 \int_0^t e^{-2\alpha_0 s} d\bar{U}_s^N \\ &= v_t^2 + (\alpha_0/\mu)^2 e^{-2\alpha_0 t} \bar{U}_t^N + 2(\alpha_0^3/\mu^2) \int_0^t e^{-2\alpha_0 s} \bar{U}_s^N ds. \end{aligned}$$

Recalling that that $\mathbb{E}[|\bar{U}_t^N|] \leq CN^{-1/2}(1+m_t)$, we infer

$$\begin{aligned} &\mathbb{E}\left[\left|(\alpha_0/\mu)^2 e^{-2\alpha_0 t} \bar{U}_t^N + 2(\alpha_0^3/\mu^2) \int_0^t e^{-2\alpha_0 s} \bar{U}_s^N ds\right|\right] \\ &\leq \frac{C}{N^{1/2}} \left(e^{-2\alpha_0 t} (1+m_t) + \int_0^t e^{-2\alpha_0 s} (1+m_s) ds \right), \end{aligned}$$

which is bounded by $CN^{-1/2}$ by Lemma 26. We have proved that $\sup_{t \geq 0} \mathbb{E}[|[W^N, W^N]_t - v_t^2|] \leq CN^{-1/2}$, from which points (a) and (b) above immediately follow. The proof is complete. \square

4. NEAREST NEIGHBOUR MODEL

We consider here the case where \mathbb{G} is a regular grid, on which particles interact (directly) only if they are neighbours. We will work on \mathbb{Z}^d , endowed with the set of edges

$$\mathcal{E} = \{(i, j) \in (\mathbb{Z}^d)^2 : |i - j| = 0 \text{ or } 1\},$$

where $|(i_1, \dots, i_d)| = (\sum_{r=1}^d i_r^2)^{1/2}$. Thus each point has $2d + 1$ neighbours (including itself). We hesitated to include self-interaction, but this avoids some needless complications due to the periodicity of the underlying random walk on \mathbb{Z}^d .

Assumption 13. (i) The graph $\mathbb{G} = (\mathcal{S}, \mathcal{E})$ is $\mathcal{S} = \mathbb{Z}^d$ (for some $d \geq 1$) endowed with the above set of edges \mathcal{E} .

(ii) There is a nonnegative locally integrable function $\varphi : [0, \infty) \mapsto [0, \infty)$ such that for all $(j, i) \in \mathcal{E}$, $\varphi_{ji} = (2d + 1)^{-1} \varphi$.

(iii) For all $i \in \mathbb{Z}^d$, there is $\mu_i \geq 0$ such that $h_i(x) = \mu_i + x$. The family $(\mu_i)_{i \in \mathbb{Z}^d}$ is bounded.

We next introduce a few notation. In the whole section, we call *vector* (and write in bold) a family of numbers indexed by \mathbb{Z}^d . We call *matrix* a family indexed by $\mathbb{Z}^d \times \mathbb{Z}^d$. The identity matrix I is of course defined as $I(i, j) = \mathbf{1}_{\{i=j\}}$. We will often use the product of a matrix and a vector. The matrix $A = (A(i, j))_{i, j \in \mathbb{Z}^d}$ defined by

$$(17) \quad A(i, j) = (2d + 1)^{-1} \mathbf{1}_{\{(i, j) \in \mathcal{E}\}}$$

will play an important role. Since A is a stochastic matrix, we can define, for any $\Lambda \in (0, 1)$,

$$(18) \quad Q_\Lambda(i, j) = \sum_{n \geq 0} \Lambda^n A^n(i, j).$$

4.1. Large-time behaviour. Under Assumption 13, we can use Theorem 6 (with $p_i = 2^{-|i|}$, see Remark 5-(ii)): there is a unique Hawkes process $(Z_t^i)_{i \in \mathbb{Z}^d, t \geq 0}$ with parameters $(\mathbb{G}, \varphi, \mathbf{h})$ such that $\sum_{i \in \mathbb{Z}^d} 2^{-|i|} \mathbb{E}[Z_t^i] < \infty$. Let us state the first results of this section. As usual, we treat separately the subcritical and supercritical cases.

Theorem 14. *Work under Assumption 13 and assume further that $\Lambda = \int_0^\infty \varphi(t) dt < 1$. Consider the unique Hawkes process $(Z_t^i)_{i \in \mathbb{Z}^d, t \geq 0}$ with parameters $(\mathbb{G}, \varphi, \mathbf{h})$. For all $i \in \mathbb{Z}^d$, $t^{-1} Z_t^i$ goes in probability, as $t \rightarrow \infty$, to $\sum_{j \in \mathbb{Z}^d} Q_\Lambda(i, j) \mu_j$.*

Theorem 15. *Work under Assumption 13 and assume further that $\Lambda = \int_0^\infty \varphi(t)dt \in (1, \infty]$ and that $t \mapsto \int_0^t |d\varphi(s)|$ has at most polynomial growth. Consider $\alpha_0 > 0$ uniquely defined by $\mathcal{L}_\varphi(\alpha_0) = 1$. Assume finally that the “mean value”*

$$(19) \quad \mu = \lim_{r \rightarrow \infty} \frac{1}{\#\{i \in \mathbb{Z}^d : |i| \leq r\}} \sum_{|i| \leq r} \mu_i \quad \text{exists and is positive.}$$

Consider the unique Hawkes process $(Z_t^i)_{i \in \mathbb{Z}^d, t \geq 0}$ with parameters $(\mathbb{G}, \varphi, \mathbf{h})$. Then for all $i \in \mathbb{Z}^d$, $e^{-\alpha_0 t} Z_t^i$ goes in probability, as $t \rightarrow \infty$, to $a_0 = \mu \alpha_0^{-2} (\int_0^\infty t \varphi(t) e^{-\alpha_0 t} dt)^{-1}$.

Let us comment on these results. In the subcritical case, the parameter $\boldsymbol{\mu} = (\mu_i)_{i \in \mathbb{Z}^d}$ is strongly present in the limiting behaviour: the limit of $t^{-1} Z_t^i$ depends on a certain mean of $\boldsymbol{\mu}$ around the site i and thus depends on i . In the supercritical case, the behaviour is very different: the limit value of $e^{-\alpha_0 t} Z_t^i$ does not depend on i , and depends on $\boldsymbol{\mu} = (\mu_i)_{i \in \mathbb{Z}^d}$ only through a *global* mean value. Observe also that for a finite-dimensional (e.g. *scalar*) Hawkes process, there is no law of large numbers: one can get a limit of something like $e^{-\alpha_0 t} Z_t^i$, but the limit is random, see Zhu [43, Section 5.4] (in particular Theorem 23 and Corollary 1). In that sense, we can say that in the supercritical case, the law of large number is reminiscent of the infinite dimension and of the interaction.

We will need a precise approximation for $A^n(i, j)$ where A is defined by (17). It is given by the local central limit theorem, since A is the transition matrix of an aperiodic symmetric random walk on \mathbb{Z}^d with bounded jumps. Precisely, we infer from Lawler-Limic [26, Theorem 2.1.1 and (2.5)] that there is a constant C such that for all $n \geq 1$, all $i \in \mathbb{Z}^d$,

$$(20) \quad |A^n(0, i) - p_n(i)| \leq \frac{C}{n^{(d+2)/2}}$$

where, for $t > 0$ and $x \in \mathbb{R}^d$,

$$(21) \quad p_t(x) = \left(\frac{2d+1}{4\pi t} \right)^{d/2} \exp\left(-\frac{(2d+1)|x|^2}{4t} \right).$$

To apply [26, Theorem 2.1.1], we needed to compute the covariance matrix Γ corresponding to our random walk, we found $\Gamma = 2(2d+1)^{-1} I_d$, I_d being the $d \times d$ identity matrix.

Lemma 16. *Consider the matrix $(A(i, j))_{i, j \in \mathbb{Z}^d}$ defined by (17).*

(i) *It holds that $\varepsilon_n = \sum_{j \in \mathbb{Z}^d} (A^n(i, j))^2$ does not depend on $i \in \mathbb{Z}^d$ and tends to 0 as $n \rightarrow \infty$.*

(ii) *Let $\boldsymbol{\mu} = (\mu_i)_{i \in \mathbb{Z}^d}$ be bounded and satisfy (19). Then for all $i \in \mathbb{Z}^d$, $\lim_{n \rightarrow \infty} (A^n \boldsymbol{\mu})_i = \mu$.*

Proof. In the following we denote by C a constant depending only on d .

Point (i) is easy: since $A^n(i, j) = A^n(0, j - i)$ and since A is stochastic, one has $\varepsilon_n = \sum_{j \in \mathbb{Z}^d} (A^n(0, j))^2 \leq \sup_{j \in \mathbb{Z}^d} A^n(0, j)$. Moreover, by (20), $A^n(0, j) \leq p_n(j) + Cn^{-(d+2)/2} \leq Cn^{-d/2}$. We conclude that $\varepsilon_n \leq Cn^{-d/2} \rightarrow 0$ as desired.

Now we turn to the proof of (ii). Let $i \in \mathbb{Z}^d$. First we show that $\lim_n [(A^n \boldsymbol{\mu})_i - \sum_{j \in \mathbb{Z}^d} p_n(j) \mu_j] = 0$. Since $(A^n \boldsymbol{\mu})_i = \sum_{j \in \mathbb{Z}^d} A^n(i, j) \mu_j = \sum_{j \in \mathbb{Z}^d} A^n(0, j - i) \mu_j$ and since the family $(\mu_j)_{j \in \mathbb{Z}^d}$ is bounded, it suffices to prove that $v_n = \sum_{j \in \mathbb{Z}^d} |A^n(0, j - i) - p_n(j)| \rightarrow 0$. We write

$$v_n \leq \sum_{|j| \leq n^{1/2+1/4d}} |A^n(0, j - i) - p_n(j)| + \sum_{|j| > n^{1/2+1/4d}} (A^n(0, j - i) + p_n(j)) = v_n^1 + v_n^2.$$

On the one hand, using that $\sum_{j \in \mathbb{Z}^d} |j|^2 A^n(0, j) \leq Cn$ (the variance of the random walk at time n is of order n), so that $\sum_{j \in \mathbb{Z}^d} |j|^2 A^n(0, j - i) = \sum_{k \in \mathbb{Z}^d} |i + k|^2 A^n(0, k) \leq C(|i|^2 + n)$ and thus

$$\sum_{|j| > n^{1/2+1/4d}} A^n(0, j - i) \leq Cn^{-1-1/2d} \sum_{j \in \mathbb{Z}^d} |j|^2 A^n(0, j - i) \leq Cn^{-1-1/2d}(|i|^2 + n).$$

Similarly, we have $\sum_{j \in \mathbb{Z}^d} |j|^2 p_n(j) \leq Cn$ and thus

$$\sum_{|j| > n^{1/2+1/4d}} p_n(j) \leq n^{-1-1/2d} \sum_{j \in \mathbb{Z}^d} |j|^2 p_n(j) \leq Cn^{-1/2d}.$$

Consequently $\lim_n v_n^2 = 0$. On the other hand,

$$v_n^1 \leq \sum_{|j| \leq n^{1/2+1/4d}} |A^n(0, j - i) - p_n(j - i)| + \sum_{|j| \leq n^{1/2+1/4d}} |p_n(j - i) - p_n(j)|.$$

From (20), the first sum is bounded by $Cn^{-(d+2)/2} \#\{j \in \mathbb{Z}^d : |j| \leq n^{1/2+1/4d}\} \leq Cn^{-3/4} \rightarrow 0$. For the second sum, we use that, with $c_d = (2d + 1)/4$,

$$|p_n(j - i) - p_n(j)| = p_n(j) \left| 1 - \exp\left(-\frac{c_d}{n}|i|^2 + \frac{2c_d}{n}i \cdot j\right) \right|.$$

Hence for $|j| \leq n^{1/2+1/4d}$ and for n large enough (e.g. so that $|i|n^{-1/2+1/4d} \leq 1$),

$$|p_n(j - i) - p_n(j)| \leq Cp_n(j)(|i|^2 n^{-1} + |i|n^{-1/2+1/4d}) \leq Cp_n(j)(1 + |i|^2)n^{-1/4}.$$

Thus $\sum_{|j| \leq n^{1/2+1/2d}} |p_n(j - i) - p_n(j)| \leq C(1 + |i|^2)n^{-1/4}$ and we deduce that $\lim_n v_n^1 = 0$.

We have shown that $\lim v_n = 0$. It only remains to check that $\lim_n \sum_{j \in \mathbb{Z}^d} \mu_j p_n(j) = \mu$. Let $(r_k)_{k \geq 0}$ be the increasing sequence of nonnegative numbers such that $\{r_k\}_{k \geq 0} = \{|j| : j \in \mathbb{Z}^d\}$ and observe that

$$\sum_{j \in \mathbb{Z}^d} \mu_j p_n(j) = \sum_{k \geq 0} p_n(r_k) \sum_{|j|=r_k} \mu_j.$$

A discrete integration by parts shows that

$$\sum_{j \in \mathbb{Z}^d} \mu_j p_n(j) = \sum_{k \geq 0} (p_n(r_k) - p_n(r_{k+1})) \sum_{|j| \leq r_k} \mu_j = \sum_{k \geq 0} v(r_k)(p_n(r_k) - p_n(r_{k+1})) \frac{1}{v(r_k)} \sum_{|j| \leq r_k} \mu_j,$$

where $v(r) = \#\{j \in \mathbb{Z}^d : |j| \leq r\}$. We easily conclude that $\lim_n \sum_{j \in \mathbb{Z}^d} \mu_j p_n(j) = \mu$ as desired, because

- (a) $\lim_{k \rightarrow \infty} \frac{1}{v(r_k)} \sum_{|j| \leq r_k} \mu_j = \mu$;
- (b) for all $k \geq 0$ fixed, $\lim_n v(r_k)(p_n(r_k) - p_n(r_{k+1})) = 0$;
- (c) $\lim_{n \rightarrow \infty} \sum_{k=0}^{\infty} v(r_k)(p_n(r_k) - p_n(r_{k+1})) = 1$.

Point (a) follows from our condition (19) on μ , point (b) is obvious (because $|v(r_k)(p_n(r_k) - p_n(r_{k+1}))| \leq v(r_k) \sup_{i \in \mathbb{Z}^d} p_n(i) \leq Cv(r_k)n^{-d/2} \rightarrow 0$). To check (c), we write $\sum_{k=0}^{\infty} v(r_k)(p_n(r_k) - p_n(r_{k+1})) = \sum_{j \in \mathbb{Z}^d} p_n(j) = \sum_{j \in \mathbb{Z}^d} A^n(0, j) + \sum_{j \in \mathbb{Z}^d} [p_n(j) - A^n(0, j)] = 1 + \sum_{j \in \mathbb{Z}^d} [p_n(j) - A^n(0, j)]$. This tends to 1, because $\lim_n \sum_{j \in \mathbb{Z}^d} |p_n(j) - A^n(0, j)| = 0$, as seen in the first part of the proof (this is v_n in the special case where $i = 0$).

□

Let us now give the

Proof of Theorem 14. We split the proof into several steps. We assume that there is at least one $i \in \mathbb{Z}^d$ such that $\mu_i > 0$, because else the result is obvious (because then $Z_t^i = 0$ for all $i \in \mathbb{Z}^d$, all $t \geq 0$). The first step will also be used in the supercritical case.

Step 1. We write as usual, for some i.i.d. family $(\pi^i(ds dz))_{i \geq 1}$ of Poisson measures on $[0, \infty) \times [0, \infty)$ with intensity measure $ds dz$,

$$Z_t^i = \int_0^t \int_0^\infty \mathbf{1}_{\{z \leq \mu_i + (2d+1)^{-1} \sum_{j \rightarrow i} \int_0^{s-} \varphi(s-u) dZ_u^j\}} \pi^i(ds dz).$$

Let us put $m_t^i = \mathbb{E}[Z_t^i]$ and $\mathbf{m}_t = (m_t^i)_{i \in \mathbb{Z}^d}$. A simple computation (using one more time Lemma 22) gives us, for all $i \in \mathbb{Z}^d$,

$$m_t^i = \mu_i t + \int_0^t (2d+1)^{-1} \sum_{j \rightarrow i} \varphi(t-s) m_s^j ds.$$

Using the vector formalism, this rewrites $\mathbf{m}_t = \boldsymbol{\mu} t + \int_0^t \varphi(t-s) (A \mathbf{m}_s) ds$. We furthermore know (from Theorem 6) that for all $t \geq 0$, $\sum_{i \in \mathbb{Z}^d} 2^{-|i|} m_t^i < \infty$. Applying Lemma 27, we see that m^i is of class C^1 on $[0, \infty)$ for each $i \in \mathbb{Z}^d$, that

$$(22) \quad \mathbf{m}'_t = \boldsymbol{\mu} + \int_0^t \varphi(t-s) A \mathbf{m}'_s ds$$

and that

$$(23) \quad \mathbf{m}'_t = \left(I + \sum_{n \geq 1} A^n \int_0^t \varphi^{*n}(s) ds \right) \boldsymbol{\mu}.$$

Lemma 27 also tells us that $u_t = \sup_{i \in \mathbb{Z}^d} \sup_{[0,t]} (m_s^i)'$ is locally bounded, which of course implies that $\sup_{i \in \mathbb{Z}^d} \sup_{[0,t]} m_s^i$ is also locally bounded (because $m_0^i = 0$ for all $i \in \mathbb{Z}^d$), and that

$$(24) \quad u_t \leq C + \int_0^t \varphi(t-s) u_s ds.$$

We introduce the martingales, for $i \in \mathbb{Z}^d$, (we use a tilde for compensation),

$$M_t^i = \int_0^t \int_0^\infty \mathbf{1}_{\{z \leq \mu_i + (2d+1)^{-1} \sum_{j \rightarrow i} \int_0^{s-} \varphi(s-u) dZ_u^j\}} \tilde{\pi}^i(ds dz)$$

and observe as usual that $[M^i, M^j]_t = 0$ when $i \neq j$ (because these martingales a.s. never jump at the same time) while $[M^i, M^i]_t = Z_t^i$. We finally introduce $U_t^i = Z_t^i - m_t^i$, the vectors $\mathbf{U}_t = (U_t^i)_{i \in \mathbb{Z}^d}$ and $\mathbf{M}_t = (M_t^i)_{i \in \mathbb{Z}^d}$ and observe that

$$(25) \quad \mathbf{U}_t = \mathbf{M}_t + \int_0^t \varphi(t-s) A \mathbf{U}_s ds.$$

Indeed, for every $i \in \mathbb{Z}^d$, using Lemma 22 and the equation satisfied by m_t^i , we find

$$U_t^i = M_t^i + \int_0^t (2d+1)^{-1} \sum_{j \rightarrow i} \varphi(t-s) (Z_s^j - m_s^j) ds = M_t^i + \int_0^t \varphi(t-s) (A \mathbf{U}_s)_i ds.$$

Equation (25) can be solved as usual as

$$(26) \quad \mathbf{U}_t = \left(\mathbf{M}_t + \sum_{n \geq 1} \int_0^t \varphi^{*n}(t-s) A^n \mathbf{M}_s ds \right).$$

Finally, we easily check that $v_t = \sup_{i \in \mathbb{Z}^d} \sup_{[0,t]} \mathbb{E}[|U_s^i|]$ is locally bounded (because $\mathbb{E}[|U_t^i|] \leq \mathbb{E}[|Z_t^i|] + m_t^i \leq 2m_t^i$) and satisfies (start from (25), use that $\mathbb{E}[|M_t^i|] \leq \mathbb{E}[|M^i, M^i|_t]^{1/2} = \mathbb{E}[|Z_t^i|]^{1/2} = (m_t^i)^{1/2} \leq (\int_0^t u_s ds)^{1/2}$) and that A is stochastic

$$(27) \quad v_t \leq \left(\int_0^t u_s ds \right)^{1/2} + \int_0^t \varphi(t-s) v_s ds.$$

Step 2. Here we prove that there is a constant C such that for all $i \in \mathbb{Z}^d$, $(m_t^i)' \leq C$ (and thus also $m_t^i \leq Ct$). This follows from (24), which implies that $u_t \leq C + \Lambda u_t$, whence $u_t \leq C/(1-\Lambda)$.

Step 3. For all $i \in \mathbb{Z}^d$, $(m_t^i)' \sim (Q_\Lambda \boldsymbol{\mu})_i$, whence also $m_t^i \sim (Q_\Lambda \boldsymbol{\mu})_i t$, as $t \rightarrow \infty$. Indeed, starting from (23), using the monotone convergence theorem and that $\int_0^\infty \varphi^{*n}(s) ds = (\int_0^\infty \varphi(s) ds)^n = \Lambda^n$,

$$\lim_{t \rightarrow \infty} (m_t^i)' = \left(\left(I + \sum_{n \geq 1} \Lambda^n A^n \right) \boldsymbol{\mu} \right)_i = (Q_\Lambda \boldsymbol{\mu})_i.$$

Step 4. There is a constant C such that for all $i \in \mathbb{Z}^d$, all $t \geq 0$, $\mathbb{E}[|U_t^i|] \leq Ct^{1/2}$. Indeed, this follows from (27) and Step 2, which imply that $v_t \leq Ct^{1/2} + \Lambda v_t$, whence $v_t \leq Ct^{1/2}/(1-\Lambda)$.

Step 5. The conclusion follows immediately, writing

$$\mathbb{E} \left[\left| \frac{Z_t^i}{t} - (Q_\Lambda \boldsymbol{\mu})_i \right| \right] \leq \mathbb{E} \left[\left| \frac{U_t^i}{t} \right| \right] + \left| \frac{m_t^i}{t} - (Q_\Lambda \boldsymbol{\mu})_i \right|,$$

which tends to 0 as $t \rightarrow \infty$ by Steps 3 and 4. \square

We now turn to the supercritical case.

Proof of Theorem 15. We consider m (not to be confused with \mathbf{m}) the unique solution to $m_t = \mu t + \int_0^t \varphi(t-s) m_s ds$, where μ is the mean value defined by (19). This equation is studied in details in Lemma 26: with α_0 and a_0 defined in the statement, we have $m_t \sim a_0 e^{\alpha_0 t}$ and $m_t' \sim a_0 \alpha_0 e^{\alpha_0 t}$ as $t \rightarrow \infty$, as well as $\Gamma(t) = \sum_{n \geq 1} \varphi^{*n}(t) \sim (a_0 \alpha_0^2 / \mu) e^{\alpha_0 t}$ and $\Upsilon(t) = \int_0^t \Gamma(s) ds \sim (a_0 \alpha_0 / \mu) e^{\alpha_0 t}$.

Step 1. We adopt the notation introduced in Step 1 of the proof of Theorem 14.

Step 2. Here we check that there is C such that for all $i \in \mathbb{Z}^d$, $(m_t^i)' \leq C e^{\alpha_0 t}$ (and thus $m_t^i \leq C e^{\alpha_0 t}$). This follows from (24), which tells us that $u_t = \sup_{i \in \mathbb{Z}^d} (m_t^i)'$ is locally bounded and satisfies $u_t \leq C + \int_0^t \varphi(t-s) u_s ds$. Setting $h_t = u_t - \int_0^t \varphi(t-s) u_s ds$, we see that h_t is locally bounded (from above and from below), because u is locally bounded and φ is locally integrable. We furthermore have $u = h + u \star \varphi$. Applying Lemma 26-(e), we deduce that $u = h + h \star \Gamma$. But h is bounded from above by C . Consequently, $u \leq C + C \star \Gamma = C(1 + \Upsilon)$, where Υ was defined in Lemma 26. The conclusion follows from Lemma 26-(b).

Step 3. We now show that for all $i \in \mathbb{Z}^d$, $(m_t^i)' \sim (m_t)'$ (whence $m_t^i \sim m_t$) as $t \rightarrow \infty$. Let us fix $i \in \mathbb{Z}^d$ and set $r_t^i = (m_t^i)' - (m_t)'$, which satisfies $r_t^i = \mu_i - \mu + \sum_{n \geq 1} ((A^n \boldsymbol{\mu})_i - \mu) \int_0^t \varphi^{*n}(s) ds$. We know from Lemma 16-(ii) that $\eta_n = (A^n \boldsymbol{\mu})_i - \mu$ tends to 0 as $n \rightarrow \infty$. Consequently, Lemma 26-(d) tells us that $e^{-\alpha_0 t} \sum_{n \geq 1} ((A^n \boldsymbol{\mu})_i - \mu) \int_0^t \varphi^{*n}(s) ds$ tends to 0 as $t \rightarrow \infty$. Hence $e^{-\alpha_0 t} r_t^i \rightarrow 0$ as $t \rightarrow \infty$. Using finally that $m_t' \sim a_0 \alpha_0 e^{\alpha_0 t}$ as $t \rightarrow \infty$, we conclude that $(m_t^i)' / m_t' = 1 + r_t^i / m_t' \sim 1 + (a_0 \alpha_0)^{-1} e^{-\alpha_0 t} r_t^i \rightarrow 1$ as desired.

Step 4. Here we check that for every $i \in \mathbb{Z}^d$, $e^{-\alpha_0 t} \mathbb{E}[|U_t^i|]$ tends to 0 as $t \rightarrow \infty$. We start from (26) to write

$$|U_t^i| \leq |M_t^i| + \sum_{n \geq 1} \int_0^t \varphi^{*n}(t-s) |(A^n \mathbf{M}_s)_i| ds.$$

But $\mathbb{E}[|M_t^i|] \leq \mathbb{E}[Z_t^i]^{1/2} = (m_t^i)^{1/2} \leq C e^{\alpha_0 t/2}$ by Step 2 and $\mathbb{E}[(A^n \mathbf{M}_t)_i^2] = \sum_j (A^n(i, j))^2 m_s^j \leq C e^{\alpha_0 t} \sum_j (A^n(i, j))^2 \leq C e^{\alpha_0 t} \varepsilon_n$ by Lemma 16-(i), with $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$. Consequently,

$$e^{-\alpha_0 t} \mathbb{E}[|U_t^i|] \leq C e^{-\alpha_0 t/2} + C e^{-\alpha_0 t} \sum_{n \geq 1} \varepsilon_n^{1/2} \int_0^t \varphi^{*n}(t-s) e^{\alpha_0 s/2} ds.$$

Lemma 26-(d) allows us to conclude.

Step 5. The conclusion follows, writing

$$\mathbb{E} \left[\left| \frac{Z_t^i}{a_0 e^{\alpha_0 t}} - 1 \right| \right] \leq \mathbb{E} \left[\left| \frac{U_t^i}{a_0 e^{\alpha_0 t}} \right| \right] + \left| \frac{m_t^i}{a_0 e^{\alpha_0 t}} - 1 \right|,$$

and using Steps 3, 4, and that $m_t \sim a_0 e^{\alpha_0 t}$ by Lemma 26-(a). \square

4.2. Study of an impulsion. Here we want to study how an *impulsion* at time 0 at $i = 0$ propagates. To this end, we work under Assumption 13 with $\mu_i = 0$ for all $i \in \mathbb{Z}^d$, but we assume that Z^0 has a jump at time 0. Such a study is of course important: it allows us to measure, in some sense, the range of the interaction.

We first define precisely the process under study.

Definition 17. *We work under Assumption 13-(i)-(ii) and consider a family $(\pi^i(ds dz), i \in \mathbb{Z}^d)$ of i.i.d. $(\mathcal{F}_t)_{t \geq 0}$ -Poisson measures on $[0, \infty) \times [0, \infty)$ with intensity measure $ds dz$. We say that a family $(Z_t^i)_{i \in \mathbb{Z}^d, t \geq 0}$ of $(\mathcal{F}_t)_{t \geq 0}$ -adapted counting processes is an impulsion Hawkes process if*

$$\forall i \in \mathbb{Z}^d, \quad Z_t^i = \int_0^t \int_0^\infty \mathbf{1}_{\left\{ z \leq \sum_{j \rightarrow i} (2d+1)^{-1} \left[\int_0^{s^-} \varphi(s-u) dZ_u^j + \varphi(s) \mathbf{1}_{\{j=0\}} \right] \right\}} \pi^i(ds dz).$$

As said previously, the term $\sum_{j \rightarrow i} \varphi(s) \mathbf{1}_{\{j=0\}}$ is interpreted as an *excitation* due to a *forced* jump of Z^0 at time 0: simply rewrite it as $\mathbf{1}_{\{0 \rightarrow i\}} \int_0^{s^-} \varphi(s-u) \delta_u(ds)$.

The following proposition is easy.

Proposition 18. *Adopt the assumptions and notation of Definition 17. There exists a pathwise unique impulsion Hawkes process $((Z_t^i)_{i \in \mathbb{Z}^d, t \geq 0})$ such that $\sum_{i \in \mathbb{Z}^d} \mathbb{E}[Z_t^i] < \infty$ for all $t \geq 0$.*

Exactly as Theorem 6, this result can be deduced from Massoulié [28, Theorem 2] in the sub-critical case where $\int_0^\infty \varphi(s) ds < 1$.

Proof. The proof resembles much that of Theorem 6, so we only sketch it. We start with uniqueness and thus consider two impulsion Hawkes processes $(Z_t^i)_{i \in \mathbb{Z}^d, t \geq 0}$ and $(\tilde{Z}_t^i)_{i \in \mathbb{Z}^d, t \geq 0}$ such that $\sum_{i \in \mathbb{Z}^d} \mathbb{E}[Z_t^i + \tilde{Z}_t^i] < \infty$. We set $\Delta_t^i = \int_0^t |d(Z_s^i - \tilde{Z}_s^i)|$, $\delta_t^i = \mathbb{E}[\Delta_t^i]$ and $\delta_t = \sum_{i \in \mathbb{Z}^d} \delta_t^i$ (which is locally bounded by assumption). We may check that $\delta_t^i \leq (2d+1)^{-1} \int_0^t \sum_{j \rightarrow i} \varphi(t-s) \delta_s^j ds$ exactly as in the proof of Theorem 6. Summing in i and recalling that each site has $2d+1$ neighbours, we find $\delta_t \leq \int_0^t \varphi(t-s) \delta_s ds$. Lemma 23-(i) tells us that $\delta_t = 0$ for all t , whence pathwise uniqueness.

Existence follows from a Picard iteration. Let us only check an *a priori* estimate implying that $\sum_{i \in \mathbb{Z}^d} \mathbb{E}[Z_t^i] < \infty$ for all $t \geq 0$. Set $m_t^i = \mathbb{E}[Z_t^i]$ and $m_t = \sum_{i \in \mathbb{Z}^d} m_t^i$. A direct computation using Lemma 22 shows that $m_t^i = (2d+1)^{-1} \sum_{j \rightarrow i} [\int_0^t \varphi(t-s) m_s^j ds + \mathbf{1}_{\{j=0\}} \int_0^t \varphi(s) ds]$. Summing in i , we find $m_t = \int_0^t \varphi(t-s) m_s ds + \int_0^t \varphi(s) ds$. Using Lemma 23-(i), that φ is locally integrable (and that $\int_0^t \varphi(s) ds$ is locally bounded), we deduce that $\sup_{[0, T]} m_t \leq C(T, \varphi)$ as desired. \square

We next compute the probability of the extinction event. Point 1 is a noticeable property that makes the result very easy and precise.

Theorem 19. *Adopt the assumptions and notation of Definition 17 and consider the impulsion Hawkes process $((Z_t^i)_{i \in \mathbb{Z}^d, t \geq 0})$.*

1. *The process $Z_t = \sum_{i \in \mathbb{Z}^d} Z_t^i$ is a scalar impulsion Hawkes process with excitation function φ . In other words, $(Z_t)_{t \geq 0}$ is a counting process with compensator $A_t = \int_0^t \lambda_s ds$, where*

$$\lambda_t = \varphi(t) + \int_0^{t-} \varphi(t-s) dZ_s.$$

2. *We introduce the extinction event defined by $\Omega_e = \{\lim_{t \rightarrow \infty} \sum_{i \in \mathbb{Z}^d} Z_t^i < \infty\}$. Setting $\Lambda = \int_0^\infty \varphi(s) ds$, we have (i) $\Pr(\Omega_e) = 0$ if $\Lambda = \infty$; (ii) $\Pr(\Omega_e) = \exp(-\gamma_\Lambda \Lambda)$ if $\Lambda \in (1, \infty)$, where $\gamma_\Lambda \in (0, 1)$ is characterised by $\gamma_\Lambda \Lambda + \log(1 - \gamma_\Lambda) = 0$; (iii) $\Pr(\Omega_e) = 1$ if $\Lambda \in (0, 1]$.*

Of course, we can sometimes use this theorem, by a simple comparison argument, if φ_{ji} depends on i, j . For example, to guarantee non-extinction with probability one, it suffices that all the φ_{ji} are bounded below by some $(2d+1)^{-1} \varphi$ such that $\int_0^\infty \varphi(s) ds = \infty$.

When $\Lambda \in (0, 1)$, a detailed study of the tail distribution of the extinction time is handled by Brémaud-Nappo-Torriani [9], for a more general scalar model with marks.

Proof. Point 1 is immediate: the compensator of the counting process $(Z_t)_{t \geq 0}$ is

$$\begin{aligned} A_t &= \int_0^t \sum_{i \in \mathbb{Z}^d} \sum_{j \rightarrow i} (2d+1)^{-1} [\int_0^{s-} \varphi(s-u) dZ_u^j + \varphi(s) \mathbf{1}_{\{j=0\}}] ds \\ &= \int_0^t \sum_{j \in \mathbb{Z}^d} [\int_0^{s-} \varphi(s-u) dZ_u^j + \varphi(s) \mathbf{1}_{\{j=0\}}] ds \\ &= \int_0^t [\varphi(s) + \int_0^{s-} \varphi(s-u) dZ_u] ds. \end{aligned}$$

We next prove point 2. It is well-known Folklore that a scalar impulsion Hawkes process can be related to a Poisson Galton-Watson process with Poisson(Λ) reproduction law, but we give a direct proof for the sake of completeness. If $\Lambda = \infty$, it suffices to note that $\Pr(\Omega_e) = \Pr(Z_\infty < \infty) = \Pr(A_\infty < \infty) \leq \Pr(\int_0^\infty \varphi(s) ds < \infty) = 0$. When $\Lambda < \infty$, we introduce the martingale, for $\gamma \in (0, 1)$, $N_t^\gamma = -\gamma(Z_t - A_t) = -\gamma Z_t + \gamma \int_0^t \varphi(s) ds + \gamma \int_0^t \varphi(t-s) Z_s ds$ by Lemma 22. We denote by $M_t^\gamma = \mathcal{E}(N^\gamma)_t = e^{\gamma A_t} \prod_{s \leq t} (1 - \gamma \Delta Z_s)$ its Doléans-Dade exponential, see Jacod-Shiryaev [23, Chapter 1, Section 4f]. Since Z is a counting process, we see that

$$M_t^\gamma = \exp\left(\gamma \int_0^t \varphi(s) ds + \gamma \int_0^t \varphi(t-s) Z_s ds + \log(1 - \gamma) Z_t\right).$$

If $\gamma\Lambda + \log(1-\gamma) \leq 0$ then M^γ is bounded (because $\gamma \int_0^t \varphi(s)ds + \gamma \int_0^t \varphi(t-s)Z_s ds + \log(1-\gamma)Z_t \leq \gamma\Lambda + (\gamma\Lambda + \log(1-\gamma))Z_t \leq \gamma\Lambda$), and thus converges in L^1 . Consequently, $\mathbb{E}[M_\infty^\gamma] = 1$ and (one easily verifies, using that Z is non-decreasing, that $\lim_{t \rightarrow \infty} \int_0^t \varphi(t-s)Z_s ds = \Lambda Z_\infty$)

$$\mathbb{E}\left[\exp\left(\gamma\Lambda + (\log(1-\gamma) + \gamma\Lambda)Z_\infty\right)\right] = 1.$$

But for all $x > 0$, there is a unique $\gamma(x) \in (0, 1)$ such that $\gamma(x)\Lambda + \log(1-\gamma(x)) = -x$, whence

$$\mathbb{E}\left[\exp\left(-xZ_\infty\right)\right] = \exp(-\Lambda\gamma(x)).$$

Consequently,

$$\Pr(\Omega_e) = \Pr(Z_\infty < \infty) = \lim_{x \rightarrow 0^+} \mathbb{E}\left[\exp\left(-xZ_\infty\right)\right] = \exp(-\Lambda\gamma(0+)).$$

If $\Lambda \in (0, 1]$, we see that $\gamma(0+) = 0$, so that $\Pr(\Omega_e) = 1$. If now $\Lambda > 1$, $\gamma(0+)$ is the unique solution in $(0, 1)$ to $\gamma(0+)\Lambda + \log(1-\gamma(0+)) = 0$. \square

We next study more deeply, in the super-critical case, how the impulsion propagates. Unfortunately, the computations are really tedious: we decided to restrict ourselves to a particular case (φ is an exponential function) where some computations are explicit. We believe that the result below can be extended to a general class of functions φ , but a difficult technical lemma is required.

Theorem 20. *Work under Assumption 13-(i)-(ii), with $\varphi(t) = ae^{-bt}$, for some $a > b > 0$. Consider the impulsion Hawkes process $((Z_t^i)_{i \in \mathbb{Z}^d, t \geq 0})$. Since $\int_0^\infty \varphi(s)ds = a/b > 1$, we know from Theorem 19 that $\Pr(\Omega_e) \in (0, 1)$. We set $\alpha_0 = a - b$ (for which $\mathcal{L}_\varphi(\alpha_0) = 1$) and we recall that the Gaussian density $p_t(x)$ is defined by (21).*

(i) *There are some constants $C > 0$ and $t_0 > 0$ and a random variable $H \geq 0$ such that for all $i \in \mathbb{Z}^d$, all $t \geq t_0$,*

$$\mathbb{E}[|Z_t^i - Hp_{at}(i)e^{\alpha_0 t}|] \leq \frac{Ce^{\alpha_0 t}}{t^{d/2+1/3}}.$$

(ii) *For all $x \in \mathbb{R}^d$, $t^{d/2}e^{-\alpha_0 t}Z_t^{\lfloor xt^{1/2} \rfloor} \rightarrow Hp_a(x)$ in probability as $t \rightarrow \infty$. Here, we define the “integer part” $\lfloor y \rfloor$ of $y = (y_1, \dots, y_d) \in \mathbb{R}^d$, by $\lfloor y \rfloor = (\lfloor y_1 \rfloor, \dots, \lfloor y_d \rfloor) \in \mathbb{Z}^d$.*

(iii) *The random variable H is positive on the event Ω_e^c .*

This result describes quite precisely how an impulsion propagates. Conditional on non extinction, and the process $(Z_t^i)_{i \in \mathbb{Z}^d}$ resembles a Gaussian profile, with height $t^{-d/2}He^{\alpha_0 t}$ and radius \sqrt{t} , for some positive random variable H .

Compared to the previous result (Theorem 15), the growth is only very slightly slower: a single impulsion at the site 0 produces a growth in $t^{-d/2}e^{\alpha_0 t}$, while we have $e^{\alpha_0 t}$ when all the sites are regularly excited (as is e.g. the case when $\mu_i = 1$ for all i).

Finally, it is important to note that, even if the growth “near 0” of the process is very fast (exponential), the spatial propagation is quite slow (of order \sqrt{t}).

We start with some preliminary computations.

Lemma 21. *Adopt the notation and assumptions of Theorem 20. Introduce also, for $i \in \mathbb{Z}^d$ and $t \geq 0$, $\Gamma(i, t) = \sum_{n \geq 1} A^n(0, i)\varphi^{*n}(t)$.*

(i) *For all $n \geq 1$, all $t \geq 0$, $\varphi^{*n}(t) = (at)^{n-1}e^{-bt}/(n-1)!$.*

(ii) *For all $t \geq 0$, $\sum_{i \in \mathbb{Z}^d} \Gamma(i, t) = e^{\alpha_0 t}$.*

(iii) There is C such that for all $t \geq 0$, $\sum_{i \in \mathbb{Z}^d} |i|^2 \Gamma(i, t) \leq C(1+t)e^{\alpha_0 t}$.

(iv) There are some constants C and $t_0 > 0$ such that for all $t \geq t_0$, all $i \in \mathbb{Z}^d$,

$$(28) \quad |\Gamma(i, t) - p_{at}(i)e^{\alpha_0 t}| \leq \frac{Ce^{\alpha_0 t}}{t^{d/2+1/3}},$$

$$(29) \quad \left| \int_0^t \Gamma(i, s) ds - \frac{1}{\alpha_0} p_{at}(i)e^{\alpha_0 t} \right| \leq \frac{Ce^{\alpha_0 t}}{t^{d/2+1/3}}$$

Proof. Point (i) is well-known and can be checked recursively. Using that A^n is a stochastic matrix, we see that $\sum_{i \in \mathbb{Z}^d} \Gamma(i, t) = \sum_{n \geq 1} \varphi^{*n}(t)$. Hence (ii) follows from (i).

Next, we recall that A is the transition matrix of a symmetric random walk on \mathbb{Z}^d (with bounded jumps), so that there is a constant C such that for all $n \geq 0$, $\sum_{i \in \mathbb{Z}^d} |i|^2 A^n(0, i) \leq Cn$ (its variance at time n is of order n). Consequently,

$$\sum_{i \in \mathbb{Z}^d} |i|^2 \Gamma(i, t) \leq C \sum_{n \geq 1} n \varphi^{*n}(t) = C \sum_{n \geq 1} \frac{n(at)^{n-1} e^{-bt}}{(n-1)!} = C \left[\sum_{n \geq 1} \frac{(at)^{n-1} e^{-bt}}{(n-1)!} + \sum_{n \geq 2} \frac{(at)^{n-1} e^{-bt}}{(n-2)!} \right].$$

This is easily computed: it gives $Ce^{\alpha_0 t}[1 + at]$.

Point (iv) is more complicated. First, we need the Gaussian approximation of $A^n(0, i)$ given by (20). We will also need the following result, which can be found e.g. in [13, Lemma 9-(d)] (plus the fact that for $x \in (0, 1)$, $(x+1) \log(x+1) - x \geq x^2/4$): for any $\lambda > 0$, for X a Poisson(λ)-distributed random variable, for any $x \in (0, 1)$,

$$(30) \quad \Pr(|X - \lambda| \geq \lambda x) \leq 2 \exp(-\lambda x^2/4).$$

We now turn to our problem. Observing that $\sum_{n \geq 1} \varphi^{*n}(t) = e^{\alpha_0 t}$, we write

$$\Delta(i, t) = |\Gamma(i, t) - p_{at}(i)e^{\alpha_0 t}| = \left| \sum_{n \geq 1} \varphi^{*n}(t)(A^n(0, i) - p_{at}(i)) \right|.$$

We now assume that t is large enough so that $p_{t/a}(i) \leq 1$ for all i and we write

$$\begin{aligned} \Delta(i, t) &\leq \sum_{n \geq 1} \varphi^{*n}(t) \mathbf{1}_{\{|(n-1)-at| > (at)^{2/3}\}} + \sum_{n \geq 1} \varphi^{*n}(t) \mathbf{1}_{\{|(n-1)-at| \leq (at)^{2/3}\}} |A^n(0, i) - p_n(i)| \\ &\quad + \sum_{n \geq 1} \varphi^{*n}(t) \mathbf{1}_{\{|(n-1)-at| \leq (at)^{2/3}\}} |p_n(i) - p_{at}(i)| \\ &= \Delta_1(i, t) + \Delta_2(i, t) + \Delta_3(i, t). \end{aligned}$$

First, using point (i) and (30) (with $\lambda = at$ and $x = (at)^{-1/3}$), we see that (if t is large enough so that $at > 1$)

$$\Delta_1(i, t) \leq e^{\alpha_0 t} \sum_{n \geq 1} e^{-at} \frac{(at)^{n-1}}{(n-1)!} \mathbf{1}_{\{|(n-1)-at| > (at)^{2/3}\}} \leq 2e^{\alpha_0 t} e^{-(at)^{1/3}/4}.$$

We next use (20) and assume that t is large enough so that $|(n-1) - at| \leq (at)^{2/3}$ implies $n \geq at/2$:

$$\Delta_2(i, t) \leq C \sum_{n \geq 1} n^{-(d+2)/2} \varphi^{*n}(t) \mathbf{1}_{\{|(n-1)-at| \leq (at)^{2/3}\}} \leq C(at)^{-(d+2)/2} \sum_{n \geq 1} \varphi^{*n}(t) \leq Ct^{-(d+2)/2} e^{\alpha_0 t}.$$

Finally, we observe that $|\partial_t p_t(x)| \leq Ct^{-d/2-1}$, so that, if t is sufficiently large, $|(n-1)-at| \leq (at)^{2/3}$ implies $|p_n(i) - p_{at}(i)| \leq Ct^{-d/2-1/3}$. Consequently,

$$\Delta_3(i, t) \leq Ct^{-d/2-1/3} \sum_{n \geq 1} \varphi^{*n}(t) \leq Ct^{-d/2-1/3} e^{\alpha_0 t}.$$

We have proved that there are C and t_0 such that for all $t \geq t_0$, all $i \in \mathbb{Z}^d$,

$$|\Gamma(i, t) - p_{at}(i)e^{\alpha_0 t}| \leq 2e^{\alpha_0 t} e^{-(at)^{1/3}/4} + Ct^{-(d+2)/2} e^{\alpha_0 t} + Ct^{-d/2-1/3} e^{\alpha_0 t} \leq Ct^{-d/2-1/3} e^{\alpha_0 t},$$

which is (28).

It remains to deduce (29) from (28). We write

$$\begin{aligned} \delta(t, i) &= \left| \int_0^t \Gamma(i, s) ds - p_{at}(i) e^{\alpha_0 t} / \alpha_0 \right| \\ &\leq \int_0^{t-t^{1/2}} \Gamma(i, s) ds + \int_{t-t^{1/2}}^t |\Gamma(i, s) - p_{as}(i) e^{\alpha_0 s}| ds \\ &\quad + \int_{t-t^{1/2}}^t |p_{as}(i) e^{\alpha_0 s} - p_{at}(i) e^{\alpha_0 s}| ds + p_{at}(i) \left| \int_{t-t^{1/2}}^t e^{\alpha_0 s} ds - e^{\alpha_0 t} / \alpha_0 \right| \\ &= \delta_1(t, i) + \delta_2(t, i) + \delta_3(t, i) + \delta_4(t, i). \end{aligned}$$

First, point (ii) implies that

$$\delta_1(t, i) \leq \int_0^{t-t^{1/2}} e^{\alpha_0 s} ds \leq C e^{\alpha_0(t-t^{1/2})}.$$

Next, (28) tells us, if t is sufficiently large (so that $t - t^{1/2} \geq t_0$ and $t - t^{1/2} \geq t/2$), that

$$\delta_2(t, i) \leq C \int_{t-t^{1/2}}^t s^{-d/2-1/3} e^{\alpha_0 s} ds \leq Ct^{-d/2-1/3} e^{\alpha_0 t}.$$

Recalling that $|\partial_t p_t(x)| \leq Ct^{-d/2-1}$, we get (still for t large enough so that $t - t^{1/2} \geq t/2$)

$$\delta_3(t, i) \leq C \int_{t-t^{1/2}}^t s^{-d/2-1} (t-s) e^{\alpha_0 s} ds \leq Ct^{-d/2-1/2} e^{\alpha_0 t}.$$

Finally, if t is sufficiently large, we can bound $p_{at}(i)$ by 1 (for all i), whence

$$\delta_4(t, i) \leq \alpha_0^{-1} |e^{\alpha_0 t} - e^{\alpha_0(t-t^{1/2})} - e^{\alpha_0 t}| \leq C e^{\alpha_0(t-t^{1/2})}.$$

All in all, we have proved that for all t large enough, all $i \in \mathbb{Z}^d$,

$$\left| \int_0^t \Gamma(i, s) ds - p_{at}(i) e^{\alpha_0 t} / \alpha_0 \right| \leq C e^{\alpha_0(t-t^{1/2})} + Ct^{-d/2-1/3} e^{\alpha_0 t} + Ct^{-d/2-1/2} e^{\alpha_0 t} \leq Ct^{-d/2-1/3} e^{\alpha_0 t}$$

as desired. \square

We finally can give the

Proof of Theorem 20. We divide the proof in several steps.

Step 1. As usual, we write

$$Z_t^i = \int_0^t \int_0^\infty \mathbf{1}_{\{z \leq (2d+1)^{-1} \sum_{j \rightarrow i} [\int_0^{s^-} \varphi(s-u) dZ_u^j + \varphi(s) \mathbf{1}_{\{j=0\}}]\}} \pi^i(ds dz),$$

we set $m_t^i = \mathbb{E}[Z_t^i]$ and $\mathbf{m}_t = (m_t^i)_{i \in \mathbb{Z}^d}$. A simple computation, using Lemma 22, shows that

$$m_t^i = \int_0^t (2d+1)^{-1} \sum_{j \rightarrow i} \varphi(t-s) m_s^j ds + \mathbf{1}_{\{0 \rightarrow i\}} (2d+1)^{-1} \int_0^t \varphi(s) ds.$$

Using the vector formalism and introducing $\boldsymbol{\delta} = (\delta_i)_{i \in \mathbb{Z}^d}$ defined by $\delta_i = \mathbf{1}_{\{i=0\}}$, this rewrites $\mathbf{m}_t = (A\boldsymbol{\delta}) \int_0^t \varphi(s) ds + \int_0^t \varphi(t-s) (A\mathbf{m}_s) ds$. Differentiating this formula (see Lemma 27 for the justification of a very similar differentiation), we find $\mathbf{m}'_t = (A\boldsymbol{\delta})\varphi(t) + \int_0^t \varphi(t-s) A\mathbf{m}'_s ds$, which can be solved as (see Lemma 27 again) $\mathbf{m}'_t = \sum_{n \geq 1} \varphi^{*n}(t) A^n \boldsymbol{\delta}$. Hence for all $i \in \mathbb{Z}^d$,

$$(31) \quad (m_t^i)' = \sum_{n \geq 1} A^n(0, i) \varphi^{*n}(t).$$

We introduce the martingales, for $i \in \mathbb{Z}^d$, (we use a tilde for compensation),

$$M_t^i = \int_0^t \int_0^\infty \mathbf{1}_{\{z \leq (2d+1)^{-1} \sum_{j \rightarrow i} [\int_0^{s^-} \varphi(s-u) dZ_u^j + \varphi(s) \mathbf{1}_{\{j=0\}}]\}} \tilde{\pi}^i(ds dz)$$

and observe as usual that $[M^i, M^j]_t = 0$ when $i \neq j$ (because these martingales a.s. never jump at the same time) while $[M^i, M^i]_t = Z_t^i$. We will use several times that for any family $(\alpha_i)_{i \in \mathbb{Z}^d}$,

$$(32) \quad \mathbb{E}\left[\left(\sum_{i \in \mathbb{Z}^d} \alpha_i M_t^i\right)^2\right] = \sum_{i \in \mathbb{Z}^d} \alpha_i^2 m_t^i.$$

We finally introduce $U_t^i = Z_t^i - m_t^i$, the vectors $\mathbf{U}_t = (U_t^i)_{i \in \mathbb{Z}^d}$ and $\mathbf{M}_t = (M_t^i)_{i \in \mathbb{Z}^d}$ and observe, exactly as in the proof of Theorem 14-Step 1, that $\mathbf{U}_t = \mathbf{M}_t + A \int_0^t \varphi(t-s) \mathbf{U}_s ds$, whence $\mathbf{U}_t = \left(\mathbf{M}_t + \sum_{n \geq 1} A^n \int_0^t \varphi^{*n}(t-s) \mathbf{M}_s ds\right)$ and thus, for all $i \in \mathbb{Z}^d$,

$$(33) \quad Z_t^i = m_t^i + M_t^i + \sum_{j \in \mathbb{Z}^d} \sum_{n \geq 1} \int_0^t \varphi^{*n}(t-s) A^n(i, j) M_s^j ds = m_t^i + M_t^i + W_t^i,$$

the last equality defining W_t^i .

Step 2. Here we treat the terms m_t^i , M_t^i , and collect a few more information. Recall that $\Gamma(t, i) = \sum_{n \geq 1} A^n(0, i) \varphi^{*n}(t)$. Starting from (31), we see that $m_t^i = \int_0^t \Gamma(i, s) ds$ and deduce from Lemma 21-(iv) that there are C and $t_0 \geq 0$ such that for all $i \in \mathbb{Z}^d$, all $t \geq t_0$,

$$\left| m_t^i - \frac{1}{\alpha_0} p_{at}(i) e^{\alpha_0 t} \right| \leq \frac{C e^{\alpha_0 t}}{t^{d/2+1/3}}.$$

Next, Lemma 21-(ii)-(iii) imply that $\sum_{i \in \mathbb{Z}^d} m_t^i = \int_0^t e^{\alpha_0 s} ds \leq C e^{\alpha_0 t}$ and $\sum_{i \in \mathbb{Z}^d} |i|^2 m_t^i \leq C \int_0^t (1+s) e^{\alpha_0 s} ds \leq C(1+t) e^{\alpha_0 t}$. Finally, we observe that

$$\mathbb{E}[|M_t^i|] \leq (m_t^i)^{1/2} \leq \left(\sum_{j \in \mathbb{Z}^d} m_t^j \right)^{1/2} \leq C e^{\alpha_0 t/2}.$$

Step 3. We introduce

$$X = \sum_{j \in \mathbb{Z}^d} \int_0^\infty e^{-\alpha_0 s} M_s^j ds$$

and show that there are $C > 0$ and $t_0 > 0$ such that for all $i \in \mathbb{Z}^d$, all $t \geq t_0$,

$$\mathbb{E}[|W_t^i - p_{at}(i) e^{\alpha_0 t} X|] \leq \frac{C e^{\alpha_0 t}}{t^{d/2+1/3}}.$$

We observe that since $A^n(i, j) = A^n(0, i - j)$, it holds that $W_t^i = \sum_{j \in \mathbb{Z}^d} \int_0^t \Gamma(i - j, t - s) M_s^j ds$. We also introduce the auxiliary processes

$$\begin{aligned}\overline{W}_t^i &= \sum_{j \in \mathbb{Z}^d} \int_0^{t^{1/2}} \Gamma(i - j, t - s) M_s^j ds, \\ \widetilde{W}_t^i &= \sum_{j \in \mathbb{Z}^d} \int_0^{t^{1/2}} p_{a(t-s)}(i - j) e^{\alpha_0(t-s)} M_s^j ds, \\ \widehat{W}_t^i &= \sum_{j \in \mathbb{Z}^d} \int_0^{t^{1/2}} p_{at}(i) e^{\alpha_0(t-s)} M_s^j ds.\end{aligned}$$

Step 3.1. Here we show that $\mathbb{E}[|W_t^i - \overline{W}_t^i|] \leq C \exp(\alpha_0 t - (\alpha_0/2)t^{1/2})$. By definition of Γ and using (32),

$$\begin{aligned}\mathbb{E}[|W_t^i - \overline{W}_t^i|] &\leq \int_{t^{1/2}}^t \sum_{n \geq 1} \varphi^{*n}(t - s) \mathbb{E}\left[\left|\sum_{j \in \mathbb{Z}^d} A^n(i, j) M_s^j\right|\right] ds \\ &\leq \int_{t^{1/2}}^t \sum_{n \geq 1} \varphi^{*n}(t - s) \left(\sum_{j \in \mathbb{Z}^d} (A^n(i, j))^2 m_s^j\right)^{1/2} ds.\end{aligned}$$

Using that $A^n(i, j)$ is bounded by 1 and that $\sum_{i \in \mathbb{Z}^d} m_t^i \leq C e^{\alpha_0 t}$ (see Step 2), we see that $\sum_{j \in \mathbb{Z}^d} (A^n(i, j))^2 m_s^j \leq C e^{\alpha_0 s/2}$. Next, the explicit expression of φ^{*n} (see Lemma 21-(i)) gives $\sum_{n \geq 1} \varphi^{*n}(t - s) = e^{\alpha_0(t-s)}$. We finally find

$$\mathbb{E}[|W_t^i - \overline{W}_t^i|] \leq C \int_{t^{1/2}}^t e^{\alpha_0(t-s)} e^{\alpha_0 s/2} ds \leq C e^{\alpha_0 t} e^{-\alpha_0 t^{1/2}/2}.$$

Step 3.2. We next check that $\mathbb{E}[|\overline{W}_t^i - \widetilde{W}_t^i|] \leq C t^{-d/2-1/3} e^{\alpha_0 t}$. Using (32), we get

$$\mathbb{E}[|\overline{W}_t^i - \widetilde{W}_t^i|] \leq \int_0^{t^{1/2}} \left[\sum_{j \in \mathbb{Z}^d} m_s^j (\Gamma(i - j, t - s) - p_{a(t-s)}(i - j) e^{\alpha_0(t-s)})^2 \right]^{1/2} ds.$$

Using Lemma 21-(iv), $|\Gamma(i - j, t - s) - p_{a(t-s)}(i - j) e^{\alpha_0(t-s)}| \leq C(t - s)^{-d/2-1/3} e^{\alpha_0(t-s)}$ if $t - s$ is large enough (which is the case for all $s \in [0, t^{1/2}]$ if t is large enough). Since furthermore $\sum_{i \in \mathbb{Z}^d} m_t^i \leq C e^{\alpha_0 t}$ (see Step 2), we find

$$\mathbb{E}[|\overline{W}_t^i - \widetilde{W}_t^i|] \leq C e^{\alpha_0 t} \int_0^{t^{1/2}} (t - s)^{-d/2-1/3} e^{-\alpha_0 s/2} ds.$$

For t large enough, we clearly have $(t - s)^{-d/2-1/3} \leq 2t^{-d/2-1/3}$ for all $s \in [0, t^{1/2}]$, whence

$$\mathbb{E}[|\overline{W}_t^i - \widetilde{W}_t^i|] \leq C t^{-d/2-1/3} e^{\alpha_0 t} \int_0^{t^{1/2}} e^{-\alpha_0 s/2} ds \leq C t^{-d/2-1/3} e^{\alpha_0 t}.$$

Step 3.3. We now prove that $\mathbb{E}[|\widetilde{W}_t^i - \widehat{W}_t^i|] \leq C t^{-d/2-1/2} e^{\alpha_0 t}$. As usual, we start with

$$\mathbb{E}[|\widetilde{W}_t^i - \widehat{W}_t^i|] \leq \int_0^{t^{1/2}} \left(\sum_{j \in \mathbb{Z}^d} m_s^j (p_{a(t-s)}(i - j) - p_{at}(i))^2 \right)^{1/2} e^{\alpha_0(t-s)} ds.$$

But an easy computation (using that $|\partial_t p_t(x)| \leq Ct^{-d/2-1}$ and $|\nabla_x p_t(x)| \leq Ct^{-d/2-1/2}$) shows that for all $t > 0$, all $h \in (0, t/2)$, all $x, y \in \mathbb{R}^d$, $|p_{t-h}(x-y) - p_t(x)| \leq Ch t^{-d/2-1} + |y| t^{-d/2-1/2}$. Hence if t is large enough so that $t - t^{1/2} \geq t/2$, we can write

$$\begin{aligned} \mathbb{E}[|\widetilde{W}_t^i - \widehat{W}_t^i|] &\leq C \int_0^{t^{1/2}} \left(\sum_{j \in \mathbb{Z}^d} m_s^j (st^{-d/2-1} + |j|t^{-d/2-1/2})^2 \right)^{1/2} e^{\alpha_0(t-s)} ds \\ &\leq Ct^{-d/2-1/2} \int_0^{t^{1/2}} \left(\sum_{j \in \mathbb{Z}^d} m_s^j \right)^{1/2} e^{\alpha_0(t-s)} ds \\ &\quad + Ct^{-d/2-1/2} \int_0^{t^{1/2}} \left(\sum_{j \in \mathbb{Z}^d} |j|^2 m_s^j \right)^{1/2} e^{\alpha_0(t-s)} ds. \end{aligned}$$

Finally, we know from Step 2 that $\sum_{j \in \mathbb{Z}^d} m_s^j + \sum_{j \in \mathbb{Z}^d} |j|^2 m_s^j \leq C(1+s)e^{\alpha_0 s}$, whence

$$\mathbb{E}[|\widetilde{W}_t^i - \widehat{W}_t^i|] \leq Ct^{-d/2-1/2} \int_0^{t^{1/2}} (1+s)^{1/2} e^{\alpha_0 s/2} e^{\alpha_0(t-s)} ds \leq Ct^{-d/2-1/2} e^{\alpha_0 t}.$$

Step 3.4 We finally verify that $\mathbb{E}[|\widehat{W}_t^i - p_{at}(i)e^{\alpha_0 t} X|] \leq Ce^{\alpha_0 t - (\alpha_0/2)t^{1/2}}$. We note that

$$\widehat{W}_t^i - p_{at}(i)e^{\alpha_0 t} X = p_{at}(i)e^{\alpha_0 t} \sum_{j \in \mathbb{Z}^d} \int_{t^{1/2}}^t e^{-\alpha_0 s} M_s^j ds.$$

For t large enough (not depending on i), we can bound $p_{at}(i)$ by 1. Hence, we infer from (32) and the fact that $\sum_{j \in \mathbb{Z}^d} m_s^j \leq Ce^{\alpha_0 s}$ that

$$\mathbb{E}[|\widehat{W}_t^i - p_{at}(i)e^{\alpha_0 t} X|] \leq Ce^{\alpha_0 t} \int_{t^{1/2}}^t e^{-\alpha_0 s} \left(\sum_{j \in \mathbb{Z}^d} m_s^j \right)^{1/2} ds \leq Ce^{\alpha_0 t} \int_{t^{1/2}}^\infty e^{-\alpha_0 s/2} ds,$$

from which the conclusion follows.

Step 3.5. Gathering Steps 3.1 to 3.4, we conclude that indeed, there are $C > 0$ and $t_0 > 0$ such that for all $t \geq t_0$, all $i \in \mathbb{Z}^d$, $\mathbb{E}[|W_t^i - p_{at}(i)e^{\alpha_0 t} X|] \leq Ce^{\alpha_0 t} t^{-d/2-1/3}$.

Step 4. Define the random variable $H = \alpha_0^{-1} + X$. Recall (33) and write $Z_t^i - p_{at}(i)e^{\alpha_0 t} H = [m_t^i - \alpha_0^{-1} p_{at}(i)e^{\alpha_0 t}] + M_t^i + [W_t^i - p_{at}(i)e^{\alpha_0 t} X]$. Gathering Steps 2 and 3, we see that there are $C > 0$ and $t_0 > 0$ such that for all $t \geq t_0$, all $i \in \mathbb{Z}^d$,

$$\mathbb{E}[|Z_t^i - p_{at}(i)e^{\alpha_0 t} H|] \leq \frac{Ce^{\alpha_0 t}}{t^{d/2+1/3}} + Ce^{\alpha_0 t/2} \leq \frac{Ce^{\alpha_0 t}}{t^{d/2+1/3}}.$$

To prove that H is nonnegative, it suffices to use the above inequality with $i = 0$, divided by $p_{at}(0)e^{\alpha_0 t}$. Recalling that $p_{at}(0) = ct^{-d/2}$ for some constant c , we deduce that $\mathbb{E}[|H - Z_t^0 e^{-\alpha_0 t}/p_{at}(0)|] \leq Ct^{-1/3}$. Consequently, H is the limit (in L^1) of $Z_t^0 e^{-\alpha_0 t}/p_{at}(0)$, and is thus nonnegative. This ends the proof of (i).

Step 5. We now check (ii), which follows from (i): for $x \in \mathbb{R}^d$, and $t \geq t_0$,

$$\begin{aligned} \mathbb{E}[|t^{d/2} e^{-\alpha_0 t} Z_t^{\lfloor xt^{1/2} \rfloor} - H p_a(x)|] &\leq t^{d/2} e^{-\alpha_0 t} \mathbb{E}[|Z_t^{\lfloor xt^{1/2} \rfloor} - H e^{\alpha_0 t} p_{at}(\lfloor xt^{1/2} \rfloor)|] \\ &\quad + \mathbb{E}[|H|] |t^{d/2} p_{at}(\lfloor xt^{1/2} \rfloor) - p_a(x)|. \end{aligned}$$

The first term on the RHS is bounded, by (i), by $Ct^{-1/3}$, which tends to 0 as $t \rightarrow \infty$. The second term also tends to 0, simply because

$$t^{d/2} p_{at}(\lfloor xt^{1/2} \rfloor) = p_a(t^{-1/2} \lfloor xt^{1/2} \rfloor) \rightarrow p_a(x) \quad \text{as } t \rightarrow \infty.$$

Step 6. It remains to prove (iii). First note that $H = \lim_{t \rightarrow \infty} e^{-\alpha_0 t} Z_t$ in L^1 , where $Z_t = \sum_{i \in \mathbb{Z}^d} Z_t^i$. Indeed, recalling (33)

$$Z_t = \sum_{i \in \mathbb{Z}^d} m_t^i + \sum_{i \in \mathbb{Z}^d} M_t^i + \sum_{i \in \mathbb{Z}^d} W_t^i.$$

We have $\sum_{i \in \mathbb{Z}^d} m_t^i = \int_0^t e^{\alpha_0 s} ds = \alpha_0^{-1} [e^{\alpha_0 t} - 1]$ by Step 2, (32) implies that $\mathbb{E}[(\sum_{i \in \mathbb{Z}^d} M_t^i)^2] = \sum_{i \in \mathbb{Z}^d} m_t^i$ and finally $\sum_{i \in \mathbb{Z}^d} W_t^i = e^{\alpha_0 t} \int_0^t e^{-\alpha_0 s} \sum_{i \in \mathbb{Z}^d} M_s^i ds$, therefore $e^{-\alpha_0 t} Z_t$ converges to $\alpha_0^{-1} + 0 + X = H$ as $t \rightarrow \infty$ in L^1 . We also note that $\mathbb{E}[H] = \alpha_0^{-1} > 0$ since $\mathbb{E}[X] = 0$.

Next, we recall that by Theorem 19-(i), $(Z_t)_{t \geq 0}$ is a scalar impulsion Hawkes process: its compensator is given by $A_t = \int_0^t [\varphi(s) + \int_0^{s-} \varphi(s-u) dZ_u] ds$. We claim that $(Z_t)_{t \geq 0}$ has the same law as $(\tilde{Z}_t)_{t \geq 0}$ built as follows:

- consider a Poisson process $(N_t)_{t \geq 0}$ with intensity $\varphi(t) dt$, observe that N_∞ is Poisson(Λ)-distributed, denote by $0 < T_1 < \dots < T_{N_\infty}$ its times of jump (we adopt the convention that $T_i = \infty$ for $i > N_\infty$),
- consider an i.i.d. family $(\tilde{Z}_t^k)_{t \geq 0}$ of scalar impulsion Hawkes process with same law as $(Z_t)_{t \geq 0}$,
- put $\tilde{Z}_t = N_t + \sum_{i=1}^{N_\infty} \tilde{Z}_{t-T_k}^k \mathbf{1}_{\{t \geq T_k\}}$.

Indeed, $(\tilde{Z}_t)_{t \geq 0}$ is a counting process with compensator

$$\int_0^t [\varphi(s) + \sum_{i \geq 1} \mathbf{1}_{\{s > T_k\}} (\varphi(s-T_k) + \int_0^{(s-T_k)-} \varphi(s-T_k-u) d\tilde{Z}_u^k)] ds = \int_0^t [\varphi(s) + \int_0^{s-} \varphi(s-u) d\tilde{Z}_u] ds.$$

We define $\tilde{H} = \lim_{t \rightarrow \infty} e^{-\alpha_0 t} \tilde{Z}_t$ and, for each $k \geq 1$, $\tilde{H}_k = \lim_{t \rightarrow \infty} e^{-\alpha_0 t} \tilde{Z}_t^k$. We obviously have $\tilde{H} = \sum_{k=1}^{N_\infty} e^{-\alpha_0 T_k} \tilde{H}_k$. Denoting by $p = \Pr(H = 0)$ (which also equals $\Pr(\tilde{H} = 0)$ and $\Pr(\tilde{H}_k = 0)$ for all $k \geq 1$), we deduce by independence of the family $(\tilde{H}_k)_{k \geq 1}$, that $p = \sum_{n \geq 1} \Pr(N_\infty = n) p^n$. Hence $p = \sum_{n \geq 1} e^{-\Lambda} \Lambda^n p^n / n! = e^{-\Lambda(1-p)}$. Since $\mathbb{E}[H] > 0$, we cannot have $p = 1$. Hence p is the unique solution in $(0, 1)$ to $p = e^{-\Lambda(1-p)}$. Recalling that $\Pr(\Omega_e) = \exp(-\gamma_\Lambda \Lambda)$ where $\gamma_\Lambda \in (0, 1)$ is characterised by $\gamma_\Lambda \Lambda + \log(1 - \gamma_\Lambda) = 0$ by Theorem 19-2-(ii), we easily check that $p = \Pr(\Omega_e)$.

By definition of H , we have $\Omega_e \subset \{H = 0\}$. Since $\Pr(\Omega_e) = \Pr(H = 0)$ we conclude that a.s., $H > 0$ on Ω_e^c . \square

5. APPENDIX: CONVOLUTION EQUATIONS

We collect here some technical results about convolution equations. We start with an identity of constant use in the paper.

Lemma 22. *Let $\phi : [0, \infty) \mapsto \mathbb{R}$ be locally integrable and let $\alpha : [0, \infty) \mapsto \mathbb{R}$ have finite variations on compact intervals and satisfy $\alpha(0) = 0$. Then for all $t \geq 0$,*

$$\int_0^t \int_0^{s-} \phi(s-u) d\alpha(u) ds = \int_0^t \int_0^s \phi(s-u) d\alpha(u) ds = \int_0^t \phi(t-s) \alpha(s) ds.$$

Proof. First, we clearly have that $\int_0^{s^-} \phi(s-u)d\alpha(u) = \int_0^s \phi(s-u)d\alpha(u)$ for almost every $s \geq 0$, whence $\int_0^t \int_0^{s^-} \phi(s-u)d\alpha(u)ds = \int_0^t \int_0^s \phi(s-u)d\alpha(u)ds$. Using twice the Fubini theorem,

$$\begin{aligned} \int_0^t \left(\int_0^s \phi(s-u)d\alpha(u) \right) ds &= \int_0^t \left(\int_u^t \phi(s-u)ds \right) d\alpha(u) \\ &= \int_0^t \left(\int_0^{t-u} \phi(v)dv \right) d\alpha(u) \\ &= \int_0^t \left(\int_0^{t-v} d\alpha(u) \right) \phi(v)dv \\ &= \int_0^t \alpha(t-v)\phi(v)dv, \end{aligned}$$

from which the conclusion follows, using the substitution $s = t - v$. \square

We carry on with a generalized Grönwall-Picard lemma, which is more or less standard.

Lemma 23. *Let $\phi : [0, \infty) \mapsto [0, \infty)$ be locally integrable and $g : [0, \infty) \mapsto [0, \infty)$ be locally bounded.*

(i) *Consider a locally bounded nonnegative function u such that for all $t \geq 0$, $u_t \leq g_t + \int_0^t \phi(t-s)u_s ds$ for all $t \geq 0$. Then $\sup_{[0, T]} u_t \leq C_T \sup_{[0, T]} g_t$, for some constant C_T depending only on $T > 0$ and ϕ .*

(ii) *Consider a sequence of locally bounded nonnegative functions u^n such that for all $t \geq 0$, all $n \geq 0$, $u_t^{n+1} \leq \int_0^t \phi(t-s)u_s^n ds$. Then $\sup_{[0, T]} \sum_{n \geq 0} u_t^n \leq C_T$, for some constant C_T depending only on $T > 0$, u^0 and ϕ .*

(iii) *Consider a sequence of locally bounded nonnegative functions u^n such that for all $t \geq 0$, all $n \geq 0$, $u_t^{n+1} \leq g_t + \int_0^t \phi(t-s)u_s^n ds$. Then for all $T \geq 0$, $\sup_{[0, T]} \sup_{n \geq 0} u_t^n \leq C_T$, for some constant C_T depending only on $T > 0$, u^0 , g and ϕ .*

Proof. We start with point (i). Fix $T > 0$ and consider $A > 0$ such that $\int_0^T \phi(s)\mathbf{1}_{\{\phi(s) \geq A\}} ds \leq 1/2$. Then for all $t \in [0, T]$,

$$u_t \leq g_t + \int_0^t \mathbf{1}_{\{\phi(t-s) \leq A\}} \phi(t-s)u_s ds + \int_0^t \mathbf{1}_{\{\phi(t-s) > A\}} \phi(t-s)u_s ds \leq g_t + A \int_0^t u_s ds + \sup_{[0, t]} u_s / 2.$$

from which we deduce that $\sup_{[0, t]} u_s \leq 2 \sup_{[0, t]} g_s + 2A \int_0^t u_s ds$. We then can apply the standard Grönwall Lemma to get $\sup_{[0, T]} u_s \leq 2(\sup_{[0, T]} g_s)e^{2AT}$.

To check point (iii), put $v_t^n = \sup_{k=0, \dots, n} u_t^k$. One easily checks that for all $n \geq 0$, $v_t^n \leq u_t^0 + g_t + \int_0^t \phi(t-s)v_s^n ds$. By point (i), $\sup_{[0, T]} v_t^n \leq C_T \sup_{[0, T]} (g_t + u_t^0)$. Letting n increase to infinity concludes the proof.

Point (ii) follows from point (iii), since $v_t^n = \sum_{k=0}^n u_t^k$ satisfies $v_t^{n+1} \leq u_t^0 + \int_0^t \phi(t-s)v_s^n ds$. \square

We next prove an easy well-posedness result for a general convolution equation.

Lemma 24. *Let $h : \mathbb{R} \mapsto [0, \infty)$ be Lipschitz-continuous and $\varphi : [0, \infty) \mapsto \mathbb{R}$ be locally integrable. The equation*

$$(34) \quad m_t = \int_0^t h \left(\int_0^s \varphi(s-u)dm_u \right) ds$$

has a unique non-decreasing locally bounded solution. Furthermore, m is of class C^1 on $[0, \infty)$.

If $h(x) = \mu + x$ for some $\mu > 0$ and if φ is nonnegative, Equation (34) rewrites as

$$(35) \quad m_t = \mu t + \int_0^t \varphi(t-s)m_s ds.$$

Proof. Let m and \tilde{m} be two such solutions. Since h is Lipschitz-continuous,

$$v_t = \int_0^t |d(m_u - \tilde{m}_u)| \leq C \int_0^t ds \int_0^s |\varphi(s-u)| |d(m_u - \tilde{m}_u)| = C \int_0^t |\varphi(t-u)| v_u du.$$

The last inequality follows from Lemma 22. Lemma 23-(i) allows us to conclude that $v_t = 0$ for all $t \geq 0$ (because $v_t \leq m_t + \tilde{m}_t$ is locally bounded), whence $m_t = \tilde{m}_t$ for all $t \geq 0$. For the existence, we consider the sequence of non-decreasing functions $m_t^0 = 0$ and $m_t^{n+1} = \int_0^t h(\int_0^s \varphi(s-u) dm_u^n) ds$ for every $n \geq 0$. We easily check that $m_t^{n+1} \leq h(0)t + |h|_{lip} \int_0^t |\varphi(t-u)| m_u^n du$, so that $\sup_{n \geq 0} m_t^n$ is locally bounded by Lemma 23-(iii). Setting, for $n \geq 0$, $\delta_t^n = \int_0^t |d(m_u^{n+1} - m_u^n)|$, one readily gets $\delta_t^{n+1} \leq |h|_{lip} \int_0^t |\varphi(s-u)| \delta_u^n du$ for all $n \geq 0$. Lemma 23-(ii) thus implies that $\sum_{n \geq 0} \delta_t^n < \infty$. All this classically implies the existence of a locally bounded non-decreasing m such that for all $t \geq 0$, $\lim_n \int_0^t |d(m_u - m_u^n)| = 0$. Checking that m solves (34) is routine.

To prove that m is C^1 , we use the previous Picard Iteration. One easily sees, by induction, that m^n is C^1 for all $n \geq 0$ and that $(m_t^{n+1})' = h(\int_0^t \varphi(t-u)(m_u^n)' du)$ (indeed, $t \mapsto \int_0^t \varphi(t-u)(m_u^n)' du = \int_0^t \varphi(u)(m_{t-u}^n)' du$ is continuous because φ is locally integrable and because $(m_t^n)'$ is continuous by the inductive assumption). Next, a direct computation shows that $|(m_t^{n+1})' - (m_t^n)'| \leq C \int_0^t |\varphi(t-u)| |(m_u^n)' - (m_u^{n-1})'| du$. Using Lemma 23-(ii), we deduce that the sequence $(m^n)'$ is Cauchy (for the uniform convergence on compact time intervals). The conclusion classically follows.

The equivalence between (34) and (35) in the linear case directly follows from Lemma 22. \square

We now investigate the large-time behaviour of m_t in the linear case. We start with the subcritical case.

Lemma 25. *Consider $\mu > 0$ and a function $\varphi : (0, \infty) \mapsto [0, \infty)$ such that $\Lambda = \int_0^\infty \varphi(s) ds < 1$. By Lemma 24, (35) has a unique non-decreasing locally bounded solution $(m_t)_{t \geq 0}$, which is furthermore of class C^1 . It holds that $m_t' \sim a_0$ and $m_t \sim a_0 t$ as $t \rightarrow \infty$, where $a_0 = \mu/(1 - \Lambda) > 0$.*

Proof. We rather use (34), which writes $m_t = \mu t + \int_0^t \int_0^s \varphi(s-u) m_u' du ds$. Differentiating this expression, we find $m_t' = \mu + \int_0^t \varphi(t-s) m_s' ds$. We first introduce $u_t = \sup_{[0,t]} m_s'$. We have $u_t \leq \mu + \Lambda u_t$, whence $u_t \leq \mu/(1 - \Lambda)$ for all $t \geq 0$ and thus $\limsup_{t \rightarrow \infty} m_t' \leq \mu/(1 - \Lambda)$. We next introduce $v_t = \inf_{s \geq t} m_s'$, which is non-decreasing and thus has a limit $\ell \in (0, \infty]$. We have $v_t \geq \mu + v_{t/2} \int_0^{t/2} \varphi(s) ds \rightarrow \mu + \Lambda \ell$ as $t \rightarrow \infty$. Consequently $\ell \geq \mu + \Lambda \ell$, whence $\ell \geq \mu/(1 - \Lambda)$ and finally $\liminf_{t \rightarrow \infty} m_t' \geq \mu/(1 - \Lambda)$. All this proves that $m_t' \sim a_0$ and this implies that $m_t \sim a_0 t$. \square

We now turn to the supercritical case.

Lemma 26. *Consider $\mu > 0$ and a function $\varphi : [0, \infty) \mapsto [0, \infty)$ such that $\Lambda = \int_0^\infty \varphi(s) ds \in (1, \infty]$. By Lemma 24, (35) has a unique non-decreasing locally bounded solution $(m_t)_{t \geq 0}$, which is of class C^1 . Assume furthermore that $t \mapsto \int_0^t |d\varphi(s)|$ has at most polynomial growth. Let $\Gamma(t) = \sum_{n \geq 1} \varphi^{*n}(t)$ and $\Upsilon(t) = \int_0^t \Gamma(s) ds$.*

(a) There is a unique $\alpha_0 > 0$ such that $\mathcal{L}_\varphi(\alpha_0) = 1$. The function Γ is locally bounded. Setting $a_0 = \mu\alpha_0^{-2}(\int_0^\infty t\varphi(t)e^{-\alpha_0 t}dt)^{-1}$, we have, as $t \rightarrow \infty$,

$$\Gamma(t) \sim (a_0\alpha_0^2/\mu)e^{\alpha_0 t}, \quad \Upsilon(t) \sim (a_0\alpha_0/\mu)e^{\alpha_0 t}, \quad m_t \sim a_0e^{\alpha_0 t}, \quad m'_t \sim a_0\alpha_0e^{\alpha_0 t}.$$

(b) There are some constants $0 < c < C$ such that for all $t \geq 0$, $ce^{\alpha_0 t} \leq \Gamma(t) + 1 \leq Ce^{\alpha_0 t}$, $ce^{\alpha_0 t} \leq \Upsilon(t) + 1 \leq Ce^{\alpha_0 t}$, $ce^{\alpha_0 t} \leq m_t + 1 \leq Ce^{\alpha_0 t}$ and $ce^{\alpha_0 t} \leq m'_t + 1 \leq Ce^{\alpha_0 t}$.

(c) We also have

$$\lim_{t \rightarrow \infty} \int_0^t \left(\frac{\Upsilon(t-s)}{m_t} - \frac{\alpha_0}{\mu} e^{-\alpha_0 s} \right)^2 m'_s ds = 0.$$

(d) Consider a real sequence $(\eta_n)_{n \geq 1}$ such that $\lim_{n \rightarrow \infty} \eta_n = 0$. Then we have the property that $\lim_{t \rightarrow \infty} e^{-\alpha_0 t} \sum_{n \geq 1} \eta_n \int_0^t \varphi^{*n}(t-s)e^{\alpha_0 s/2} ds = 0$.

(e) For any pair of locally bounded functions $u, h : (0, \infty) \mapsto \mathbb{R}$ such that $u = h + u \star \varphi$, there holds $u = h + h \star \Gamma$.

Proof. We easily deduce from our assumptions on φ that there is some constants $C > 0$, $p > 0$ such that for all $t \geq 0$, $\varphi(t) \leq C(1+t)^p$ (in particular, φ is locally bounded). Hence its Laplace transform is clearly well-defined on $(0, \infty)$, of class C^∞ , and $\lim_{\alpha \rightarrow \infty} \mathcal{L}_\varphi(\alpha) = 0$. Furthermore, $\mathcal{L}_\varphi(0) = \int_0^\infty \varphi(t)dt \in (1, \infty]$. Hence, there indeed exists a unique $\alpha_0 > 0$ such that $\mathcal{L}_\varphi(\alpha_0) = 1$. We now divide the proof into several steps.

Step 1. We first prove that Γ is locally bounded. To this end, we introduce $\Gamma_n = \sum_{k=1}^n \varphi^{*k}(t)$ and observe that $\Gamma_{n+1} = \varphi + \Gamma_n \star \varphi$. Since φ is locally bounded, Lemma 23-(iii) allows us to conclude that $\sup_n \Gamma_n$ is locally bounded, whence the result.

Step 2. Here we prove (e). Since h is locally bounded and since φ is locally integrable, we easily deduce from Lemma 23-(i) that the equation $v = h + v \star \varphi$ (with unknown v) has at most one locally bounded solution. Since both u and $h + h \star \Gamma$ are locally bounded solutions, the conclusion follows.

Step 3. The aim of this step is to verify that $\Gamma(t) \sim (a_0\alpha_0^2/\mu)e^{\alpha_0 t}$ as $t \rightarrow \infty$. Observe that Γ solves $\Gamma = \varphi + \Gamma \star \varphi$ and introduce $u(t) = \Gamma(t)e^{-\alpha_0 t}$ and $f(t) = \varphi(t)e^{-\alpha_0 t}$. One easily checks that $u = f + u \star f$. We now apply Theorem 4 of Feller [12]. We have $\int_0^\infty f(t)dt = 1$ by definition of α_0 . We set $b_1 = \int_0^\infty tf(t)dt = \int_0^\infty t\varphi(t)e^{-\alpha_0 t}dt$ and $b_2 = \int_0^\infty t^2 f(t)dt = \int_0^\infty t^2 \varphi(t)e^{-\alpha_0 t}dt$, which clearly both converge, since $\varphi(t) \leq C(1+t)^p$. Finally, it is not difficult to check that $f(t)$, $tf(t)$ and $t^2 f(t)$ have a bounded total variation on $[0, \infty)$ since we have assumed that $t \mapsto \int_0^t |d\varphi(s)|$ has at most polynomial growth. Thus Feller [12, Theorem 4] tells us that $u(t) \rightarrow 1/b_1$ as $t \rightarrow \infty$, which gives $\Gamma(t) \sim e^{\alpha_0 t}/b_1$. This ends the proof, since $1/b_1 = a_0\alpha_0^2/\mu$ by definition of a_0 .

Step 4. We now conclude the proof of (a) and (b). Recall that $m_t = \mu t + \int_0^t \int_0^s \varphi(s-u)m'_u du ds$, so that $m' = \mu + \varphi \star m'$. Applying (e), we deduce that $m' = \mu + \mu \star \Gamma = \mu(1 + \Upsilon)$. By Step 3, we know that $\Gamma(t) \sim (a_0\alpha_0^2/\mu)e^{\alpha_0 t}$ as $t \rightarrow \infty$. This obviously implies that $\Upsilon(t) \sim (a_0\alpha_0/\mu)e^{\alpha_0 t}$, whence $m'_t \sim a_0\alpha_0e^{\alpha_0 t}$ and finally $m_t = \int_0^t m'_s ds \sim a_0e^{\alpha_0 t}$. Finally, (b) directly follows from (a) and the facts that Γ , Υ , m and m' are nonnegative and locally bounded.

Step 5. Point (d) is not very difficult: using that $\varphi(t) \leq C(1+t)^p$, we see that $\varphi^{*n}(t) \leq C_n(1+t)^{p_n}$ for some constants $C_n > 0$ and $p_n > 0$. We next introduce $\varepsilon_k = \sup_{n \geq k} |\eta_n|$, which

decreases to 0 as $k \rightarrow \infty$, and write, for any $k \geq 1$,

$$\begin{aligned} \limsup_{t \rightarrow \infty} e^{-\alpha_0 t} \sum_{n \geq 1} |\eta_n| \int_0^t \varphi^{*n}(t-s) e^{\alpha_0 s/2} ds &\leq \limsup_{t \rightarrow \infty} e^{-\alpha_0 t} \sum_{n=1}^k |\eta_n| \int_0^t C_n(t-s)^{p_n} e^{\alpha_0 s/2} ds \\ &+ \varepsilon_k \limsup_{t \rightarrow \infty} e^{-\alpha_0 t} \sum_{n=k+1}^{\infty} \int_0^t \varphi^{*n}(t-s) e^{\alpha_0 s/2} ds. \end{aligned}$$

The first term on the RHS is of course 0 (for any fixed k). We can bound the second one, using (b), by

$$\varepsilon_k \limsup_{t \rightarrow \infty} e^{-\alpha_0 t} \int_0^t \Gamma(t-s) e^{\alpha_0 s/2} ds \leq C \varepsilon_k \limsup_{t \rightarrow \infty} e^{-\alpha_0 t} \int_0^t e^{\alpha_0(t-s)} e^{\alpha_0 s/2} ds \leq C \varepsilon_k.$$

Letting k tend to infinity concludes the proof.

Step 6. It only remains to check point (c). We use the Lebesgue dominated convergence theorem. Define $h_s^t = (\Upsilon(t-s)/m_t - \alpha_0 e^{-\alpha_0 s}/\mu)^2 m_s' \mathbf{1}_{\{s \leq t\}}$. We have to prove that $\lim_{t \rightarrow \infty} \int_0^\infty h_s^t ds = 0$. First, it is obvious from (a) that for $s > 0$ fixed, $\lim_{t \rightarrow \infty} h_s^t = 0$. Next, we use (b) to write (for t large enough so that $m_t \geq c e^{\alpha_0 t}$) $h_s^t \leq C(e^{-\alpha_0 s})^2 e^{\alpha_0 s} \leq C e^{-\alpha_0 s}$, which does not depend on t and is integrable on $(0, \infty)$. \square

We next consider briefly a *vector* convolution equation.

Lemma 27. *Consider a family $\boldsymbol{\mu} = (\mu_i)_{i \in \mathbb{Z}^d}$ of real numbers such that $0 \leq \mu_i \leq C$ for all $i \in \mathbb{Z}^d$, the stochastic matrix $(A(i, j))_{i, j \in \mathbb{Z}^d}$ defined by (17), and a locally integrable function $\varphi : (0, \infty) \mapsto [0, \infty)$. The equation $\mathbf{m}_t = \boldsymbol{\mu} t + \int_0^t \varphi(t-s) A \mathbf{m}_s ds$ with unknown $\mathbf{m} = (m_t^i)_{t \geq 0, i \in \mathbb{Z}^d}$ has a unique solution such that for all $t \geq 0$, $\sum_{i \in \mathbb{Z}^d} 2^{-|i|} \sup_{[0, t]} m_s^i < \infty$. Furthermore, m^i is of class C^1 on $[0, \infty)$ for all $i \in \mathbb{Z}^d$, and it holds that $\mathbf{m}'_t = \boldsymbol{\mu} + \int_0^t \varphi(t-s) A \mathbf{m}'_s ds$ and $\mathbf{m}'_t = \left(I + \sum_{n \geq 1} A^n \int_0^t \varphi^{*n}(s) ds \right) \boldsymbol{\mu}$. Finally, $u_t = \sup_{i \in \mathbb{Z}^d} \sup_{[0, t]} (m_s^i)'$ is finite for all $t \geq 0$ and it holds that $u_t \leq C + \int_0^t \varphi(t-s) u_s ds$.*

Proof. We proceed in a few steps.

Step 1. We first note that for any vector $\mathbf{x} = (x_i)_{i \in \mathbb{Z}^d}$, it holds that $\sum_{i \in \mathbb{Z}^d} 2^{-|i|} |(A\mathbf{x})_i| \leq 2 \sum_{i \in \mathbb{Z}^d} 2^{-|i|} |x_i|$. This easily follows from the explicit form of A .

Step 2. We next check uniqueness. Consider two solutions \mathbf{m} and $\tilde{\mathbf{m}}$ satisfying the required condition and put $h_t = \sum_{i \in \mathbb{Z}^d} 2^{-|i|} \sup_{[0, t]} |m_t^i - \tilde{m}_t^i|$. We have $h_t \leq \sum_{i \in \mathbb{Z}^d} 2^{-|i|} \int_0^t \varphi(t-s) |(A(\mathbf{m}_s - \tilde{\mathbf{m}}_s))_i| ds$. Using Step 1, we deduce that $h_t \leq 2 \int_0^t \varphi(t-s) h_s ds$ and thus $h_t = 0$ by Lemma 23-(i).

Step 3. We define $\mathbf{m}'_t = \left(I + \sum_{n \geq 1} A^n \int_0^t \varphi^{*n}(s) ds \right) \boldsymbol{\mu}$. Using that A is stochastic and that $\boldsymbol{\mu}$ is bounded (by C), we easily deduce that for all $i \in \mathbb{Z}^d$, $(m_t^i)' \leq C(1 + \sum_{n \geq 1} \int_0^t \varphi^{*n}(s) ds)$. This function is locally bounded because φ is locally integrable: use that $\Upsilon_k(t) = \sum_{n=1}^k \int_0^t \varphi^{*n}(s) ds$ satisfies $\Upsilon_k(t) \leq \int_0^t \varphi(s) ds + \int_0^t \varphi(t-s) \Upsilon_k(s) ds$ and use Lemma 23-(iii), which provides a uniform (in k) bound. Consequently, $u_t = \sup_{i \in \mathbb{Z}^d} \sup_{[0, t]} (m_s^i)'$ is finite for all $t \geq 0$. Similar arguments show that $(m_t^i)'$ is continuous on $[0, \infty)$, because $|(m_{t+h}^i)' - (m_t^i)'| \leq C \sum_{n \geq 1} \int_t^{t+h} \varphi^{*n}(s) ds$.

Step 4. A straightforward consequence of the definition of \mathbf{m}'_t is that it solves $\mathbf{m}'_t = \boldsymbol{\mu} + \int_0^t \varphi(t-s) A \mathbf{m}'_s ds$. Using that A is stochastic and that $\boldsymbol{\mu}$ is bounded, we immediately deduce that

- [28] L. Massoulié, *Stability results for a general class of interacting point processes dynamics, and applications*, Stochastic Process. Appl. 75 (1998), 1–30.
- [29] Y. Ogata, *The asymptotic behaviour of maximum likelihood estimators for stationary point processes*, Ann. Instit. Math. Statist. 30 (1978), 243–261.
- [30] Y. Ogata, *On Lewis simulation method for point processes*, IEEE Transactions on Information Theory 27 (1981), 23–30.
- [31] Y. Ogata, *Seismicity analysis through point-process modeling: A review*, Pure and Applied Geophysics 155 (1999), 471–507.
- [32] M. Okatan, M.A. Wilson, E.N. Brown, *analyzing functional connectivity using a network likelihood model of ensemble neural spiking activity*, Neural Computation 17 (2005), 1927–1961.
- [33] J.W. Pillow, J. Shlens, L. Paninski, A. Scher, A.M. Litke, E.J. Chichilnisky, E.P. Simoncelli, *Spatio-temporal correlations and visual signalling in a complete neuronal population*, Nature 454 (2008), 995–999.
- [34] P. Reynaud-Bouret and S. Schbath, *Adaptive estimation for Hawkes processes: application to genome analysis*, Ann. Statist. 38 (2010), 2781–2822.
- [35] P. Reynaud-Bouret, V. Rivoirard, F. Grammont, C. Tuleau-Malot, *Goodness-of-fit tests and nonparametric adaptive estimation for spike train analysis*, Journal of Math. Neuroscience, to appear. hal-00789127
- [36] P. Reynaud-Bouret, V. Rivoirard, C. Tuleau-Malot, *Inference of functional connectivity in Neurosciences via Hawkes processes*, 1st IEEE Global Conference on Signal and Information Processing, 2013.
- [37] A.S. Sznitman, *Topics in propagation of chaos*, Ecole d’Été de Probabilités de Saint-Flour XIX-1989, Vol. 1464 of Lecture Notes in Math. Springer, 1991, 165–251.
- [38] A. Simma, M.I. Jordan, *Modeling events with cascades of Poisson processes*. <http://www.eecs.berkeley.edu/Pubs/TechRpts/2010/EECS-2010-109.html>
- [39] S.-H. Yang, H. Zha, *Mixture of mutually exciting processes for viral diffusion*, Proceedings of the 30th International Conf. on Machine Learning (ICML), 2013.
- [40] K. Zhou, H. Zha, L. Song, *Learning triggering kernels for multi-dimensional Hawkes processes*, Proceedings of the 30th International Conf. on Machine Learning (ICML), 2013.
- [41] L. Zhu, *Central limit theorem for nonlinear Hawkes processes*, J. App. Probab. 50 (2013), 760–771.
- [42] L. Zhu, *Large deviations for Markovian nonlinear Hawkes processes*, Ann. App. Probab., to appear. arXiv:1108.2432
- [43] L. Zhu, *Nonlinear Hawkes Processes*, PhD, New York University, 2013. arXiv:1304.7531

SYLVAIN DELATTRE, LABORATOIRE DE PROBABILITÉS ET MODÈLES ALÉATOIRES, UMR 7599, UNIVERSITÉ PARIS DIDEROT, CASE COURRIER 7012, AVENUE DE FRANCE, 75205 PARIS CEDEX 13, FRANCE.

E-mail address: sylvain.delattre@univ-paris-diderot.fr

NICOLAS FOURNIER, LABORATOIRE DE PROBABILITÉS ET MODÈLES ALÉATOIRES, UMR 7599, UNIVERSITÉ PIERRE-ET-MARIE CURIE, CASE 188, 4 PLACE JUSSIEU, F-75252 PARIS CEDEX 5, FRANCE.

E-mail address: nicolas.fournier@upmc.fr

MARC HOFFMANN, CEREMADE, CNRS-UMR 7534, UNIVERSITÉ PARIS DAUPHINE, PLACE DU MARÉCHAL DE LATTRE DE TASSIGNY 75775 PARIS CEDEX 16, FRANCE.

E-mail address: hoffmann@ceremade.dauphine.fr