

# A non-expanding transport distance for some structured equations

Nicolas Fournier\*      Benoît Perthame<sup>†‡</sup>

September 6, 2021

## Abstract

Structured equations are a standard modeling tool in mathematical biology. They are integro-differential equations where the unknown depends on one or several variables, representing the state or phenotype of individuals. A large literature has been devoted to many aspects of these equations and in particular to the study of measure solutions. Here we introduce a transport distance closely related to the Monge-Kantorovich distance, which appears to be non-expanding for several (mainly linear) examples of structured equations.

---

2010 *Mathematics Subject Classification*: 35A05; 35K55; 60J99; 28A33

*Keywords and phrases*: Transport distances; Monge-Kantorovich distance; Coupling; Structured equations; Mathematical biology.

## Introduction

The subject of structured equations arises in several areas of biology and extends ordinary differential equations by including parameters chosen because they bring some influence on the population dynamics, see the books [14, 30, 38]. This leads to various integro-differential equations and partial differential equation (P.D.E.) which also appear in many other areas as physics, communication science and industry. Besides the interesting modeling issues, the questions which have been considered are about existence of solutions, entropy properties and, mostly, long term convergence to steady states, with possibly exponential rate of convergence. Another question concerns measure solutions, possibly after renormalization [2, 15]. Furthermore, in the context of a nonlinear neuroscience problem, convergence of a particle system has recently been proved using transport costs with specific costs precisely adapted to the coefficients [22].

The present paper aims at showing that a simple variant of the Monge-Kantorovich transport distance appears to be non-expanding along several structured equations. These include the renewal equation and a few other models listed below.

We work in a state space that we denote by  $\mathcal{J}$ , which can be  $[0, \infty)$ ,  $[0, \infty) \times T$  with  $T$  a discrete torus,  $[0, \infty)^2$  or  $[0, \infty) \times \mathbb{R}^d$  and we always use a cost function  $\varrho : \mathcal{J} \times \mathcal{J} \mapsto [0, \infty)$  which satisfies

---

\*Sorbonne Université, CNRS, Laboratoire de Probabilité, Statistique et Modélisation, F-75005 Paris, France. Email: nicolas.fournier@sorbonne-universite.fr

<sup>†</sup>Sorbonne Université, CNRS, Université de Paris, Inria, Laboratoire Jacques-Louis Lions, F-75005 Paris, France. Email: benoit.perthame@sorbonne-universite.fr.

<sup>‡</sup>B.P. has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 740623).

$\varrho(x, x) = 0$  and  $\varrho(x, y) = \varrho(y, x) > 0$  for  $x \neq y$  and will typically take the form  $\varrho(y, x) = \min(|x - y|, a)$  for some  $a > 0$  chosen according to the equation at hand. We recall that for two probability measures  $u_1, u_2 \in \mathcal{P}(\mathcal{J})$ , the transport cost is defined as

$$\begin{cases} \mathcal{T}_\varrho(u_1, u_2) = \inf_{v \in \mathcal{H}(u_1, u_2)} \iint \varrho(x, y) v(dx, dy), \\ \mathcal{H}(u_1, u_2) = \{v \in \mathcal{P}(\mathcal{J} \times \mathcal{J}) \text{ with marginals } u_1 \text{ and } u_2\}. \end{cases} \quad (1)$$

When  $\varrho$  is a distance on  $\mathcal{J}$ ,  $\mathcal{T}_\varrho$  is a distance on  $\mathcal{P}(\mathcal{J})$  and, with a slight abuse of language, one refers to the Monge-Kantorovich distance. Recent accounts about the wide theory of optimal transport can be found in the books [41, 1, 39]. We use a specific distance which applies to a number of equations, but for each of our models, other distances and couplings (see below) may apply with their own advantages. For instance the case of the renewal equation was inspired from another one in [22]. Also for TCP connections (which is a particular case of the growth-fragmentation equation), the classical Wasserstein distance is used in [3, 11]. The motivation for the distance at hand, besides its simplicity, is that it applies to many models, with the same strategy for treating the inhomogeneity, and with the same coupling choice. This coupling choice consists in making jump both variables simultaneously as often as possible.

Our approach relies on the coupling method for the integro-differential equation under investigation, see the survey paper [23]. The first example of use of the coupling method, to our knowledge, can be traced back to Dobrushin [18], where the nonlinear Vlasov equation is derived as mean-field limit of a deterministic system of interacting particles, making use of some transport cost. No P.D.E. is written for the coupling in [18], because everything may be expressed in terms of characteristics. See [26, Section 3] for a P.D.E. analogue to Dobrushin's argument. In the same spirit, the Euler equation is derived from a deterministic system of interacting vortices in Marchioro-Pulvirenti [29, Section 5.3], using also a coupling argument, see also [27] for a result with the strong transport distance  $d_\infty$ .

The paper is organized as follows. We begin with the renewal equation in order to present in details the results and method. Building on this, we extend the method to a system of renewal equations, some space-age structured equation, the multi-time renewal equation, the growth-fragmentation equation, and to an age-size coupled model. All these equations are linear. We complete our study with a model with sexual reproduction, which is quadratic, and generates new difficulties. Our setting is very general and does not use uniqueness of solutions, therefore we complete them with a technical appendix devoted to a uniqueness result by the Hilbert duality method when further regularity on the coefficients is assumed. A second appendix mentions some hints toward the non-conservative models.

## 1 The renewal equation

Our first example, also the simplest, is the general renewal equation. It allows to introduce the method and to explain the choice of cost within the setting of the equation

$$\begin{cases} \frac{\partial u_t(x)}{\partial t} + \frac{\partial [g(x)u_t(x)]}{\partial x} + d(x)u_t(x) = b(x)N(t), & t \geq 0, x \geq 0, \\ u_t(x=0) = 0, & N(t) = \int_0^\infty d(x)u_t(dx), & t \geq 0. \end{cases} \quad (2)$$

When  $b = \delta_0$ , the Dirac mass at 0, and  $g \equiv 1$ , we find the classical renewal equation [20]. The more general version at hand is motivated by various models proposed in mathematical neuroscience,

[22, 34, 37]. We assume that

$$g, d \in C([0, \infty)), \quad b \in \mathcal{P}([0, \infty)), \quad g \text{ is non-increasing}, \quad g(0) \geq 0, \quad d \geq 0. \quad (3)$$

We will further suppose that

$$\exists a > 0 \quad \text{such that} \quad a \leq \inf_{|x-y| \leq a} \frac{|x-y| \max(d(x), d(y))}{|d(x) - d(y)|}. \quad (4)$$

Observe that this last condition holds true with  $a = \min(a_0, 1)$  as soon as  $a_0 > 0$ , where

$$a_0 = \inf_{|x-y| \leq 1} \frac{|x-y| \max(d(x), d(y))}{|d(x) - d(y)|}.$$

For example,  $d(x) = \alpha + \beta x^p$  satisfies such a condition, provided  $\alpha > 0$ ,  $\beta \geq 0$  and  $p \geq 1$ , as well as any Lipschitz and uniformly positive function  $d$ .

**Theorem 1** *Assume (3)-(4). We consider the cost function on  $(0, \infty) \times (0, \infty)$  defined by*

$$\varrho(x, y) = \min(a, |x - y|).$$

For any  $u_0^1, u_0^2 \in \mathcal{P}([0, \infty))$ , there exists a pair of weak measure solutions  $(u_t^1)_{t \geq 0}, (u_t^2)_{t \geq 0} \subset \mathcal{P}([0, \infty))$  to (2) starting from  $u_0^1$  and  $u_0^2$ , i.e., such that for  $i = 1, 2$  and all  $t \geq 0$ ,

$$\int_0^t \int_0^\infty d(x) u_s^i(dx) ds < \infty \quad (5)$$

and for all  $t \geq 0$ , all  $\varphi \in C_c^1([0, \infty))$ ,

$$\int_0^\infty \varphi(x) u_t^i(dx) = \int_0^\infty \varphi(x) u_0^i(dx) + \int_0^t \int_0^\infty \left[ g(x) \varphi'(x) + d(x) \int_0^\infty (\varphi(z) - \varphi(x)) b(dz) \right] u_s^i(dx) ds. \quad (6)$$

Moreover, for all  $t \geq 0$ , we have

$$\mathcal{T}_\varrho(u_t^1, u_t^2) \leq \mathcal{T}_\varrho(u_0^1, u_0^2).$$

Notice that, even if we did not mention it, it follows from (6) that  $u \in C_w([0, \infty); \mathcal{P}([0, \infty)))$ . Also, the generality of this statement relies on the price that the solutions  $u_t^1, u_t^2$  may depend on the choice of the pair  $(u_0^1, u_0^2)$ . However, when  $g \in C_b^1$ , the solutions are unique in distributional sense as proved in the appendix, and then the result is more standard. The regularity of  $g$  can certainly be lowered in view of the theory developed in [6, 7, 17].

Throughout the paper, in the proofs, we use the notations  $a \vee b := \max(a, b)$  and  $a \wedge b := \min(a, b)$ .

**Proof.** We assume (3), fix  $u_0^1, u_0^2 \in \mathcal{P}([0, \infty))$  and consider any  $v_0 \in \mathcal{H}(u_0^1, u_0^2)$ . There exists a family  $(v_t)_{t \geq 0}$  of probability measures on  $[0, \infty)^2$ , starting from  $v_0$ , such that for all  $t \geq 0$ ,

$$\int_0^t \iint [d(x) + d(y)] v_s(dx, dy) ds < \infty, \quad (7)$$

and which weakly solves

$$\begin{aligned} \frac{\partial v_t}{\partial t} + \frac{\partial[g(x)v_t]}{\partial x} + \frac{\partial[g(y)v_t]}{\partial y} + [d(x) \vee d(y)]v_t &= b(x)\delta(x-y) \iint [d(x') \wedge d(y')] v_t(dx', dy') \\ &+ b(x) \int (d(x') - d(y))_+ v_t(dx', y) + b(y) \int (d(y') - d(x))_+ v_t(x, dy'). \end{aligned}$$

This means that for all  $t \geq 0$ , all  $\varphi \in C_c^1([0, \infty)^2)$ ,

$$\begin{aligned} \iint \varphi(x, y)v_t(dx, dy) &= \iint \varphi(x, y)v_0(dx, dy) + \int_0^t \iint \left[ g(x) \frac{\partial \varphi(x, y)}{\partial x} + g(y) \frac{\partial \varphi(x, y)}{\partial y} \right] v_s(dx, dy) ds \\ &+ \int_0^t \iiint [\varphi(z, z) - \varphi(x, y)] [d(x) \wedge d(y)] b(dz) v_s(dx, dy) ds \\ &+ \int_0^t \iiint [\varphi(z, y) - \varphi(x, y)] (d(x) - d(y))_+ b(dz) v_s(dx, dy) ds \\ &+ \int_0^t \iiint [\varphi(x, z) - \varphi(x, y)] (d(y) - d(x))_+ b(dz) v_s(dx, dy) ds. \end{aligned} \quad (8)$$

Intuitively, this coupling equation drifts independently the two variables  $x, y$  according to  $g$  between jumps, makes jump simultaneously the two variables  $x, y$  at rate  $d(x) \wedge d(y)$  to the same value  $z$  chosen according to the distribution  $b$ , makes jump only  $x$  at rate  $d(x) - d(y)$  when  $d(x) > d(y)$ , and makes jump only  $y$  at rate  $d(y) - d(x)$  when  $d(y) > d(x)$ .

Using (7) and then (8) with a function  $\varphi$  depending only on  $x$ , observing that  $v_0 \in \mathcal{H}(u_0^1, u_0^2)$  and that

$$d(x) \wedge d(y) + (d(x) - d(y))_+ = d(x),$$

we deduce that the first marginal  $u_t^1(dx) = \int_{y \in [0, \infty)} v_t(dx, dy)$  satisfies (5)-(6). Similarly, the second marginal  $u_t^2(dy) = \int_{x \in [0, \infty)} v_t(dx, dy)$  satisfies (5)-(6). And it holds that  $v_t \in \mathcal{H}(u_t^1, u_t^2)$  for all  $t \geq 0$ .

The existence for (7)-(8) follows from classical arguments, using e.g., an approximate problem where  $g, d$  are replaced by smooth and bounded functions, and from the following *a priori* tightness estimate. By the de la Vallée Poussin theorem (see [28], Theorem 8, p. 208), there exists a function  $h : [0, \infty) \rightarrow [0, \infty)$  such that  $\lim_{x \rightarrow \infty} h(x) = \infty$  and such that

$$C := \iint [h(x) + h(y)] v_0(dx, dy) + \int h(z) b(dz) < \infty.$$

One can moreover choose  $h$  smooth and satisfying  $0 \leq h' \leq 1$ . Applying (8) with  $\varphi(x, y) = h(x) + h(y)$ , one immediately concludes that for all  $t \geq 0$ ,

$$\begin{aligned} \iint [h(x) + h(y)] v_t(dx, dy) &= \iint [h(x) + h(y)] v_0(dx, dy) + \int_0^t \iint [g(x)h'(x) + g(y)h'(y)] v_s(dx, dy) ds \\ &+ \int_0^t \iint \left[ d(x) \left( \int h(z) b(dz) - h(x) \right) + d(y) \left( \int h(z) b(dz) - h(y) \right) \right] v_s(dx, dy) ds \\ &\leq C + 2\bar{C}t + \int_0^t \iint \left[ d(x)(C - h(x)) + d(y)(C - h(y)) \right] v_s(dx, dy) ds, \end{aligned}$$

where  $\bar{C} = \sup_{x \geq 0} g(x)h'(x)$  is finite because  $g$  is continuous and non-increasing and because  $h'$  is  $[0, 1]$ -valued. Since now  $d$  is continuous, non-negative and since  $h$  increases to infinity, there is  $L > 0$  such that  $d(x)(C - h(x)) \leq L - d(x)h(x)/2$ , whence finally

$$\iint [h(x) + h(y)]v_t(dx, dy) + \frac{1}{2} \int_0^t \iint [d(x)h(x) + d(y)h(y)]v_s(dx, dy)ds \leq C + 2\bar{C}t + 2Lt. \quad (9)$$

This last *a priori* tightness estimate is sufficient to prove existence for (7)-(8). Indeed, consider, for each  $\varepsilon \in (0, 1]$ , the solution  $(v_t^\varepsilon)_{t \geq 0}$  to an approximate problem, e.g. (7)-(8) with  $g, d$  are replaced by smooth and bounded functions, starting from  $v_0$ . Then (9) holds for  $v^\varepsilon$ , uniformly in  $\varepsilon \in (0, 1]$ . Since  $\lim_{x \rightarrow \infty} h(x) = \infty$ , this is sufficient to show that the family  $((v_t^\varepsilon)_{t \geq 0}, \varepsilon \in (0, 1])$  is relatively compact in  $C([0, \infty), \mathcal{P}([0, \infty)^2))$ , the set  $\mathcal{P}([0, \infty)^2)$  being endowed with the topology of weak convergence (i.e. when taking continuous and bounded test functions). Hence one can find  $\varepsilon_k \rightarrow 0$  such that  $(v_t^{\varepsilon_k})_{t \geq 0}$  converges to some  $(v_t)_{t \geq 0}$  in  $C([0, \infty), \mathcal{P}([0, \infty)^2))$ . It is then routine, using again that (9) holds for  $v^\varepsilon$ , uniformly in  $\varepsilon \in (0, 1]$ , and that  $\lim_{x \rightarrow \infty} h(x) = \infty$ , to show that  $(v_t)_{t \geq 0}$  satisfies (7)-(8).

We next fix  $a > 0$ , set  $\rho(x, y) = |x - y| \wedge a$  and we choose a coupling  $v_0 \in \mathcal{H}(u_0^1, u_0^2)$  such that  $\iint \rho(x, y)v_0(dx, dy) = \mathcal{T}_\rho(u_0^1, u_0^2)$ . We apply (8) with  $\varphi = \rho$  (or more precisely, firstly to some smooth and compactly supported approximation  $\rho_\varepsilon$  of  $\rho$  and then let  $\varepsilon \rightarrow 0$ ). We notice that

$$g(x) \frac{\partial \rho(x, y)}{\partial x} + g(y) \frac{\partial \rho(x, y)}{\partial y} = \mathbb{1}_{\{|x-y| \leq a\}} \operatorname{sgn}(x-y)[g(x) - g(y)] \leq 0$$

because  $g$  is non-increasing. Since  $b$  is a probability measure, we obtain

$$\begin{aligned} \iint \rho(x, y)v_t(dx, dy) &\leq \mathcal{T}_\rho(u_0^1, u_0^2) - \int_0^t \iint \rho(x, y)[d(x) \vee d(y)]v_s(dx, dy)ds \\ &\quad + \int_0^t \iiint [\rho(z, y)(d(x) - d(y))_+ + \rho(x, z)(d(y) - d(x))_+] b(dz)v_s(dx, dy)ds. \end{aligned}$$

Recalling that  $\mathcal{T}_\rho(u_t^1, u_t^2) \leq \iint \rho(x, y)v_t(dx, dy)$ , in order to complete the proof, it is therefore sufficient to verify the inequality, for all  $x, y \geq 0$ ,

$$I(x, y) := \frac{(d(x) - d(y))_+}{d(x) \vee d(y)} \int \varrho(z, y)b(dz) + \frac{(d(y) - d(x))_+}{d(x) \vee d(y)} \int \varrho(x, z)b(dz) \leq \varrho(x, y). \quad (10)$$

Since  $\varrho(x, y) = a \wedge |x - y|$  and since  $b$  is a probability measure, we have

$$I(x, y) \leq \frac{(d(x) - d(y))_+}{d(x) \vee d(y)} a + \frac{(d(y) - d(x))_+}{d(x) \vee d(y)} a = \frac{|d(x) - d(y)|}{d(x) \vee d(y)} a \leq |x - y| \wedge a.$$

Indeed, the last inequality is obvious if  $|x - y| \geq a$  and follows from (4) otherwise.  $\square$

**Remark 2 (Strict contraction)** *Strict contraction*,  $\mathcal{T}_\varrho(u_t^1, u_t^2) \leq e^{-\theta t} \mathcal{T}_\varrho(u_0^1, u_0^2)$ , can also be obtained under the assumption that there are  $a > 0$ ,  $\theta \in (0, 1)$  such that for all  $x, y \geq 0$ ,

$$\frac{|d(x) - d(y)|}{\max(d(x), d(y))} a \leq (1 - \theta) \min(|x - y|, a).$$

For example, this condition holds true when  $d$  is nondecreasing,  $K$ -Lipschitz-continuous and  $[c, C]$ -valued for some  $C > c > 0$ , with  $\theta = c/C$  and  $a = c(C - c)/(KC)$ .

## 2 A system of renewal equations

In mathematical biology, it is usual to describe the cell cycle, see [30, 38], using a system of renewal equations coupled at their boundaries as follows

$$\begin{cases} \frac{\partial u_t(x,i)}{\partial t} + \frac{\partial [g_i(x)u_t(x,i)]}{\partial x} + d_i(x)u_t(x,i) = 0, & t \geq 0, x \geq 0, i = 1, \dots, I, \\ u_t(x=0, i) = \int_0^\infty d_{i-1}(x)u_t(dx, i-1), & t \geq 0, i = 1, \dots, I, \end{cases} \quad (11)$$

with the convention  $u_t(x, 0) = u_t(x, I)$ . In terms of stochastic processes, a cell with age  $x$  in state  $i$  ages chronologically (when, say,  $g_i \equiv 1$ ) until, with rate  $d_i(x)$ , it changes state to  $i+1$  where it starts with age  $x=0$ . So the state space is

$\mathcal{J} = [0, \infty) \times T$ , with  $T = \{1, \dots, I\}$  the torus, i.e., states 0 and  $I+1$  are identified to states  $I$  and 1.

We assume that for all  $i \in T$ ,

$$g_i, d_i \in C([0, \infty)), \quad g_i \text{ is non-increasing}, \quad g_i(0) \geq 0, \quad d_i \geq 0, \quad (12)$$

and that

$$\exists a > 0 \quad \text{such that} \quad a \leq \min_{1 \leq i \leq I} \inf_{|x-y| \leq a} \frac{|x-y| \max(d_i(x), d_i(y))}{|d_i(x) - d_i(y)|}. \quad (13)$$

This condition is satisfied if for all  $i \in T$ , there are  $\alpha_i > 0$ ,  $\beta_i \geq 0$  and  $p_i \geq 1$  such that  $d_i(x) = \alpha_i + \beta_i x^{p_i}$ , or if all  $d_i$ 's are Lipschitz and uniformly positive.

**Theorem 3** *Assume (12)-(13). We consider the cost on  $\mathcal{J} \times \mathcal{J}$  defined by*

$$\varrho(x, i, y, j) = \min(a, |x-y|) \mathbf{1}_{\{i=j\}} + a \mathbf{1}_{\{i \neq j\}}.$$

For any  $u_0^1, u_0^2 \in \mathcal{P}(\mathcal{J})$ , there exists a pair of weak measure solutions  $(u_t^1)_{t \geq 0}, (u_t^2)_{t \geq 0} \subset \mathcal{P}(\mathcal{J})$  to (11) starting from  $u_0^1$  and  $u_0^2$ , i.e., such that for  $j = 1, 2$  and all  $t \geq 0$

$$\int_0^t \int_{\mathcal{J}} d_i(x) u_s^j(dx, di) ds < \infty \quad (14)$$

and for  $j = 1, 2$ , all  $t \geq 0$  and all  $\varphi \in C_c^1(\mathcal{J})$ ,

$$\begin{aligned} \int_{\mathcal{J}} \varphi(x, i) u_t^j(dx, di) &= \int_{\mathcal{J}} \varphi(x, i) u_0^j(dx, di) \\ &+ \int_0^t \int_{\mathcal{J}} \left[ g_i(x) \varphi'(x, i) + d_i(x) [\varphi(0, i+1) - \varphi(x, i)] \right] u_s^j(dx, di) ds. \end{aligned} \quad (15)$$

Moreover, for all  $t \geq 0$ , we have

$$\mathcal{T}_\varrho(u_t^1, u_t^2) \leq \mathcal{T}_\varrho(u_0^1, u_0^2).$$

Note that if  $u_t(x, i)$  is a strong solution to (11), then  $u_t(dx, di) := \sum_{k \in T} u_t(x, k) dx \delta_k(di)$  is a weak measure solution to (11). Note also that the comments after Theorem 1 apply here too.

**Proof.** We assume (12)-(13) and consider any coupling  $v_0 \in \mathcal{H}(u_0^1, u_0^2)$ . There exists a family  $(v_t)_{t \geq 0}$  of probability measures on  $\mathcal{J}^2$  such that for all  $t \geq 0$ ,

$$\int_0^t \iint [d_i(x) + d_j(y)] v_s(dx, di, dy, dj) ds < \infty, \quad (16)$$

and which weakly solves

$$\begin{aligned} \frac{\partial v_t}{\partial t} + \frac{\partial [g_i(x)v_t]}{\partial x} + \frac{\partial [g_j(y)v_t]}{\partial y} + [(d_i(x) \vee d_j(y)) \mathbf{1}_{\{i=j\}} + (d_i(x) + d_j(y)) \mathbf{1}_{\{i \neq j\}}] v_t = \\ \delta_{(0,0)}(x, y) \mathbf{1}_{\{i=j\}} \iint (d_{i-1}(x') \wedge d_{i-1}(y')) v_t(dx', i-1, dy', j-1) \\ + \delta_0(x) \mathbf{1}_{\{j=i-1\}} \int (d_{i-1}(x') - d_j(y))_+ v_t(dx', i-1, y, j) \\ + \delta_0(y) \mathbf{1}_{\{i=j-1\}} \int (d_{j-1}(y') - d_i(x'))_+ v_t(x, i, dy', j-1) \\ + \delta_0(x) \mathbf{1}_{\{j \neq i-1\}} \int d_{i-1}(x') v_t(dx', i-1, y, j) + \delta_0(y) \mathbf{1}_{\{i \neq j-1\}} \int d_{j-1}(y') v_t(x, i, dy', j-1). \end{aligned}$$

This means that for all  $\varphi \in C_c^1(\mathcal{J}^2)$ , all  $t \geq 0$ ,

$$\begin{aligned} \iint \varphi(x, i, y, j) v_t(dx, di, dy, dj) = \iint \varphi(x, i, y, j) v_0(dx, di, dy, dj) \\ + \int_0^t \iint \left[ g_i(x) \frac{\partial \varphi(x, i, y, j)}{\partial x} + g_j(y) \frac{\partial \varphi(x, i, y, j)}{\partial y} \right] v_s(dx, di, dy, dj) ds \\ + \int_0^t \iint \left( [\varphi(0, i+1, 0, j+1) - \varphi(x, i, y, j)] \mathbf{1}_{\{i=j\}} (d_i(x) \wedge d_j(y)) \right. \\ \quad + [\varphi(0, i+1, y, j) - \varphi(x, i, y, j)] \mathbf{1}_{\{i=j\}} (d_i(x) - d_j(y))_+ \\ \quad + [\varphi(x, i, 0, j+1) - \varphi(x, i, y, j)] \mathbf{1}_{\{i=j\}} (d_j(y) - d_i(x))_+ \\ \quad + [\varphi(0, i+1, y, j) - \varphi(x, i, y, j)] \mathbf{1}_{\{i \neq j\}} d_i(x) \\ \quad \left. + [\varphi(x, i, 0, j+1) - \varphi(x, i, y, j)] \mathbf{1}_{\{i \neq j\}} d_j(y) \right) v_s(dx, di, dy, dj) ds. \quad (17) \end{aligned}$$

Using (16) and (17) with a function  $\varphi$  depending only on  $(x, i)$  (or  $(y, j)$ ), we see that the first marginal  $u_t^1(dx, di) = \int_{(y,j) \in \mathcal{J}} v_t(dx, di, dy, dj)$  (or the second marginal  $u_t^2(dy, dj) = \int_{(x,i) \in \mathcal{J}} v_t(dx, di, dy, dj)$ ) satisfies (14) and (15).

As before, the existence for (16)-(17) follows from classical arguments, using an approximate problem where  $g_i, d_i$  are replaced by smooth and bounded functions, and from the following *a priori* tightness estimate. By the de la Vallée Poussin theorem, there exists a function  $h : [0, \infty) \rightarrow [0, \infty)$  such that  $\lim_{x \rightarrow \infty} h(x) = \infty$  and such that

$$C := \iint [h(x) + h(y)] v_0(dx, di, dy, dj) < \infty.$$

One can moreover choose  $h$  smooth, satisfying  $h(0) = 0$  and  $0 \leq h' \leq 1$ . Applying (17) with

$\varphi(x, i, y, j) = h(x) + h(y)$ , one immediately concludes that for all  $t \geq 0$ ,

$$\begin{aligned} \iint [h(x) + h(y)]v_t(dx, di, dy, dj) &= \iint [h(x) + h(y)]v_0(dx, di, dy, dj) \\ &\quad + \int_0^t \iint [g_i(x)h'(x) + g_j(y)h'(y)]v_s(dx, di, dy, dj)ds \\ &\quad - \int_0^t \iint [d_i(x)h(x) + d_j(y)h(y)]v_s(dx, di, dy, dj)ds. \end{aligned}$$

Setting  $\bar{C} = \sup_{i \in T, x \geq 0} g_i(x)h'(x)$ , which is finite because  $g_i$  is continuous, non-increasing and because  $h'$  is  $[0, 1]$ -valued, we end up with

$$\iint [h(x) + h(y)]v_t(dx, di, dy, dj) + \int_0^t \iint [d_i(x)h(x) + d_j(y)h(y)]v_s(dx, di, dy, dj)ds \leq C + 2\bar{C}t.$$

This last *a priori* tightness estimate is sufficient to prove existence for (16)-(17).

We now apply (17) with  $\varphi = \varrho$ , after regularization as before, where we recall that  $\varrho(x, i, y, j) = (a \wedge |x - y|)\mathbb{1}_{\{i=j\}} + a\mathbb{1}_{\{i \neq j\}}$ . Since  $g_i$  is non-increasing, we have

$$g_i(x) \frac{\partial \varrho(x, i, y, j)}{\partial x} + g_j(y) \frac{\partial \varrho(x, i, y, j)}{\partial y} = \mathbb{1}_{\{i=j\}} \mathbb{1}_{\{|x-y| \leq a\}} \text{sgn}(x-y)[g_i(x) - g_j(y)] \leq 0.$$

Hence we find, choosing  $v_0$  such that  $\iint \varrho(x, i, y, j)v_0(dx, di, dy, dj) = \mathcal{T}_\rho(u_0^1, u_0^2)$ ,

$$\begin{aligned} &\iint \varrho(x, i, y, j)v_t(dx, di, dy, dj) \\ &\leq \mathcal{T}_\rho(u_0^1, u_0^2) + \int_0^t \iint \left( [\varrho(0, i+1, 0, j+1) - \varrho(x, i, y, j)] \mathbb{1}_{\{i=j\}} (d_i(x) \wedge d_j(y)) \right. \\ &\quad + [\varrho(0, i+1, y, j) - \varrho(x, i, y, j)] \mathbb{1}_{\{i=j\}} (d_i(x) - d_j(y))_+ \\ &\quad + [\varrho(x, i, 0, j+1) - \varrho(x, i, y, j)] \mathbb{1}_{\{i=j\}} (d_j(y) - d_i(x))_+ \\ &\quad + [\varrho(0, i+1, y, j) - \varrho(x, i, y, j)] \mathbb{1}_{\{i \neq j\}} d_i(x) \\ &\quad \left. + [\varrho(x, i, 0, j+1) - \varrho(x, i, y, j)] \mathbb{1}_{\{i \neq j\}} d_j(y) \right) v_s(dx, di, dy, dj)ds. \end{aligned}$$

With our choice of  $\varrho$ , the two last lines are non-positive. We thus arrive at

$$\iint \varrho(x, i, y, j)v_t(dx, di, dy, dj) \leq \mathcal{T}_\rho(u_0^1, u_0^2) - \int_0^t \iint \mathbb{1}_{\{i=j\}} \Delta_i(x, y)v_s(dx, di, dy, dj)ds,$$

where

$$\begin{aligned} \Delta_i(x, y) &= (d_i(x) \vee d_i(y))(|x - y| \wedge a) - (d_i(x) - d_i(y))_+ a - (d_i(y) - d_i(x))_+ a \\ &= (d_i(x) \vee d_i(y))(|x - y| \wedge a) - |d_i(x) - d_i(y)|a. \end{aligned}$$

Next, we check that  $\Delta_i$  is always non-negative. If  $|x - y| \geq a$ , this is obvious. If  $|x - y| \leq a$ , this follows from (13). Since  $\mathcal{T}_\rho(u_t^1, u_t^2) \leq \iint \rho(x, i, y, j)v_t(dx, di, dy, dj)$ , the proof is complete.  $\square$



### 3 Space and age structure

Next, we consider an example similar to that in the previous section, when the discrete parameter  $i$  is replaced by a continuous parameter  $z \in \mathbb{R}^d$ , which represents space or a physiological trait. The formalism makes the link with the heat equation through a standard physical process used in particular to describe diffusion or anomalous diffusion, see recent analyses in [36, 9, 4]. We depart from the equation

$$\begin{cases} \varepsilon^2 \frac{\partial u_t(x, z)}{\partial t} + \frac{\partial u_t(x, z)}{\partial x} + d(x)u_t(x, z) = 0, & t \geq 0, x > 0, z \in \mathbb{R}^d, \\ u_t(x = 0, z) = \int_0^\infty \int_{\mathbb{R}^d} d(x)u_t(dx, z + \varepsilon\eta)k(d\eta), & t \geq 0, z \in \mathbb{R}^d. \end{cases} \quad (18)$$

This equation models particles characterized by their age  $x$  and position  $z$ . When in state  $x, z$ , the particle's age  $x$  grows linearly until there is a jump, at rate  $d(x)$ , resulting in the particle moving from  $z$  to  $z - \varepsilon\eta$  (with  $\eta$  chosen according to the probability density  $k$ ). At each jump, the age is reset to 0. Hence the state space is here  $\mathcal{J} = [0, \infty) \times \mathbb{R}^d$ . Under a few assumptions, it is known that, for  $u_\varepsilon$  the solution to (18),  $U_\varepsilon(t, z) = \int_0^\infty u_\varepsilon(t, x, z)dx$  converges, as  $\varepsilon \rightarrow 0$ , to the solution of the heat equation in  $\mathbb{R}^d$ .

We assume that

$$d \in C([0, \infty)), \quad d \geq 0, \quad k \in \mathcal{P}(\mathbb{R}^d), \quad (19)$$

and, again, that

$$\exists a > 0 \quad \text{such that} \quad a \leq \inf_{|x-y| \leq a} \frac{|x-y| \max(d(x), d(y))}{|d(x) - d(y)|}. \quad (20)$$

**Theorem 4** *Assume (19)-(20) and fix  $\varepsilon > 0$ . We consider the cost on  $\mathcal{J} \times \mathcal{J}$  defined by*

$$\varrho(x, z, y, r) = \min(a, |x - y| + |z - r|).$$

*For any  $u_0^1, u_0^2 \in \mathcal{P}(\mathcal{J})$ , there exists a pair of weak measure solutions  $(u_t^1)_{t \geq 0}, (u_t^2)_{t \geq 0} \subset \mathcal{P}(\mathcal{J})$  to (18) starting from  $u_0^1$  and  $u_0^2$ , i.e., such that for  $i = 1, 2$  and all  $t \geq 0$ ,*

$$\int_0^t \int_{\mathcal{J}} d(x)u_s^i(dx, dz)ds < \infty \quad (21)$$

*and for  $i = 1, 2$ , all  $t \geq 0$  and all  $\varphi \in C_c^1(\mathcal{J})$ ,*

$$\begin{aligned} \int_{\mathcal{J}} \varphi(x, z)u_t^i(dx, dz) &= \int_{\mathcal{J}} \varphi(x, z)u_0^i(dx, dz) \\ &+ \varepsilon^{-2} \int_0^t \int_{\mathcal{J}} \left[ \frac{\partial \varphi(x, z)}{\partial x} + d(x) \int_{\mathbb{R}^d} [\varphi(0, z - \varepsilon\eta) - \varphi(x, z)]k(d\eta) \right] u_s^i(dx, dz)ds. \end{aligned} \quad (22)$$

*Moreover, for all  $t \geq 0$ , we have*

$$\mathcal{T}_\varrho(u_t^1, u_t^2) \leq \mathcal{T}_\varrho(u_0^1, u_0^2).$$

**Proof.** We assume (19)-(20), consider  $u_0^1, u_0^2 \in \mathcal{P}(\mathcal{J})$  and a coupling  $v_0 \in \mathcal{H}(u_0^1, u_0^2)$ . There exists a family  $(v_t)_{t \geq 0}$  of probability measures on  $\mathcal{J}^2$  such that for all  $t \geq 0$

$$\int_0^t \iint [d(x) + d(y)] v_t(dx, dz, dy, dr) < \infty, \quad (23)$$

and which weakly solves

$$\begin{aligned} \varepsilon^2 \frac{\partial v_t}{\partial t} + \frac{\partial v_t}{\partial x} + \frac{\partial v_t}{\partial y} + (b(x) \vee b(y)) v_t = \\ \delta(x) \delta(y) \int_0^\infty \int_0^\infty \int_{\mathbb{R}^d} (b(x') \wedge b(y')) v_t(dx', z + \varepsilon \eta, dy', r + \varepsilon \eta) k(\eta) d\eta \\ + \delta(x) \int_{\mathcal{J}} (b(x') - b(y))_+ v_t(dx', z + \varepsilon \eta, y, r + \varepsilon \eta) k(\eta) d\eta \\ + \delta(y) \int_{\mathcal{J}} (b(y') - b(x))_+ v_t(x, z + \varepsilon \eta, dy', r + \varepsilon \eta) k(\eta) d\eta. \end{aligned}$$

This means that for all  $\varphi \in C_c^1(\mathcal{J}^2)$ , all  $t \geq 0$ ,

$$\begin{aligned} \iint \varphi(x, z, y, r) v_t(dx, dz, dy, dr) = \iint \varphi(x, z, y, r) v_0(dx, dz, dy, dr) \\ + \varepsilon^{-2} \int_0^t \iint \left[ \frac{\partial \varphi(x, z, y, r)}{\partial x} + \frac{\partial \varphi(x, z, y, r)}{\partial y} \right] v_s(dx, dz, dy, dr) ds \\ + \varepsilon^{-2} \int_0^t \iint \left( [(d(x) \wedge d(y)) \int_{\mathbb{R}^d} [\varphi(0, z - \varepsilon \eta, 0, r - \varepsilon \eta) - \varphi(x, z, y, r)] k(d\eta) \right. \\ \left. + (d(x) - d(y))_+ \int_{\mathbb{R}^d} [\varphi(0, z - \varepsilon \eta, y, r) - \varphi(x, z, y, r)] k(d\eta) \right. \\ \left. + (d(y) - d(x))_+ \int_{\mathbb{R}^d} [\varphi(x, z, 0, r - \varepsilon \eta) - \varphi(x, z, y, r)] k(d\eta) \right) v_s(dx, dz, dy, dr) ds. \quad (24) \end{aligned}$$

Using (23) and (24) with a function  $\varphi$  depending only on  $(x, z)$  (or  $(y, r)$ ), we see that the first marginal  $u_t^1(dx, dz) = \int_{(y,r) \in \mathcal{J}} v_t(dx, dz, dy, dr)$  (and the second one  $u_t^2(dy, dr) = \int_{(x,z) \in \mathcal{J}} v_t(dx, dz, dy, dr)$ ) satisfies (21) and (22).

The existence for (16)-(17) follows as usual from the following *a priori* tightness estimate. By the de la Vallée Poussin theorem, there is a function  $h : [0, \infty) \rightarrow [0, \infty)$  such that  $\lim_{x \rightarrow \infty} h(x) = \infty$  and

$$C := \iint [h(x) + h(y) + h(|z|) + h(|r|)] v_0(dx, dz, dy, dr) + \int h(\varepsilon|\eta|) k(d\eta) < \infty.$$

One can moreover choose  $h$  smooth, satisfying  $h(0) = 0$  and  $0 \leq h' \leq 1$ . Applying (24) with  $\varphi(x, z, y, r) = h(x) + h(y) + h(|z|) + h(|r|)$ , one immediately concludes that for all  $t \geq 0$ ,

$$\begin{aligned} \iint [h(x) + h(y) + h(|z|) + h(|r|)] v_t(dx, dz, dy, dr) \leq C + \varepsilon^{-2} \int_0^t \iint [h'(x) + h'(y)] v_s(dx, dz, dy, dr) ds \\ + \varepsilon^{-2} \int_0^t \iint \left[ d(x) \int [h(0) + h(|z - \varepsilon \eta|) - h(x) - h(|z|)] k(d\eta) \right. \\ \left. + d(y) \int [h(0) + h(|r - \varepsilon \eta|) - h(y) - h(|r|)] k(d\eta) \right] v_s(dx, dz, dy, dr) ds. \end{aligned}$$

Using that  $h'$  is  $[0, 1]$ -valued and that  $h(0) = 0$ , we observe that

$$\int [h(0) + h(|z - \varepsilon\eta|) - h(x) - h(|z|)]k(d\eta) \leq \int h(|\varepsilon\eta|)(d\eta) - h(x) \leq C - h(x).$$

Using now that  $d(x)[C - h(x)] \leq L - d(x)h(x)/2$  for some constant  $L > 0$ , we end up with

$$\begin{aligned} \iint [h(x) + h(y) + h(|z|) + h(|r|)]v_t(dx, dz, dy, dr) + \frac{1}{2\varepsilon^2} \int_0^t \iint [d(x)h(x) + d(y)h(y)]v_s(dx, dz, dy, dr)ds \\ \leq C + 2\varepsilon^{-2}t + 2\varepsilon^{-2}Lt. \end{aligned}$$

This *a priori* tightness estimate is sufficient, as usual, to prove existence for (23)-(24).

We now apply (24) with  $\varphi = \varrho$ , where we recall that  $\varrho(x, z, y, r) = a \wedge (|x - y| + |z - r|)$ . Since

$$\frac{\partial \varrho(x, z, y, r)}{\partial x} + \frac{\partial \varrho(x, z, y, r)}{\partial y} = 0,$$

we find, choosing  $v_0$  such that  $\iint \varrho(x, z, y, r)v_0(dx, dz, dy, dr) = \mathcal{T}_\rho(u_0^1, u_0^2)$ ,

$$\iint \varrho(x, z, y, r)v_t(dx, dz, dy, dr) \leq \mathcal{T}_\rho(u_0^1, u_0^2) + \varepsilon^{-2} \int_0^t \iint \Delta(x, z, y, r)v_s(dx, dz, dy, dr)ds,$$

where

$$\begin{aligned} \Delta(x, z, y, r) = & (d(x) \wedge d(y)) \int_{\mathbb{R}^d} [\varrho(0, z - \varepsilon\eta, 0, r - \varepsilon\eta) - \varrho(x, z, y, r)]k(d\eta) \\ & + (d(x) - d(y))_+ \int_{\mathbb{R}^d} [\varrho(0, z - \varepsilon\eta, y, r) - \varrho(x, z, y, r)]k(d\eta) \\ & + (d(y) - d(x))_+ \int_{\mathbb{R}^d} [\varrho(x, z, 0, r - \varepsilon\eta) - \varrho(x, z, y, r)]k(d\eta). \end{aligned}$$

Since  $\mathcal{T}_\rho(u_t^1, u_t^2) \leq \iint \rho(x, z, y, r)v_t(dx, dz, dy, dr)$ , it thus only remains to check that  $\Delta$  is always non-positive. But we have, assuming e.g. that  $d(x) \geq d(y)$ ,

$$\begin{aligned} \Delta(x, z, y, r) = & -d(x)[(|x - y| + |z - r|) \wedge a] + d(y)(|z - r| \wedge a) \\ & + (d(x) - d(y)) \int_{\mathbb{R}^d} [(|y| + |z - \varepsilon\eta - r|) \wedge a]k(d\eta) \\ \leq & -d(x)[(|x - y| + |z - r|) \wedge a] + d(y)(|z - r| \wedge a) + (d(x) - d(y))a. \end{aligned}$$

If  $|x - y| + |z - r| \geq a$ , we have

$$\Delta(x, z, y, r) \leq -d(x)a + d(y)a + (d(x) - d(y))a = 0.$$

If  $|x - y| + |z - r| \leq a$ , still assuming that  $d(x) \geq d(y)$ , we have

$$\begin{aligned} \Delta(x, z, y, r) \leq & -d(x)(|x - y| + |z - r|) + d(y)|z - r| + (d(x) - d(y))a \\ \leq & -d(x)|x - y| + (d(x) - d(y))a \leq 0 \end{aligned}$$

thanks to (20), since  $|x - y| + |z - r| \leq a$  implies that  $|x - y| \leq a$ . Therefore, we always have  $\Delta(x, z, y, r) \leq 0$  and the proof is complete.  $\square$

## 4 The multiple time renewal equation

Several applications use multi-time renewal equations to describe a population density subjected to aging or to time-evolution. Recently, for evaluating the efficiency of tracing softwares, it was used to take into account secondary infections, see [21]. In neuroscience, the interpretation is that neurons keep memory of their last spikes in the process of deciding when to fire again, see [12]. For two times memory, the equation reads

$$\begin{cases} \frac{\partial u_t(x_1, x_2)}{\partial t} + \frac{\partial u_t(x_1, x_2)}{\partial x_1} + \frac{\partial u_t(x_1, x_2)}{\partial x_2} + d(x_1, x_2)u_t(x_1, x_2) = 0, & x_2 \geq x_1 \geq 0, t \geq 0, \\ u_t(x_1 = 0, x_2) = \int_{x_1}^{\infty} d(x_2, z)u_t(x_2, dz), & x_2 \geq 0, t \geq 0. \end{cases} \quad (25)$$

An example of stochastic interpretation is that particles are individuals producing *events* at a rate depending on the ages of its two last events. Here the variables  $x_1$  and  $x_2$  represent the ages of the two last events, so these ages increase linearly until, at rate  $d(x_1, x_2)$ , they are reset to the values  $(0, x_1)$ . Hence our state space is now  $\mathcal{J} = \{(x_1, x_2) \in [0, \infty)^2 : x_2 > x_1\}$ . We assume that

$$d \in C(\mathcal{J}), \quad d \geq 0 \quad (26)$$

and that

$$\exists a > 0 \quad \text{such that} \quad a \leq \inf_{2|x_1 - \tilde{x}_1| + |x_2 - \tilde{x}_2| \leq a} \frac{(|x_1 - \tilde{x}_1| + |x_2 - \tilde{x}_2|) \max(d(x_1, x_2), d(\tilde{x}_1, \tilde{x}_2))}{|d(x_1, x_2) - d(\tilde{x}_1, \tilde{x}_2)|}. \quad (27)$$

One can check that  $d(x_1, x_2) = \alpha + \beta x_1^{p_1} + \gamma x_2^{p_2}$ , with  $\alpha > 0$ ,  $\beta \geq 0$ ,  $\gamma \geq 0$ ,  $p_1 \geq 1$  and  $p_2 \geq 1$  satisfies such an assumption, as well as Lipschitz and uniformly positive functions.

**Theorem 5** *We assume (26)-(27). We consider the cost on  $\mathcal{J} \times \mathcal{J}$  defined by*

$$\rho(x_1, x_2, \tilde{x}_1, \tilde{x}_2) = \min(a, 2|x_1 - \tilde{x}_1| + |x_2 - \tilde{x}_2|).$$

*For any  $u_0^1, u_0^2 \in \mathcal{P}(\mathcal{J})$ , there exists a pair of weak measure solutions  $(u_t^1)_{t \geq 0}, (u_t^2)_{t \geq 0} \subset \mathcal{P}(\mathcal{J})$  to (25) starting from  $u_0^1$  and  $u_0^2$ , i.e., such that for  $i = 1, 2$  and all  $t \geq 0$*

$$\int_0^t \int_{\mathcal{J}} d(x_1, x_2) u_s^i(dx_1, dx_2) ds < \infty \quad (28)$$

*and for  $i = 1, 2$ , all  $t \geq 0$  and all  $\varphi \in C_c^1(\mathcal{J})$ ,*

$$\begin{aligned} \int_{\mathcal{J}} \varphi(x_1, x_2) u_t^i(dx_1, dx_2) &= \int_{\mathcal{J}} \varphi(x_1, x_2) u_0^i(dx_1, dx_2) \\ &+ \int_0^t \int_{\mathcal{J}} \left[ \frac{\partial \varphi(x_1, x_2)}{\partial x_1} + \frac{\partial \varphi(x_1, x_2)}{\partial x_2} + d(x_1, x_2) [\varphi(0, x_1) - \varphi(x_1, x_2)] \right] u_s^i(dx_1, dx_2) ds. \end{aligned} \quad (29)$$

*Moreover, for all  $t \geq 0$ , we have*

$$\mathcal{T}_\rho(u_t^1, u_t^2) \leq \mathcal{T}_\rho(u_0^1, u_0^2).$$

**Proof.** We assume (26)-(27), consider  $u_0^1, u_0^2 \in \mathcal{P}(\mathcal{J})$  and a coupling  $v_0 \in \mathcal{H}(u_0^1, u_0^2)$ . There exists a family  $(v_t)_{t \geq 0}$  of probability measures on  $\mathcal{J}^2$  such that for all  $t \geq 0$

$$\int_0^t \iint [d(x_1, x_2) + d(\tilde{x}_1, \tilde{x}_2)] v_t(dx_1, dx_2, d\tilde{x}_1, d\tilde{x}_2) < \infty \quad (30)$$

and which weakly solves, with zero flux boundary conditions at  $x_1 = 0$  and  $x_2 = 0$ ,

$$\begin{aligned} \frac{\partial v_t}{\partial t} + \frac{\partial v_t}{\partial x_1} + \frac{\partial v_t}{\partial x_2} + \frac{\partial v_t}{\partial \tilde{x}_1} + \frac{\partial v_t}{\partial \tilde{x}_2} + (d(x_1, x_2) \vee d(\tilde{x}_1, \tilde{x}_2)) v_t \\ = \delta(x_1) \delta(\tilde{x}_1) \iint (d(x_2, z) \vee d(\tilde{x}_2, \tilde{z})) v_t(x_2, dz, \tilde{x}_2, d\tilde{z}) \\ + \delta(x_1) \iint (d(x_2, z) - d(\tilde{x}_1, \tilde{x}_2))_+ v_t(x_2, dz, \tilde{x}_1, \tilde{x}_2) \\ + \delta(\tilde{x}_1) \iint (d(x_1, x_2) - d(\tilde{x}_2, \tilde{z}))_+ v_t(x_1, x_2, \tilde{x}_2, d\tilde{z}). \end{aligned}$$

This means that for all  $\varphi \in C_c^1(\mathcal{J}^2)$ , all  $t \geq 0$ ,

$$\begin{aligned} \iint \varphi(x_1, x_2, \tilde{x}_1, \tilde{x}_2) v_t(dx_1, dx_2, d\tilde{x}_1, d\tilde{x}_2) = \iint \varphi(x_1, x_2, \tilde{x}_1, \tilde{x}_2) v_0(dx_1, dx_2, d\tilde{x}_1, d\tilde{x}_2) \\ + \int_0^t \iint \left[ \frac{\partial \varphi}{\partial x_1} + \frac{\partial \varphi}{\partial x_2} + \frac{\partial \varphi}{\partial \tilde{x}_1} + \frac{\partial \varphi}{\partial \tilde{x}_2} \right] (x_1, x_2, \tilde{x}_1, \tilde{x}_2) v_s(dx, dz, dy, dr) ds \\ + \int_0^t \iint \left( (d(x_1, x_2) \wedge d(\tilde{x}_1, \tilde{x}_2)) [\varphi(0, x_1, 0, \tilde{x}_1) - \varphi(x_1, x_2, \tilde{x}_1, \tilde{x}_2)] \right. \\ \left. + (d(x_1, x_2) - d(\tilde{x}_1, \tilde{x}_2))_+ [\varphi(0, x_1, \tilde{x}_1, \tilde{x}_2) - \varphi(x_1, x_2, \tilde{x}_1, \tilde{x}_2)] \right. \\ \left. + (d(\tilde{x}_1, \tilde{x}_2) - d(x_1, x_2))_+ [\varphi(x_1, x_2, 0, \tilde{x}_1) - \varphi(x_1, x_2, \tilde{x}_1, \tilde{x}_2)] \right) v_s(dx_1, dx_2, d\tilde{x}_2, d\tilde{x}_2) ds. \end{aligned} \quad (31)$$

Using (30) and (31) with a function  $\varphi$  depending only on  $(x_1, x_2)$  (or  $(\tilde{x}_1, \tilde{x}_2)$ ), we see that the marginals of  $v$  satisfy (28) and (29).

The existence for (30)-(31) follows as usual from the following *a priori* tightness estimate. By the de la Vallée Poussin theorem, there is a function  $h : [0, \infty) \rightarrow [0, \infty)$  such that  $\lim_{x \rightarrow \infty} h(x) = \infty$  and

$$C := \iint [h(x_1) + h(x_2) + h(\tilde{x}_1) + h(\tilde{x}_2)] v_0(dx_1, dx_2, d\tilde{x}_1, d\tilde{x}_2) < \infty.$$

Choosing moreover  $h$  smooth, satisfying  $h(0) = 0$  and  $0 \leq h' \leq 1$ , applying (31) with  $\varphi(x_1, x_2, \tilde{x}_1, \tilde{x}_2) = h(x_1) + h(x_2) + h(\tilde{x}_1) + h(\tilde{x}_2)$ , one easily concludes as usual that for all  $t \geq 0$ ,

$$\begin{aligned} \iint [h(x_1) + h(x_2) + h(\tilde{x}_1) + h(\tilde{x}_2)] v_t(dx_1, dx_2, d\tilde{x}_1, d\tilde{x}_2) \\ + \int_0^t \iint (d(x_1, x_2) h(x_2) + d(\tilde{x}_1, \tilde{x}_2) h(\tilde{x}_2)) v_s(dx_1, dx_2, d\tilde{x}_1, d\tilde{x}_2) ds \leq C + 4t. \end{aligned}$$

Recalling that  $h$  increases to infinity and that  $h(x_1) \leq h(x_2)$  for all  $(x_1, x_2) \in \mathcal{J}$ , this *a priori* tightness estimate is enough to prove existence for (30)-(31).

We now apply (24) with  $\varphi = \varrho$ , where  $\varrho(x_1, x_2, \tilde{x}_1, \tilde{x}_2) = a \wedge (2|x_1 - \tilde{x}_1| + |x_2 - \tilde{x}_2|)$ . Since

$$\frac{\partial \varphi}{\partial x_1} + \frac{\partial \varphi}{\partial x_2} + \frac{\partial \varphi}{\partial \tilde{x}_1} + \frac{\partial \varphi}{\partial \tilde{x}_2} = 0,$$

we find, choosing  $v_0$  such that  $\iint \varrho(x_1, x_2, \tilde{x}_1, \tilde{x}_2) v_0(dx_1, dx_2, d\tilde{x}_1, d\tilde{x}_2) = \mathcal{T}_\rho(u_0^1, u_0^2)$ ,

$$\iint \varrho(x_1, x_2, \tilde{x}_1, \tilde{x}_2) v_t(dx_1, dx_2, d\tilde{x}_1, d\tilde{x}_2) \leq \mathcal{T}_\rho(u_0^1, u_0^2) + \int_0^t \iint \Delta(x_1, x_2, \tilde{x}_1, \tilde{x}_2) v_s(dx_1, dx_2, d\tilde{x}_1, d\tilde{x}_2) ds,$$

where

$$\begin{aligned} \Delta(x_1, x_2, \tilde{x}_1, \tilde{x}_2) = & - (d(x_1, x_2) \vee d(\tilde{x}_1, \tilde{x}_2)) ([2|x_1 - \tilde{x}_1| + |x_2 - \tilde{x}_2|] \wedge a) \\ & + (d(x_1, x_2) \wedge d(\tilde{x}_1, \tilde{x}_2)) (|x_1 - \tilde{x}_1| \wedge a) \\ & + (d(x_1, x_2) - d(\tilde{x}_1, \tilde{x}_2))_+ ([2|\tilde{x}_1| + |x_1 - \tilde{x}_2|] \wedge a) \\ & + (d(\tilde{x}_1, \tilde{x}_2) - d(x_1, x_2))_+ ([2|x_1| + |\tilde{x}_1 - x_2|] \wedge a). \end{aligned}$$

Since  $\mathcal{T}_\rho(u_t^1, u_t^2) \leq \iint \rho(x_1, x_2, \tilde{x}_1, \tilde{x}_2) v_t(dx_1, dx_2, d\tilde{x}_1, d\tilde{x}_2)$ , it only remains to check that  $\Delta$  is always non-positive.

When first  $2|x_1 - \tilde{x}_1| + |x_2 - \tilde{x}_2| \geq a$ , this is obvious.

When  $2|x_1 - \tilde{x}_1| + |x_2 - \tilde{x}_2| \leq a$ , it suffices to verify that

$$\begin{aligned} & (d(x_1, x_2) \vee d(\tilde{x}_1, \tilde{x}_2)) [2|x_1 - \tilde{x}_1| + |x_2 - \tilde{x}_2|] \\ & \geq (d(x_1, x_2) \wedge d(\tilde{x}_1, \tilde{x}_2)) |x_1 - \tilde{x}_1| + |d(x_1, x_2) - d(\tilde{x}_1, \tilde{x}_2)| a. \end{aligned}$$

This follows from the fact that, by (27), we have

$$(d(x_1, x_2) \vee d(\tilde{x}_1, \tilde{x}_2)) [|x_1 - \tilde{x}_1| + |x_2 - \tilde{x}_2|] \geq |d(x_1, x_2) - d(\tilde{x}_1, \tilde{x}_2)| a.$$

□

## 5 Growth-fragmentation

The growth-fragmentation equation arises in several areas of biology. The variable represents for instance the size of cells or the length of biopolymers. A large literature is available on the subject and we refer for instance to [30, 15, 5, 33]. It also arises in communication science for TCP connections, see [35] and the references therein. In particular, in [11, 3], the authors prove some strict contraction in *classical* Wasserstein distance, for a less general model, by coupling the stochastic processes in a very similar way as we do.

The model combines growth with a rate  $g$  and fragmentation with a rate  $d$  and it is written

$$\begin{cases} \frac{\partial u_t(x)}{\partial t} + \frac{\partial [g(x) u_t(x)]}{\partial x} + d(x) u_t(x) = \int_x^\infty d(x') \kappa(x, x') u_t(dx'), & t \geq 0, x \geq 0, \\ u_t(x=0) = 0, & t \geq 0. \end{cases} \quad (32)$$

Usual conditions on the fragmentation kernel are expressed through the identities

$$\kappa(x, x') = 0 \text{ for } x > x', \quad \int_0^{x'} \kappa(x, x') dx = 1 \quad (33)$$

which lead to the conservation law

$$\int_0^\infty u_t(x)dx = \int_0^\infty u_0(x)dx = 1.$$

To go further, we can specify

$$\kappa(x, x') = \frac{1}{x'} \beta\left(\frac{x}{x'}\right) \quad \text{for some } \beta \in \mathcal{P}([0, 1]), \quad (34)$$

see the end of the section for a more general possible setting. We then assume that

$$g, d \in C([0, \infty)), \quad g \text{ is non-increasing, } \quad g(0) \geq 0, \quad d \geq 0, \quad (35)$$

and

$$\exists a > 0 \quad \text{such that} \quad a \leq \left(1 - \int_0^1 r\beta(dr)\right) \inf_{|x-y| \leq a} \frac{|x-y| \max(d(x), d(y))}{|d(x) - d(y)|}. \quad (36)$$

If  $\beta$  is non-trivial in that  $\int_0^1 r\beta(dr) \in [0, 1)$ , then this assumption is verified if e.g.  $d(x) = \alpha + \beta x^p$ , provided  $\alpha > 0$ ,  $\beta \geq 0$  and  $p \geq 1$ .

**Theorem 6** *We assume (34)-(35)-(36). We choose again, on  $[0, \infty) \times [0, \infty)$  the cost*

$$\varrho(x, y) = \min(a, |x - y|).$$

*For any  $u_0^1, u_0^2 \in \mathcal{P}([0, \infty))$ , there exists a pair of weak measure solutions  $(u_t^1)_{t \geq 0}, (u_t^2)_{t \geq 0} \subset \mathcal{P}([0, \infty))$  to (32), starting from  $u_0^1$  and  $u_0^2$ , i.e., such that for  $i = 1, 2$ , for all  $t \geq 0$ , all  $A \geq 1$ ,*

$$\int_0^t \int_{A+1}^\infty d(x) \int_0^{A/x} \beta(dr) u_s^i(dx) ds < \infty \quad (37)$$

*and for all  $t \geq 0$ , all  $\varphi \in C_c^1([0, \infty))$ ,*

$$\int_0^\infty \varphi(x) u_t^i(dx) = \int_0^\infty \varphi(x) u_0^i(dx) + \int_0^t \int_0^\infty \left[ g(x) \varphi'(x) + d(x) \int_0^1 (\varphi(rx) - \varphi(x)) \beta(dr) \right] u_s^i(dx) ds. \quad (38)$$

*Moreover, for all  $t \geq 0$ , we have*

$$\mathcal{T}_\varrho(u_t^1, u_t^2) \leq \mathcal{T}_\varrho(u_0^1, u_0^2).$$

Observe that (38) makes sense thanks to (37): for  $\varphi \in C_c^1([0, \infty))$  supported in  $[0, A]$ ,

$$\left| d(x) \int_0^1 (\varphi(rx) - \varphi(x)) \beta(dr) \right| \leq \mathbf{1}_{\{x \leq A+1\}} 2 \|\varphi\|_\infty \sup_{[0, A+1]} d + \mathbf{1}_{\{x > A+1\}} \|\varphi\|_\infty d(x) \int_0^{A/x} \beta(dr),$$

where we used a rough upper bound by  $A + 1$  to fit our assumption.

**Proof.** Consider  $u_0^1, u_0^2 \in \mathcal{P}([0, \infty))$  and a coupling  $v_0 \in \mathcal{H}(u_0^1, u_0^2)$ . There exists a family  $(v_t)_{t \geq 0}$  of probability measures on  $[0, \infty)^2$  satisfying, for all  $t \geq 0$ , all  $A \geq 1$ ,

$$\int_0^t \iint \left[ \mathbf{1}_{\{x \geq A+1\}} d(x) \int_0^{A/x} \beta(dr) + \mathbf{1}_{\{y \geq A+1\}} d(y) \int_0^{A/y} \beta(dr) \right] v_s(dx, dy) ds < \infty \quad (39)$$

and solving weakly

$$\begin{aligned} \frac{\partial v_t}{\partial t} + \frac{\partial}{\partial x} [g(x) v_t] + \frac{\partial}{\partial y} [g(y) v_t] + (d(x) \vee d(y)) v_t = \\ \iint (d(x') \wedge d(y')) \frac{1}{x'y'} \beta\left(\frac{x}{x'}\right) \delta\left(\frac{x}{x'} - \frac{y}{y'}\right) \beta\left(\frac{x}{x'}\right) v_t(dx', dy') \\ + \int (d(x') - d(y))_+ \frac{1}{x'} \beta\left(\frac{x}{x'}\right) v_t(dx', y) + \int (d(y') - d(x))_+ \frac{1}{y'} \beta\left(\frac{y}{y'}\right) v(x, dy'). \end{aligned}$$

This means that for all  $t \geq 0$ , all  $\varphi \in C_c^1([0, \infty)^2)$ ,

$$\begin{aligned} \iint \varphi(x, y) v_t(dx, dy) = \iint \varphi(x, y) v_0(dx, dy) + \int_0^t \iint \left[ g(x) \frac{\partial \varphi(x, y)}{\partial x} + g(y) \frac{\partial \varphi(x, y)}{\partial y} \right] v_s(dx, dy) ds \\ + \int_0^t \iint (d(x) \wedge d(y)) \int_0^1 [\varphi(rx, ry) - \varphi(x, y)] \beta(dr) v_s(dx, dy) ds \\ + \int_0^t \iint (d(x) - d(y))_+ \int_0^1 [\varphi(rx, y) - \varphi(x, y)] \beta(dr) v_s(dx, dy) ds \\ + \int_0^t \iint (d(y) - d(x))_+ \int_0^1 [\varphi(x, ry) - \varphi(x, y)] \beta(dr) v_s(dx, dy) ds. \end{aligned} \quad (40)$$

One checks as usual that the two marginals of  $v_t$  solve (37)-(38). Next, as in the previous sections, one finds that there is a function  $h : [0, \infty) \rightarrow [0, \infty)$ , strictly increasing to infinity, which taken as a test function gives

$$\iint [h(x) + h(y)] v_t(dx, dy) + \int_0^t \iint [d(x)H(x) + d(\tilde{x})H(\tilde{x})] v_s(dx, dy) ds \leq C + 2\bar{C}t,$$

where  $H(x) = \int_0^1 (h(x) - h(rx)) \beta(dr)$  and  $\bar{C} = \sup_{x \geq 0} g(x)h'(x)$ . This *a priori* estimate is sufficient to prove tightness and construct a solution to (37)-(38), because for any  $A \geq 1$ , any  $x \geq A + 1$ , one has

$$\int_0^{A/x} \beta(dr) \leq \int_0^{A/x} \frac{h(x) - h(rx)}{h(A+1) - h(A)} \beta(dr) \leq \frac{H(x)}{h(A+1) - h(A)}.$$

Applying now the above equation to  $\varphi = \varrho$ , where  $\varrho(x, y) = a \wedge |x - y|$ , one finds as usual, if choosing  $v_0$  correctly and using that  $g$  is non-increasing, that

$$\iint \varrho(x, y) v_t(dx, dy) \leq \mathcal{T}_\varrho(u_0^1, u_0^2) + \int_0^t \iint \Delta(x, y) v_s(dx, dy) ds,$$

where

$$\begin{aligned} \Delta(x, y) = - (d(x) \vee d(y)) \varrho(x, y) + (d(x) \wedge d(y)) \int_0^1 \varrho(rx, ry) \beta(dr) \\ + (d(x) - d(y))_+ \int_0^1 \varrho(rx, y) \beta(dr) + (d(y) - d(x))_+ \int_0^1 \varrho(x, ry) \beta(dr). \end{aligned} \quad (41)$$

We finally show that  $\Delta$  is always non-positive. First, when  $|x - y| \geq a$ , it is enough to check that

$$a(d(x) \vee d(y)) \geq a(d(x) \wedge d(y)) + a(d(x) - d(y))_+ + a(d(y) - d(x))_+,$$



which is obvious. Next, when  $|x - y| \leq a$ , it suffices to check that

$$(d(x) \vee d(y))|x - y| \geq (d(x) \wedge d(y)) \left[ \int_0^1 r\beta(dr) \right] |x - y| + a(d(x) - d(y))_+ + a(d(y) - d(x))_+,$$

which follows from the fact that, thanks to (36),

$$\left[ 1 - \int_0^1 r\beta(dr) \right] (d(x) \vee d(y))|x - y| \geq a|d(x) - d(y)|.$$

□

Here it does not seem possible to find a general condition under which one can obtain a strict contraction. As already mentioned, a strict contraction was obtained in [11, 3], for a specific model, with the usual  $W_1$ -Wasserstein distance, i.e. with the choice  $a = \infty$ .

A little study shows that the result still holds true if assuming, instead of (34), that the family of fragmentation kernels  $(\kappa(\cdot, x))_{x \in \mathbb{R}_+} \subset \mathcal{P}(\mathbb{R}_+)$  satisfies  $\kappa([0, x], x) = 1$  for all  $x \geq 0$  and

$$\exists m \in [0, 1), \quad \text{for all } x, y \in \mathbb{R}_+, \quad W_1(\kappa(\cdot, x), \kappa(\cdot, y)) \leq m|x - y|,$$

where  $W_1$  is the usual Monge-Kantorovich distance on  $\mathcal{P}(\mathbb{R}_+)$ , and replacing (36) by

$$\exists a > 0 \quad \text{such that} \quad a \leq (1 - m) \inf_{|x-y| \leq a} \frac{|x - y| \max(d(x), d(y))}{|d(x) - d(y)|}.$$

Indeed, it suffices to apply the usual strategy, starting from the coupling equation

$$\begin{aligned} \iint \varphi(x, y)v_t(dx, dy) &= \iint \varphi(x, y)v_0(dx, dy) + \int_0^t \iint \left[ g(x) \frac{\partial \varphi(x, y)}{\partial x} + g(y) \frac{\partial \varphi(x, y)}{\partial y} \right] v_s(dx, dy) ds \\ &+ \int_0^t \iint (d(x) \wedge d(y)) \int_0^x \int_0^y [\varphi(x', y') - \varphi(x, y)] \bar{\kappa}(dx', dy', x, y) v_s(dx, dy) ds \\ &+ \int_0^t \iint (d(x) - d(y))_+ \int_0^x [\varphi(x', y) - \varphi(x, y)] \kappa(dx', x) v_s(dx, dy) ds \\ &+ \int_0^t \iint (d(y) - d(x))_+ \int_0^y [\varphi(x, y') - \varphi(x, y)] \kappa(dy', y) v_s(dx, dy) ds, \end{aligned}$$

where for each  $x, y \in \mathbb{R}_+$ ,  $\bar{\kappa}(\cdot, \cdot, x, y) \in \mathcal{H}(\kappa(\cdot, x), \kappa(\cdot, y))$  satisfies

$$\iint |x' - y'| \bar{\kappa}(dx', dy', x, y) = W_1(\kappa(\cdot, x), \kappa(\cdot, y)).$$

## 6 Age and size structure

Models have been proposed which use several stucture variables. For instance, age or size only are not enough to predict cell division. But a combination of both (or other physiological variables as size increment) have been used, see [40, 19], leading to write

$$\left\{ \begin{array}{ll} \frac{\partial u_t(x, z)}{\partial t} + \frac{\partial u_t(x, z)}{\partial x} + \frac{\partial [g(z) u_t(x, z)]}{\partial z} + d(x, z) u_t(x, z) = 0, & t \geq 0, x > 0, z > 0, \\ u_t(x, z = 0) = 0, & t \geq 0, x > 0, \\ u_t(x = 0, z) = \int_{x=0}^{\infty} \int_{z'=z}^{\infty} d(x', z') \kappa(z, z') u_t(dx', dz'), & t \geq 0, z > 0 \end{array} \right. \quad (42)$$

The state space is  $\mathcal{J} = [0, \infty)^2$ . We assume that the fragmentation kernel has the specific form (34), that

$$g \in C([0, \infty)), \quad g \text{ is non-increasing,} \quad g(0) \geq 0, \quad d \in C(\mathcal{J}), \quad d \geq 0. \quad (43)$$

and that

$$\exists a > 0 \quad \text{such that} \quad a \leq \left(1 - \int_0^1 r\beta(dr)\right) \inf_{|x-z|+|\tilde{x}-\tilde{z}|\leq a} \frac{|x-\tilde{x}|+|z-\tilde{z}|}{|d(x,z)-d(\tilde{x},\tilde{z})|} \max(d(x,z), d(\tilde{x},\tilde{z})). \quad (44)$$

**Theorem 7** Assume (34)-(43)-(44) and consider the cost

$$\varrho(x, z, \tilde{x}, \tilde{z}) = \min(a, |x - \tilde{x}| + |z - \tilde{z}|).$$

For any  $u_0^1, u_0^2 \in \mathcal{P}(\mathcal{J})$ , there exists a pair of weak measure solutions  $(u_t^1)_{t \geq 0}, (u_t^2)_{t \geq 0} \subset \mathcal{P}(\mathcal{J})$  to (42), starting from  $u_0^1$  and  $u_0^2$ , i.e., such that for  $i = 1, 2$  and all  $t \geq 0$ , all  $A \geq 1$ ,

$$\int_0^t \int_{\mathcal{J}} d(x, z) \left[ \mathbf{1}_{\{x \geq A+1\}} + \mathbf{1}_{\{x \leq A+1, z \geq A+1\}} \int_0^{A/z} \beta(dr) \right] u_s^i(dx, dz) ds < \infty \quad (45)$$

and for all  $t \geq 0$ , all  $\varphi \in C_c^1(\mathcal{J})$ ,

$$\begin{aligned} \int_{\mathcal{J}} \varphi(x, z) u_t^i(dx, dz) &= \int_{\mathcal{J}} \varphi(x, z) u_0^i(dx, dz) \\ &+ \int_0^t \int_{\mathcal{J}} \left[ \frac{\partial \varphi(x, z)}{\partial x} + g(z) \frac{\partial \varphi(x, z)}{\partial z} + d(x, z) \int_0^1 (\varphi(0, rz) - \varphi(x, z)) \beta(dr) \right] u_s^i(dx, dz) ds. \end{aligned} \quad (46)$$

Moreover, for all  $t \geq 0$ , we have

$$\mathcal{T}_{\varrho}(u_t^1, u_t^2) \leq \mathcal{T}_{\varrho}(u_0^1, u_0^2).$$

Here again, (46) makes sense thanks to (45): if  $\varphi \in C_c(\mathcal{J})$  is supported in  $[0, A]^2$ , then

$$\begin{aligned} d(x, z) \left| \int_0^1 (\varphi(0, rz) - \varphi(x, z)) \beta(dr) \right| &\leq \mathbf{1}_{\{x \leq A+1, z \leq A+1\}} 2 \|\varphi\|_{\infty} \sup_{[0, A+1]^2} d + \mathbf{1}_{\{x \geq A+1\}} \|\varphi\|_{\infty} d(x, z) \\ &+ \mathbf{1}_{\{x \leq A+1, z \geq A+1\}} \|\varphi\|_{\infty} d(x, z) \int_0^{A/z} \beta(dr). \end{aligned}$$

**Proof.** As before, we consider  $u_0^1, u_0^2 \in \mathcal{P}(\mathcal{J})$  and a coupling  $v_0 \in \mathcal{H}(u_0^1, u_0^2)$ . There exists a family  $(v_t)_{t \geq 0}$  of probability measures on  $\mathcal{J}^2$  such that, for all  $t \geq 0$ ,

$$\begin{aligned} \int_0^t \int_{\mathcal{J}} \left( d(x, z) \left[ \mathbf{1}_{\{x \geq A+1\}} + \mathbf{1}_{\{x \leq A+1, z \geq A+1\}} \int_0^{A/z} \beta(dr) \right] \right. \\ \left. + d(\tilde{x}, \tilde{z}) \left[ \mathbf{1}_{\{\tilde{x} \geq A+1\}} + \mathbf{1}_{\{\tilde{x} \leq A+1, \tilde{z} \geq A+1\}} \int_0^{A/\tilde{z}} \beta(dr) \right] \right) v_s(dx, dz, d\tilde{x}, d\tilde{z}) ds < \infty \end{aligned} \quad (47)$$

and that weakly solves

$$\begin{aligned}
& \frac{\partial}{\partial t} v_t(x, z, \tilde{x}, \tilde{z}) + \frac{\partial v_t}{\partial x} + \frac{\partial v_t}{\partial \tilde{x}} + \frac{\partial}{\partial z} [g(z) v_t] + \frac{\partial v_t}{\partial \tilde{z}} [g(\tilde{z}) v_t] + b(x, z) \vee d(\tilde{x}, \tilde{z}) v_t \\
&= \delta(x) \delta(\tilde{x}) \iint (d(x', z') \wedge d(\tilde{x}', \tilde{z}')) \frac{1}{z' \tilde{z}'} \beta\left(\frac{z}{z'}\right) \delta\left(\frac{z}{z'} - \frac{\tilde{z}}{\tilde{z}'}\right) v_t(dx', dz', d\tilde{x}', d\tilde{z}') \\
&+ \delta(x) \int (d(x', z') - d(\tilde{x}, \tilde{z}))_+ \frac{1}{z'} \beta\left(\frac{z}{z'}\right) v_t(dx', dz', \tilde{x}, \tilde{z}) \\
&+ \delta(\tilde{x}) \int (d(\tilde{x}', \tilde{z}') - d(x, z))_+ \frac{1}{\tilde{z}'} \beta\left(\frac{\tilde{z}}{\tilde{z}'}\right) v_t(x, z, d\tilde{x}', d\tilde{z}')
\end{aligned}$$

This equation means that for all  $t \geq 0$ , for all  $\varphi \in C_c^1(\mathcal{J}^2)$ ,

$$\begin{aligned}
& \iint \varphi(x, z, \tilde{x}, \tilde{z}) v_t(dx, dz, d\tilde{x}, d\tilde{z}) = \iint \varphi(x, z, \tilde{x}, \tilde{z}) v_0(dx, dz, d\tilde{x}, d\tilde{z}) \\
&+ \int_0^t \iint \left[ \frac{\partial \varphi}{\partial x} + \frac{\partial \varphi}{\partial \tilde{x}} + g(z) \frac{\partial \varphi}{\partial z} + g(\tilde{z}) \frac{\partial \varphi}{\partial \tilde{z}} \right] (x, z, \tilde{x}, \tilde{z}) v_s(dx, dz, d\tilde{x}, d\tilde{z}) ds \\
&+ \int_0^t \iint \left[ (d(x, z) \wedge d(\tilde{x}, \tilde{z})) \int_0^1 [\varphi(0, rz, 0, r\tilde{z}) - \varphi(x, z, \tilde{x}, \tilde{z})] \beta(dr) \right. \\
&\quad \left. + (d(x, z) - d(\tilde{x}, \tilde{z}))_+ \int_0^1 [\varphi(0, rz, \tilde{x}, \tilde{z}) - \varphi(x, z, \tilde{x}, \tilde{z})] \beta(dr) \right. \\
&\quad \left. + (d(\tilde{x}, \tilde{z}) - d(x, z))_+ \int_0^1 [\varphi(x, z, 0, r\tilde{z}) - \varphi(x, z, \tilde{x}, \tilde{z})] \beta(dr) \right] v_s(dx, dz, d\tilde{x}, d\tilde{z}) ds. \tag{48}
\end{aligned}$$

The two marginals of  $v_t$  solve (45)-(46). Next, one finds as usual that there is a function  $h : [0, \infty) \rightarrow [0, \infty)$ , strictly increasing to infinity, such that

$$\begin{aligned}
& \iint [h(x) + h(z) + h(\tilde{x}) + h(\tilde{z})] v_t(dx, dz, d\tilde{x}, d\tilde{z}) \\
&+ \int_0^t \iint [d(x, z) H(x, z) + d(\tilde{x}, \tilde{z}) H(\tilde{x}, \tilde{z})] v_s(dx, dz, d\tilde{x}, d\tilde{z}) ds \leq C + 2(1 + \bar{C})t,
\end{aligned}$$

where  $H(x, z) = h(x) + \int_0^1 (h(x) - h(rx)) \beta(dr)$  and  $\bar{C} = \sup_{x \geq 0} g(x) h'(x)$ . This *a priori* estimate is sufficient to construct a solution to (47)-(48), because for any  $A \geq 1$

$$\mathbf{1}_{\{x \geq A+1\}} + \mathbf{1}_{\{x \leq A+1, z \geq A+1\}} \int_0^{A/z} \beta(dr) \leq \frac{h(x)}{h(A+1)} + \int_0^1 \frac{h(x) - h(rx)}{h(A+1) - h(A)} \beta(dr) \leq \frac{H(x)}{h(A+1) - h(A)}.$$

Following the usual procedure to prove the decay property, that we do not repeat again, it suffices, to conclude the proof, to check that

$$\begin{aligned}
& (d(x, z) \vee d(\tilde{x}, \tilde{z})) [a \wedge (|x - \tilde{x}| + |z - \tilde{z}|)] \\
&\geq (d(x, z) \wedge d(\tilde{x}, \tilde{z})) \int_0^1 a \wedge (r|z - \tilde{z}|) \beta(dr) + a |d(x, z) - d(\tilde{x}, \tilde{z})|.
\end{aligned}$$

When the min on the left hand side is achieved by  $a$  the inequality is obvious. Otherwise, we have to check that

$$(d(x, z) \vee d(\tilde{x}, \tilde{z})) [|x - \tilde{x}| + |z - \tilde{z}|] \geq \int_0^1 r \beta(dr) (d(x, z) \wedge d(\tilde{x}, \tilde{z})) |z - \tilde{z}| + a |d(x, z) - d(\tilde{x}, \tilde{z})|.$$

This again is satisfied if

$$\left(1 - \int_0^1 r\beta(dr)\right)(d(x, z) \vee d(\tilde{x}, \tilde{z}))[|x - \tilde{x}| + |z - \tilde{z}|] \geq a |d(x, z) - d(\tilde{x}, \tilde{z})|,$$

which is the condition (44), recall that we are in the case where  $|x - \tilde{x}| + |z - \tilde{z}| \leq a$ .  $\square$

## 7 Sexually structured populations

Here a female of type  $x'$  mates with a male of type  $x'_*$ , chosen with the probability  $u_t(\cdot)$ , the newborn is distributed with type  $x$  according to the law  $K(x; x', x'_*)$ . As often in this theory, we assume the distribution of males and females are identical, and rely on the formalism which can be found in [8, 32, 10] for instance. Using transport costs for nonlinear equations is not as standard as for linear ones, the closest known case is the homogeneous Boltzmann equation with Maxwellian collisions, see [23] for a simple proof based on coupling with the same fundamental method of 'common jumps'. For the problem at hand, an approach by Kantorovich duality is available in [16]. The (homogeneous) model reads

$$\partial_t u_t(x) + u_t(x) = \iint_{\mathbb{R}^{2d}} K(x; x', x'_*) u_t(dx') u_t(dx'_*), \quad t \geq 0, x \in \mathbb{R}^d. \quad (49)$$

For keeping the total population constant, the kernel  $K \geq 0$  satisfies

$$\iint_{\mathbb{R}^d} K(dx; x', x'_*) = 1.$$

For instance, we can think of two extreme cases of either a Dirac concentration or a uniform distribution,

$$K(x; x', x'_*) = \delta_{\theta x' + (1-\theta)x'_*}(x), \quad \theta \in (0, 1), \quad \text{or} \quad K(x; x', x'_*) = \frac{1}{|x' - x'_*|} \mathbf{1}_{\{x \in (x', x'_*)\}}. \quad (50)$$

These distributions can be generalized to the form

$$K(dx; x', x'_*) = \int_0^1 \delta_{x'\sigma + x'_*(1-\sigma)}(x) h(d\sigma), \quad (51)$$

with  $h$  a probability distribution on  $[0, 1]$  such that  $\int_0^1 \sigma h(d\sigma) = \theta \in (0, 1)$ , which is the form we use in the sequel.

**Theorem 8** *With the notations and assumptions above, we choose, for some  $p \geq 1$ ,*

$$\varrho(x, y) = |x - y|^p.$$

*For any  $u_0^1, u_0^2 \in \mathcal{P}_p(\mathbb{R}^d)$ , there exists a pair of weak measure solutions  $(u_t^1)_{t \geq 0}, (u_t^2)_{t \geq 0} \subset \mathcal{P}_p(\mathbb{R}^d)$  to (49), starting from  $u_0^1$  and  $u_0^2$ , i.e., such that for  $i = 1, 2$ , all  $t \geq 0$  and all  $\varphi \in C_c^1(\mathbb{R}^d)$ ,*

$$\begin{aligned} \int_{\mathbb{R}^d} \varphi(x) u_t^i(dx) &= \int_{\mathbb{R}^d} \varphi(x) u_0^i(dx) \\ &+ \int_0^t \iiint [\varphi(x) - \theta \varphi(x') - (1-\theta) \varphi(x'_*)] K(dx; x', x'_*) u_s^i(dx', dx'_*) ds. \end{aligned} \quad (52)$$

Moreover, for all  $t \geq 0$ , we have

$$\mathcal{T}_\rho(u_t^1, u_t^2) \leq \mathcal{T}_\rho(u_0^1, u_0^2).$$

**Proof.** We use a coupling  $\bar{K}(x, y; x', x'_*, y', y'_*)$ , to be chosen later, with the property

$$\int_{\mathbb{R}^d} \bar{K}(x, dy; x', x'_*, y', y'_*) = K(x; x', x'_*), \quad \int_{\mathbb{R}^d} \bar{K}(dx, y; x', x'_*, y', y'_*) = K(y; y', y'_*).$$

Then, we introduce the coupling equation

$$\partial_t v_t(x, y) + v_t(x, y) = \iiint \bar{K}(x, y; x', x'_*, y', y'_*) v_t(dx', dy') v_t(dx'_*, dy'_*).$$

This means that for all  $t \geq 0$ , for all  $\varphi \in C_c^1(\mathbb{R}^{2d})$ ,

$$\begin{aligned} \int_{\mathbb{R}^d} \varphi(x, y) v_t(dx, dy) &= \int_{\mathbb{R}^d} \varphi(x, y) v_0(dx, dy) + \int_0^t \iiint \left[ \varphi(x, y) - \theta \varphi(x', y') - (1 - \theta) \varphi(x'_*, y'_*) \right] \\ &\quad \bar{K}(dx, dy; x', x'_*, y', y'_*) v_s(dx', dy') v_s(dx'_*, dy'_*) ds, \end{aligned} \quad (53)$$

where  $\theta \in (0, 1)$  is the same as in (51). For existence, we have to check the tightness in  $\mathcal{P}_p(\mathbb{R}^d)$ . By the de la Vallée Poussin theorem, there is a function  $h : \mathbb{R}^d \rightarrow [0, \infty)$  such that  $\lim_{|x| \rightarrow \infty} h(x) = \infty$  and

$$\bar{C} := \iint [h(x)|x|^p + h(y)|y|^p] v_0(dx, dy) < \infty.$$

One can moreover choose  $h$  smooth, satisfying  $h(0) = 0$  and such that  $x \mapsto h(x)|x|^p$  convex. Choosing  $h(x)|x|^p + h(y)|y|^p$  as a test function in (53), we conclude the bound, for all  $t \geq 0$ ,

$$\iint [h(x)|x|^p + h(y)|y|^p] v_t(dx, dy) \leq \bar{C},$$

because, by convexity, the second term in the right hand side of (53) is non-positive.

For the non-expansion property, we just have to show that the right hand side is non-positive for  $\rho(x, y) = |x - y|^p$  (arguing again after truncation, regularization), if choosing as coupling kernel

$$\bar{K}(dx, dy; x', x'_*, y', y'_*) = \int_0^1 h(d\sigma) \delta_{\sigma x' + (1-\sigma)x'_*}(x) \delta_{\sigma y' + (1-\sigma)y'_*}(y).$$

The duality formula raises the condition

$$\int_0^1 |\sigma x' + (1 - \sigma)x'_* - \sigma y' - (1 - \sigma)y'_*|^p h(d\sigma) \leq \theta |x' - y'|^p + (1 - \theta) |x'_* - y'_*|^p$$

which, by convexity, is immediate. □

## A Uniqueness of measure solutions

The coupling method is most powerful when the measure solutions are unique. This uniqueness problem, in particular for coefficients with low regularity, can lead to several deep developments, [2, 15]. Here, we consider regular coefficients so that the Hilbert Uniqueness Method can be applied without difficulty both to the Structured Equations under consideration and to the coupled equations. We treat in details the example of the renewal equation, i.e., (2) when  $b = \delta$ , a case which can also be found in [24]. We assume that

$$d \in C([0, \infty)), \quad g(x) \in C_b^1(\mathbb{R}^+), \quad g(0) \geq 0, \quad (54)$$

We define the weak solutions (or distributional solutions), as follows.

**Definition 9** *A function  $(u_t)_{t \geq 0} \subset \mathcal{P}(0, \infty)$  satisfies the renewal equation (2) in the distribution sense, if for all  $T > 0$  and all test function  $\psi \in C_{\text{comp}}^1([0, T] \times [0, \infty])$  such that  $\psi(x, T) \equiv 0$ , we have*

$$-\int_0^T \int_0^\infty \left[ \frac{\partial \psi(x, t)}{\partial t} + g(x) \frac{\partial \psi(x, t)}{\partial x} - d(x) \psi(x, t) + d(x) \psi(0, t) \right] u_t(dx) dt = \int_0^\infty \psi(x, 0) u_0(dx).$$

**Theorem 10 (Well posedness)** *We assume (54). There is a unique weak solution of the renewal equation (2).*

For the existence part, we refer to [2, 15] where more elaborate equations are treated. For uniqueness, we need to study the inhomogeneous dual problem. We introduce a source term  $S(x, t)$  on a given time interval  $[0, T]$  and

$$\begin{cases} -\frac{\partial}{\partial t} \psi(x, t) - g(x) \frac{\partial}{\partial x} \psi(x, t) + d(x) \psi(x, t) = \psi(0, t) d(x) + S(x, t), \\ \psi(x, T) = 0. \end{cases} \quad (55)$$

This problem is backward in  $t$  and  $x$ , therefore it does not use a boundary condition at  $x = 0$ .

**Lemma 11 (Existence for the dual problem)** *Assume (54),  $S \in C_{\text{comp}}^1([0, T] \times \mathbb{R}^+)$  and  $d \in C^1(0, \infty)$ , then there is a unique  $C^1$  solution to the dual equation (55). Moreover  $\psi(x, t)$  vanishes for  $x \geq R > 0$  for some  $R$  depending on the data and  $T$ , and the bound holds*

$$\sup_{0 \leq t \leq T, x \in \mathbb{R}^+} |\psi(x, t)| \leq C(T) \|S\|_\infty.$$

**Proof.** We use the method of characteristics based on the solution of the differential system parametrized by the Cauchy data  $(x, t)$  which is fixed

$$\begin{cases} \frac{d}{ds} X_s = g(X_s), & 0 \leq s \leq T, \\ X_t = x \geq 0. \end{cases}$$

It is well-posed thanks to the Cauchy-Lipschitz theorem and  $X_s \geq 0$  thanks to assumption  $g(0) \geq 0$ . It might be useful to keep in mind that  $X_s$  depends on  $(x, t)$  and thus the notation  $X_s \equiv X_s(x, t)$ .

Then, we set

$$\begin{aligned}\tilde{\psi}(s; x, t) &= \psi(s, X_s) e^{\int_s^t d(\sigma, X_\sigma) d\sigma}, & \tilde{d}(s; x, t) &= d(s, X_s) e^{\int_s^t d(\sigma, X_\sigma) d\sigma}, \\ \tilde{S}(s; x, t) &= S(s, X_s) e^{\int_s^t d(\sigma, X_\sigma) d\sigma},\end{aligned}$$

and ignore the parameter  $(x, t)$  when the statements are clear enough. We rewrite equation (55) as

$$\begin{aligned}\frac{d}{ds} \tilde{\psi}(s) &= \left[ \frac{\partial}{\partial t} \psi + g \frac{\partial}{\partial x} \psi - d \psi \right] e^{\int_s^t d(\sigma, X_\sigma) d\sigma} \Big|_{(s, X_s)} \\ &= -\psi(0, s) \tilde{d}(s) - \tilde{S}(s),\end{aligned}$$

Next, we integrate between  $s = t$  and  $s = T$ , use the Cauchy data at  $t = T$  and the identity  $\tilde{\psi}(t) = \psi(x, t)$ , and we obtain

$$\psi(x, t) = \int_t^T [\psi(0, s) \tilde{d}(s; x, t) + \tilde{S}(s; x, t)] ds. \quad (56)$$

This integral equation can be solved first for  $x = 0$ . Then, equation (56) is reduced to the Volterra equation

$$\psi(0, t) = \int_t^T [\psi(0, s) \tilde{d}(s; 0, t) + \tilde{S}(s; 0, t)] ds, \quad 0 \leq t \leq T$$

which, thanks to the (backward) Cauchy-Lipschitz theorem, has a unique solution that vanishes for  $t = T$ . By the  $C^1$  regularity of the data, we also have  $\psi(0, t) \in C^1([0, T])$ .

Since  $\psi(0, t)$  is now known, formula (56) gives us the explicit form of the solution for all  $(x, t)$ . Notice that, in the compact support statement,  $\tilde{\psi}(x, t)$  vanishes for  $x \geq R$  where  $R$  denotes the size of the support of  $S$  in  $x$ , plus  $T \|g\|_\infty$ . The uniform bound on  $\psi$  also follows from formula (56),  $\square$

**Proof. [Uniqueness for the renewal equation.]** With the help of the dual problem, we can use the Hilbert Uniqueness Method. The idea is simple: when the coefficients  $d, g$  satisfy the assumptions of Lemma 11, we can use the solution  $\psi$  of (55) as a test function in the weak formulation of Definition 9. For the difference  $u = u^2 - u^1$  between two possible solutions  $u^2, u^1$  with the same initial data, we arrive at

$$\int_0^T \int_0^\infty \left[ \frac{\partial \psi(x, t)}{\partial t} + g(x) \frac{\partial \psi(x, t)}{\partial x} - d(x) \psi(x, t) + d(x) \psi(0, t) \right] u_t(dx) dt = 0.$$

for  $\psi(\cdot, \cdot) \in C^1$  which is the case when  $d \in C^1$ . Then, taking into account (55), we arrive at

$$\int_0^T \int_0^\infty S(x, t) u_t(dx) dt = 0,$$

for all  $T > 0$  and all functions  $S \in C_{\text{comp}}^1$ , and this implies  $u \equiv 0$ .

When  $d$  is merely continuous, we consider a regularized family  $d_p \rightarrow d$  where the convergence holds locally uniformly. Then, for a given function  $S \in C_{\text{comp}}^1$ , we solve (55) with  $d_p$  in place of  $d$  and call  $\psi_p$  its solution (which is uniformly bounded with compact support). Inserting it in the definition of weak solutions, we obtain

$$\int_0^T \int_0^\infty S(x, t) u_t(dx) dt = R_p,$$

$$R_p = \int_0^T \int_0^\infty [d_p - d(x)][\psi_p(x, t) - \psi_p(0, t)]u_t(dx) dt,$$

and using that  $\psi_p$  is uniformly bounded, we deduce that

$$|R_p| \leq T \|\psi\|_\infty \|d_p - d\|_\infty \xrightarrow{p \rightarrow \infty} 0.$$

Therefore, we have recovered the identity  $\int_0^T \int_0^\infty S(x, t)u_t(dx)dt = 0$ , for all functions  $S \in C_{\text{comp}}^1$ , and this implies again  $u \equiv 0$ .

This concludes the uniqueness result stated in Theorem 10.  $\square$

## B Non-conservative equations

To treat non-conservative equations, several routes are possible. Using transport costs between measures with different masses is a possibility, see [13, 25] and the references therein, but the adaptation has to be worked out. Another strategy is to transform the equation by a change of unknown using the eigenelements, as it is performed for the method of relative entropy [38, 31]. We give two examples.

The first one concerns the general form of the renewal equation (2)

$$\begin{cases} \frac{\partial u_t(x)}{\partial t} + \frac{\partial [g(x)u_t(x)]}{\partial x} + d(x)u_t(x) = b(x)N(t), & t \geq 0, x \geq 0, \\ u_t(x=0) = 0, & N(t) = \int_0^\infty c(x)u_t(dx), & t \geq 0. \end{cases} \quad (57)$$

with  $c$  nonnegative, bounded and continuous. Assume that there is a first eigenvalue  $\lambda$  and a positive dual eigenfunction  $\psi(x)$ , that means

$$-g(x)\frac{\partial \psi(x)}{\partial x} + (d(x) + \lambda)\psi(x) = c(x) \int_0^\infty b(x')\psi(x')dx'.$$

Then, the new unknown  $w_t = u_t\psi e^{-\lambda t}$  satisfies, with  $\bar{c} = \frac{c}{\psi}$  and  $\bar{b} = b\psi$ , the conservative equation

$$\begin{cases} \frac{\partial w_t(x)}{\partial t} + \frac{\partial [g(x)w_t(x)]}{\partial x} + \int_0^\infty \bar{b}(x')dx' \bar{c}(x)w_t(x) = \bar{b}(x) \int_0^\infty \bar{c}(x')w_t(x')dx', & t \geq 0, x \geq 0, \\ w_t(x=0) = 0, & t \geq 0. \end{cases}$$

A non-conservative growth fragmentation equation (32) can be treated in a similar way departing from

$$\begin{cases} \frac{\partial u_t(x)}{\partial t} + \frac{\partial [g(x)u_t(x)]}{\partial x} + d(x)u_t(x) = \int_x^\infty c(x')\kappa(x, x')u_t(dx'), & t \geq 0, x \geq 0, \\ u_t(x=0) = 0, & t \geq 0, \end{cases} \quad (58)$$

with  $c$  nonnegative, bounded and continuous. The same construction and notations lead to the conservative equation

$$\begin{cases} \frac{\partial w_t(x)}{\partial t} + \frac{\partial [g(x)w_t(x)]}{\partial x} + \int_0^x \kappa(x', x)\psi(x')dx' \bar{c}(x)w_t(x) = \psi(x) \int_x^\infty \bar{c}(x')\kappa(x, x')w_t(x')dx', & t \geq 0, x \geq 0, \\ w_t(x=0) = 0, & t \geq 0. \end{cases}$$



## References

- [1] L. AMBROSIO, N. GIGLI, AND G. SAVARÉ, *Gradient flows in metric spaces and in the space of probability measures*, Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, second ed., 2008.
- [2] V. BANSAYE, B. CLOEZ, AND P. GABRIEL, *Ergodic behavior of non-conservative semigroups via generalized Doeblin's conditions*, Acta Appl. Math., 166 (2020), pp. 29–72.
- [3] J.-B. BARDET, A. CHRISTEN, A. GUILLIN, F. MALRIEU, AND P.-A. ZITT, *Total variation estimates for the TCP process*, Electron. J. Probab., 18 (2013), pp. no. 10, 21.
- [4] H. BERRY, T. LEPOUTRE, AND A. M. GONZÁLEZ, *Quantitative convergence towards a self-similar profile in an age-structured renewal equation for subdiffusion*, Acta Appl. Math., 145 (2016), pp. 15–45.
- [5] J. BERTOIN AND A. R. WATSON, *The strong Malthusian behavior of growth-fragmentation processes*, Ann. H. Lebesgue, 3 (2020), pp. 795–823.
- [6] S. BIANCHINI AND M. GLOYER, *An estimate on the flow generated by monotone operators*, Comm. Partial Differential Equations, 36 (2011), pp. 777–796.
- [7] F. BOUCHUT, F. JAMES, AND S. MANCINI, *Uniqueness and weak stability for multi-dimensional transport equations with one-sided Lipschitz coefficient*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), 4 (2005), pp. 1–25.
- [8] R. BÜRGER, *The mathematical theory of selection, recombination, and mutation*, Wiley Series in Mathematical and Computational Biology, John Wiley & Sons, Ltd., Chichester, 2000.
- [9] V. CALVEZ, P. GABRIEL, AND A. MATEOS GONZÁLEZ, *Limiting Hamilton-Jacobi equation for the large scale asymptotics of a subdiffusion jump-renewal equation*, Asymptot. Anal., 115 (2019), pp. 63–94.
- [10] V. CALVEZ, J. GARNIER, AND F. PATOUT, *Asymptotic analysis of a quantitative genetics model with nonlinear integral operator*, J. Éc. polytech. Math., 6 (2019), pp. 537–579.
- [11] D. CHAFAÏ, F. MALRIEU, AND K. PAROUX, *On the long time behavior of the TCP window size process*, Stochastic Process. Appl., 120 (2010), pp. 1518–1534.
- [12] J. CHEVALLIER, M. J. CÁCERES, M. DOUMIC, AND P. REYNAUD-BOURET, *Microscopic approach of a time elapsed neural model*, Math. Models Methods Appl. Sci., 25 (2015), pp. 2669–2719.
- [13] L. CHIZAT, G. PEYRÉ, B. SCHMITZER, AND F.-X. VIALARD, *Unbalanced optimal transport: dynamic and Kantorovich formulations*, J. Funct. Anal., 274 (2018), pp. 3090–3123.
- [14] J. M. CUSHING, *An introduction to structured population dynamics*, vol. 71 of CBMS-NSF Regional Conference Series in Applied Mathematics, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1998.

- [15] T. DEBIEC, M. DOUMIC, P. GWIAZDA, AND E. WIEDEMANN, *Relative entropy method for measure solutions of the growth-fragmentation equation*, SIAM J. Math. Anal., 50 (2018), pp. 5811–5824.
- [16] P. DEGOND, A. FROUVILLE, AND G. RAOUL, *Local stability of perfect alignment for a spatially homogeneous kinetic model*, J. Stat. Phys., 157 (2014), pp. 84–112.
- [17] R. J. DIPERNA AND P.-L. LIONS, *Ordinary differential equations, transport theory and Sobolev spaces*, Invent. Math., 98 (1989), pp. 511–547.
- [18] R. L. DOBRUŠIN, *Vlasov equations*, Funktsional. Anal. i Prilozhen., 13 (1979), pp. 48–58, 96.
- [19] M. DOUMIC, A. OLIVIER, AND L. ROBERT, *Estimating the division rate from indirect measurements of single cells*, Discrete Contin. Dyn. Syst. Ser. B, 25 (2020), pp. 3931–3961.
- [20] W. FELLER, *On the integral equation of renewal theory*, Ann. Math. Statistics, 12 (1941), pp. 243–267.
- [21] L. FERRETTI, C. WYMANT, M. KENDALL, L. ZHAO, A. NURTAY, L. ABELER-DÖRNER, M. PARKER, D. BONSALE, AND C. FRASER, *Quantifying sars-cov-2 transmission suggests epidemic control with digital contact tracing*, Science, 368 (2020).
- [22] N. FOURNIER AND E. LÖCHERBACH, *On a toy model of interacting neurons*, Ann. Inst. Henri Poincaré Probab. Stat., 52 (2016), pp. 1844–1876.
- [23] N. FOURNIER AND B. PERTHAME, *Transport distances for pdes: the coupling method*, EMS Surv. Math. Sci., 7 (2020), pp. 1–31.
- [24] P. GABRIEL, *Measure solutions to the conservative renewal equation*, in CIMPA School on Mathematical Models in Biology and Medicine, vol. 62 of ESAIM Proc. Surveys, EDP Sci., Les Ulis, 2018, pp. 68–78.
- [25] W. GANGBO, W. LI, S. OSHER, AND M. PUTHAWALA, *Unnormalized optimal transport*, J. Comput. Phys., 399 (2019), pp. 108940, 17.
- [26] F. GOLSE, C. MOUHOT, AND T. PAUL, *On the mean field and classical limits of quantum mechanics*, Comm. Math. Phys., 343 (2016), pp. 165–205.
- [27] M. HAURAY, *Wasserstein distances for vortices approximation of Euler-type equations*, Math. Models Methods Appl. Sci., 19 (2009), pp. 1357–1384.
- [28] P. LAURENÇOT, *Weak compactness techniques and coagulation equations*, in Evolutionary equations with applications in natural sciences, vol. 2126 of Lecture Notes in Math., Springer, Cham, 2015, pp. 199–253.
- [29] C. MARCHIORO AND M. PULVIRENTI, *Mathematical theory of incompressible nonviscous fluids*, vol. 96 of Applied Mathematical Sciences, Springer-Verlag, New York, 1994.
- [30] J. A. J. METZ AND O. DIEKMANN, *The dynamics of physiologically structured populations*, vol. 68 of Lecture Notes in Biomath., Springer, Berlin, 1986.

- [31] P. MICHEL, S. MISCHLER, AND B. PERTHAME, *General relative entropy inequality: an illustration on growth models*, J. Math. Pures Appl. (9), 84 (2005), pp. 1235–1260.
- [32] S. MIRRAHIMI AND G. RAOUL, *Dynamics of sexual populations structured by a space variable and a phenotypical trait*, Theoretical Population Biology, 84 (2013), pp. 87–103.
- [33] S. MISCHLER AND J. SCHER, *Spectral analysis of semigroups and growth-fragmentation equations*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 33 (2016), pp. 849–898.
- [34] S. MISCHLER AND Q. WENG, *Relaxation in time elapsed neuron network models in the weak connectivity regime*, Acta Appl. Math., 157 (2018), pp. 45–74.
- [35] P. MONMARCHÉ, *On  $\mathcal{H}^1$  and entropic convergence for contractive PDMP*, Electron. J. Probab., 20 (2015), pp. Paper No. 128, 30.
- [36] S. NORDMANN, B. PERTHAME, AND C. TAING, *Dynamics of concentration in a population model structured by age and a phenotypical trait*, Acta Appl. Math., 155 (2018), pp. 197–225.
- [37] K. PAKDAMAN, B. PERTHAME, AND D. SALORT, *Dynamics of a structured neuron population*, Nonlinearity, 23 (2010), pp. 55–75.
- [38] B. PERTHAME, *Transport equations in biology*, Frontiers in Mathematics, Birkhäuser Verlag, Basel, 2007.
- [39] F. SANTAMBROGIO, *Optimal transport for applied mathematicians*, vol. 87 of Progress in Nonlinear Differential Equations and their Applications, Birkhäuser/Springer, Cham, 2015.
- [40] S. TAHERI-ARAGHI, S. BRADDE, J. T. SAULS, N. S. HILL, P. A. LEVIN, J. PAULSSON, M. VERGASSOLA, AND S. JUN, *Cell-size control and homeostasis in bacteria*, Current Biology, (2015).
- [41] C. VILLANI, *Topics in optimal transportation*, vol. 58 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, 2003.