

Exact packing measure of the range of ψ -Super Brownian motions

Xan Duhalde¹ · Thomas Duquesne¹

Received: 16 April 2015 / Revised: 2 November 2015 / Published online: 11 December 2015
© Springer-Verlag Berlin Heidelberg 2015

Abstract We consider super processes whose spatial motion is the d -dimensional Brownian motion and whose branching mechanism ψ is critical or subcritical; such processes are called ψ -super Brownian motions. If $d > 2\gamma/(\gamma - 1)$, where $\gamma \in (1, 2]$ is the lower index of ψ at ∞ , then the total range of the ψ -super Brownian motion has an exact packing measure whose gauge function is $g(r) = (\log \log 1/r)/\varphi^{-1}((1/r \log \log 1/r)^2)$, where $\varphi = \psi' \circ \psi^{-1}$. More precisely, we show that the occupation measure of the ψ -super Brownian motion is the g -packing measure restricted to its total range, up to a deterministic multiplicative constant only depending on d and ψ . This generalizes the main result of Duquesne (Ann Probab 37(6):2431–2458, 2009) that treats the quadratic branching case. For a wide class of ψ , the constant $2\gamma/(\gamma - 1)$ is shown to be equal to the packing dimension of the total range.

Keywords Super Brownian motion · General branching mechanism · Lévy snake · Total range · Occupation measure · Exact packing measure

Mathematics Subject Classification Primary 60G57 · 60J80; Secondary 28A78

✉ Thomas Duquesne
thomas.duquesne@upmc.fr
Xan Duhalde
xan.duhalde@gmail.com

¹ Sorbonne Universités, UPMC Université Paris 06, LPMA (UMR 7599), Boîte courrier 188, 4 place Jussieu, 75252 Paris Cedex 05, France

1 Introduction

The main result of this paper provides an exact packing gauge function for the total range of super processes whose spatial motion is the d -dimensional Brownian motion and whose branching mechanism ψ is critical or subcritical. We call such super processes ψ -super Brownian motions (or ψ -SBM, for short). This generalizes the main result of [10] that concerns the Dawson–Watanabe super process corresponding to the quadratic branching mechanism $\psi(\lambda) = \lambda^2$.

Before stating precisely our main results, let us briefly recall previous works related to fine geometric properties of super processes. Most of these results concern the Dawson–Watanabe super process $(Z_t)_{t \geq 0}$. Dawson and Hochberg [6] have proved that a.s. for all $t > 0$, the Hausdorff dimension of the topological support of Z_t is equal to $2 \wedge d$. In [8], Dawson and Perkins prove that in supercritical dimensions $d \geq 3$, the Dawson–Watanabe super process Z_t is a.s. equal to the h_1 -Hausdorff measure restricted to the topological support of Z_t , where $h_1(r) = r^2 \log \log 1/r$ (see also Perkins [34, 35] for a close result holding a.s. for all times t). By use of the Brownian snake, Le Gall and Perkins [31] prove a similar result in the critical dimension $d = 2$ with the gauge function $h_2(r) = r^2 \log 1/r \log \log 1/r$. In [32], Le Gall, Perkins and Taylor have proved that in dimension $d \geq 3$, the topological support of Z_t has no exact packing measure.

Dawson and Hochberg in [6] also proved that the Hausdorff dimension of the total range of the Dawson–Watanabe super process is a.s. equal to $4 \wedge d$. In [7], Dawson, Iscoe and Perkins investigate the fine geometric properties of the total occupation measure $\mathbf{M} = \int_0^\infty Z_t dt$ of the Dawson–Watanabe super process: they prove that in supercritical dimensions $d \geq 5$, \mathbf{M} is a.s. equal to the h_3 -Hausdorff measure restricted on the total range of the super process, where $h_3(r) = r^4 \log \log 1/r$. In [28], Le Gall considers the critical dimension $d = 4$ and he proves a similar result with respect to the gauge function $h_4(r) = r^4 \log 1/r \log \log 1/r$. In [10], the occupation measure \mathbf{M} is also shown to coincide (in dimension $d \geq 5$) a.s. with the g_1 -packing measure restricted to the total range of the super process, where $g_1(r) = r^4 (\log \log 1/r)^{-3}$.

For super Brownian motions whose branching mechanism is general, less results are available: in [9], Delmas computes the Hausdorff dimension of the total range of super Brownian motions whose branching mechanism is stable; this result is eventually extended in [14] to general branching mechanism ψ thanks to geometric considerations on ψ -Lévy trees. The ψ -Lévy trees are the actual genealogical structures of the ψ -SBM; they are compact random real trees coded by the height process (introduced by Le Gall and Le Jan [30] and further studied in [13]) and they appear as the scaling limits of Galton–Watson trees; their geometric properties are discussed in [11, 12, 14, 15]. In particular, it is proved in [12] that Lévy trees have an exact packing measure, which is closely related to the main result of our article.

Let us introduce precisely the main results of our paper. We first fix a branching mechanism ψ that is critical or subcritical: namely, $\psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is the Laplace exponent of a spectrally positive Lévy process that is of the following Lévy–Khintchine form:

$$\forall \lambda \in \mathbb{R}_+, \quad \psi(\lambda) = \alpha\lambda + \beta\lambda^2 + \int_{(0, \infty)} (e^{-\lambda r} - 1 + \lambda r) \pi(dr), \quad (1)$$

where $\alpha, \beta \in \mathbb{R}_+$, and π is the Lévy measure that satisfies $\int_{(0,\infty)} (r \wedge r^2) \pi(dr) < \infty$. The branching mechanism ψ is the main parameter that governs the law of the processes that are considered in this paper. We introduce two exponents that compare ψ with power functions at infinity:

$$\gamma = \sup \left\{ c \in \mathbb{R}_+ : \lim_{\lambda \rightarrow \infty} \psi(\lambda)\lambda^{-c} = \infty \right\}, \quad \eta = \inf \left\{ c \in \mathbb{R}_+ : \lim_{\lambda \rightarrow \infty} \psi(\lambda)\lambda^{-c} = 0 \right\}. \tag{2}$$

The lower index γ and the upper index η have been introduced by Blumenthal and Gettoor [4]: they appear in the fractal dimensions and the regularity of the processes that we consider. The statements of the paper also involve a third exponent:

$$\delta = \sup \left\{ c \in \mathbb{R}_+ : \exists C \in (0, \infty) \text{ such that } C\psi(\mu)\mu^{-c} \leq \psi(\lambda)\lambda^{-c}, 1 \leq \mu \leq \lambda \right\} \tag{3}$$

that has been introduced in [12]. It is easy to check that $1 \leq \delta \leq \gamma \leq \eta \leq 2$. If ψ is regularly varying at ∞ , all these exponents coincide. In general, they are however distinct and we mention that there exist branching mechanisms ψ of the form (1) such that $\delta = 1 < \gamma = \eta$ (see Lemma 2.3 and Lemma 2.4 in [12] for more details). In our paper we shall often assume that $\delta > 1$ which is a mild regularity assumption on ψ (see Comment 1.4 below).

The space \mathbb{R}^d stands for the usual d -dimensional Euclidian space. We denote by $M_f(\mathbb{R}^d)$ the space of finite Borel measures equipped with the topology of weak convergence. For all $\mu \in M_f(\mathbb{R}^d)$ and for all Borel measurable functions $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$, we use the following notation:

$$\langle f, \mu \rangle = \int_{\mathbb{R}^d} f(x) \mu(dx) \quad \text{and} \quad \langle \mu \rangle = \mu(\mathbb{R}^d).$$

Then, $\langle \mu \rangle$ is the total mass of μ . We shall also denote by $\text{Supp}(\mu)$ the topological support of μ that is the smallest closed subset supporting μ .

Unless the contrary is explicitly mentioned, all the random variables that we consider are defined on the same measurable space (Ω, \mathcal{F}) . We first introduce a \mathbb{R}^d -valued continuous process $\xi = (\xi_t)_{t \geq 0}$; for all $x \in \mathbb{R}^d$, we let \mathbf{P}_x be a probability measure on (Ω, \mathcal{F}) such that ξ under \mathbf{P}_x is distributed as a standard d -dimensional Brownian motion starting from x . We also introduce $Z = (Z_t)_{t \in \mathbb{R}_+}$ that is a $M_f(\mathbb{R}^d)$ -valued càglàd process defined on (Ω, \mathcal{F}) , and for all $\mu \in M_f(\mathbb{R}^d)$, we let \mathbb{P}_μ be a probability measure on (Ω, \mathcal{F}) such that Z under \mathbb{P}_μ is distributed as a super Brownian motion with branching mechanism ψ (a ψ -SBM for short). Namely, under \mathbb{P}_μ , Z is a Markov process whose transitions are characterized as follows: for all bounded Borel measurable functions $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$ and for all $s, t \in \mathbb{R}_+$,

$$\mathbb{P}_\mu\text{-a.s.} \quad \mathbb{E}_\mu \left[\exp(-\langle Z_{t+s}, f \rangle) \mid Z_s \right] = \exp(-\langle Z_s, v_t \rangle), \tag{4}$$

where the function $(v_t(x))_{t \in \mathbb{R}_+, x \in \mathbb{R}^d}$ is the unique nonnegative solution of the integral equation

$$v_t(x) + \mathbf{E}_x \left[\int_0^t \psi(v_{t-s}(\xi_s)) \, ds \right] = \mathbf{E}_x [f(\xi_t)] \quad , \quad x \in \mathbb{R}^d, \quad t \in [0, \infty).$$

Dawson–Watanabe super processes correspond, up to scaling in time and space, to the branching mechanism $\psi(\lambda) = \lambda^2$. Super diffusions with general branching mechanisms of the form (1) have been introduced by Dynkin [16]; for a detailed account on super processes, we refer to the books of Dynkin [17, 18], Le Gall [29], Perkins [5], Etheridge [20] and Li [33].

We easily check that, under \mathbb{P}_μ , the process $(\langle Z_t \rangle)_{t \geq 0}$ of the total mass of the ψ -SBM is a *continuous states branching process* with branching mechanism ψ . Continuous states branching processes have been introduced by Jirina [24] and Lamperti [26, 27], and further studied by Bingham [3]. The assumption $\delta > 1$, implies $\gamma > 1$, which easily entails $\int^\infty d\lambda/\psi(\lambda) < \infty$ that is called the *Grey condition*. Under this condition, standard results on continuous states branching processes (see Bingham [3]) imply that $\langle Z \rangle$ is absorbed in 0 in finite time: namely, $\mathbb{P}_\mu(\exists t \in \mathbb{R}_+ : Z_t = 0) = 1$. Thus the following definition makes sense:

$$\mathbf{M} = \int_0^\infty Z_t \, dt \tag{5}$$

and \mathbf{M} is therefore a random finite Borel measure on \mathbb{R}^d : it is the *occupation* measure of the ψ -SBM Z . We also define the *total range* of Z by

$$\mathbf{R} = \bigcup_{\varepsilon > 0} \overline{\bigcup_{t \geq \varepsilon} \text{Supp}(Z_t)} \tag{6}$$

where for any subset B in \mathbb{R}^d , \overline{B} stands for its closure. We recall here a result due to Sheu [38] that gives a condition on ψ for \mathbf{R} to be bounded:

$$\mathbb{P}_\mu\text{-a.s. } \mathbf{R} \text{ is bounded} \iff \int_1^\infty \frac{db}{\sqrt{\int_1^b \psi(a) da}} < \infty \tag{7}$$

and $\text{Supp}(\mu)$ is compact.

See also Hesse and Kyprianou [23] for a simple probabilistic proof. Note that if $\gamma > 1$, then (7) holds true.

We next denote by \dim_H and \dim_p respectively the *Hausdorff* and the *packing dimensions* on \mathbb{R}^d ; we also denote by $\underline{\dim}$ and $\overline{\dim}$ the *lower* and the *upper box dimensions*: see Falconer [21] for precise definitions. We recall here [14, Theorem 6.3] that asserts that for all $\mu \in M_f(\mathbb{R}^d)$ distinct from the null measure, the following holds true.

$$\text{If } \gamma > 1, \text{ then } \mathbb{P}_\mu\text{-a.s. } \dim_H(\mathbf{R}) = d \wedge \frac{2\eta}{\eta - 1}. \tag{8}$$

If furthermore $\text{Supp}(\mu)$ is compact, then \mathbf{R} is bounded (by (7)) and [14, Theorem 6.3] also asserts that $\mathbb{P}_\mu\text{-a.s. } \underline{\dim}(\mathbf{R}) = d \wedge \frac{2\eta}{\eta - 1}$. As already mentioned (8) generalizes the work of Delmas [9] that treats SBMs whose branching mechanism is stable. Note that

the assumption $\gamma > 1$ is not completely satisfactory for $\dim_H(\mathbf{R})$ only depends on d and η (see [14, Proposition 5.7] and the discussion in Section 5.3 of this article). The first result of our paper computes the packing dimension of \mathbf{R} under more restrictive assumptions.

Theorem 1.1 *Let $\mu \in M_f(\mathbb{R}^d)$ be distinct from the null measure. Let ψ be of the form (1). Let \mathbf{R} be the total range of the ψ -SBM with initial measure μ , as defined in (6). Assume that $\delta > 1$ and that $d > \frac{2\delta}{\delta-1}$. Then,*

$$\mathbb{P}_\mu\text{-a.s. } \dim_p(\mathbf{R}) = \frac{2\gamma}{\gamma - 1}. \tag{9}$$

If furthermore $\text{Supp}(\mu)$ is compact, then $\mathbb{P}_\mu\text{-a.s. } \overline{\dim}(\mathbf{R}) = \frac{2\gamma}{\gamma-1}$.

Comment 1.1 [14, Theorem 5.5] shows that if $\gamma > 1$, the Hausdorff and the packing dimensions of the ψ -Lévy tree are resp. $\frac{\eta}{\eta-1}$ and $\frac{\gamma}{\gamma-1}$. Thus, for all sufficiently large d , (8) and (9) imply that the Hausdorff and the packing dimensions of \mathbf{R} are *twice* that of the ψ -Lévy tree. This can be informally explained by the fact that \mathbf{R} appears as the range of the ψ -Lévy snake that can be viewed as a Gaussian process indexed by the ψ -Lévy tree that is $(\frac{1}{2} - \varepsilon)$ -Hölder regular for any $\varepsilon \in (0, 1/2)$ (see [14, Lemma 6.4] and see also (35) in Sect. 2.4 of the present paper for more details). \square

Comment 1.2 The above mentioned Hölder-regularity of the Lévy snake studied in [14] also entails that if $\text{Supp}(\mu)$ is compact and if $\gamma > 1$, then $\overline{\dim}(\mathbf{R}) \leq d \wedge \frac{2\gamma}{\gamma-1}$. Since $\dim_p(A) \leq \overline{\dim}(A)$ for any bounded $A \subset \mathbb{R}^d$ (see e.g. [21]), an easy argument entails that for all non-null finite measure μ , if $\gamma > 1$, then $\mathbb{P}_\mu\text{-a.s. } \dim_p(\mathbf{R}) \leq d \wedge \frac{2\gamma}{\gamma-1}$. This, combined with (8) entails that

$$\text{if } \eta = \gamma > 1, \text{ then } \mathbb{P}_\mu\text{-a.s. } \dim_H(\mathbf{R}) = \dim_p(\mathbf{R}) = d \wedge \frac{2\gamma}{\gamma - 1}, \tag{10}$$

with the same equality for the lower and the upper box dimensions if $\text{Supp}(\mu)$ is compact. The equality $\gamma = \eta$ holds true for instance if ψ is regularly varying at ∞ . Therefore, the novelty of Theorem 1.1 only concerns the cases where $\gamma \neq \eta$. \square

Comment 1.3 In Theorem 1.1, note that the assumption $\delta > 1$ is not optimal since the value of $\dim_p(\mathbf{R})$ only depends on γ . Actually, the assumptions $\delta > 1$ and $d > \frac{2\delta}{\delta-1}$ are used in Lemma 3.10 and in the proof of (134) that entails the lower bound $\dim_p(\mathbf{R}) \geq \frac{2\gamma}{\gamma-1}$: in this proof, we compare the occupation measure of balls with specific powers of their radius, and if ψ is irregular, then these arguments fail to prove (9) when $d \in (\frac{2\gamma}{\gamma-1}, \frac{2\delta}{\delta-1})$. We conjecture that *if $\gamma > 1$, then $\mathbb{P}_\mu\text{-a.s. } \dim_p(\mathbf{R}) = d \wedge \frac{2\gamma}{\gamma-1}$.* \square

Let us set $\varphi = \psi' \circ \psi^{-1}$. The main properties of that increasing function are stated in Sect. 2.2. Here, we just notice that the inverse function of φ , that is denoted by φ^{-1} ,

is defined from $[\alpha, \infty)$ to $[0, \infty)$. Then, we set

$$g(r) = \frac{\log \log \frac{1}{r}}{\varphi^{-1} \left(\left(\frac{1}{r} \log \log \frac{1}{r} \right)^2 \right)}, \quad r \in (0, r_0) \tag{11}$$

where $r_0 = \min(\alpha^{-1/2}, e^{-e})$, with the convention $\alpha^{-1/2} = \infty$ if $\alpha = 0$. We check (see Sect. 2.2) that g is a continuous increasing function such that $\lim_{0+} g = 0$.

We next denote by \mathcal{P}_g the g -packing measure on \mathbb{R}^d , whose definition is recalled in Sect. 2.1. The following theorem is the main result of the paper.

Theorem 1.2 *Let $\mu \in M_f(\mathbb{R}^d)$ be distinct from the null measure. Let ψ be of the form (1). Let Z be a ψ -SBM starting from μ ; let \mathbf{R} be its total range, as defined by (6), and let \mathbf{M} be its occupation measure, as defined by (5). Let g be defined by (11). Assume that*

$$\delta > 1 \quad \text{and} \quad d > \frac{2\gamma}{\gamma - 1}.$$

Then, there exists a positive constant $\kappa_{d,\psi}$ that only depends on d and ψ such that

$$\mathbb{P}_\mu\text{-a.s.} \quad \mathbf{M} = \kappa_{d,\psi} \mathcal{P}_g(\cdot \cap \mathbf{R}). \tag{12}$$

Comment 1.4 Unlike Theorem 1.1, we think that the assumptions of Theorem 1.2 are optimal: indeed, since $d \wedge \frac{2\gamma}{\gamma-1}$ is thought to be the packing dimension of \mathbf{R} , (12) probably does not hold true when $d \leq \frac{2\gamma}{\gamma-1}$. Moreover, arguing as in [12, Lemma 2.3], we easily check that $\delta > 1$ if and only if g satisfies a *doubling condition* (see (13), Sect. 2.1). Although it is possible to define packing measures with respect to irregular gauge functions (see Edgar [19]), the doubling condition is the minimal assumption on a gauge function which implies that the corresponding packing measure has nice properties (regularity, comparison lemmas). In this sense, Assumption $\delta > 1$ is required to stay within the framework of standard geometric measures. \square

Comment 1.5 In the stable cases where $\psi(\lambda) = \lambda^\gamma$ with $\gamma \in (1, 2]$, then

$$\forall r \in (0, \infty), \quad g(r) = r^{\frac{2\gamma}{\gamma-1}} (\log \log 1/r)^{-\frac{\gamma+1}{\gamma-1}}.$$

If $\gamma = 2$, then $g(r) = r^4 (\log \log 1/r)^{-3}$ and we recover the result from [10]. Moreover, note that in the stable cases, Theorem 1.1 is a straightforward consequence of Theorem 1.2. However, when $\gamma \neq \eta$, it turns out that Theorem 1.1 cannot be simply derived from Theorem 1.2: indeed, one important ingredient of the proof of Theorem 1.2 consists in computing the lower local density of \mathbf{M} with respect to g ; this lower limit is achieved along a deterministic sequence of radii $(\rho_n)_{n \in \mathbb{N}}$ defined in Lemma 3.7 (and Lemma 3.6) and it seems difficult to compare $g(\rho_n)$ with a power of ρ_n . \square

The paper is organised as follows. In Sect. 2, we recall definitions and basic results: Sect. 2.1 is devoted to packing measures and to two comparison lemmas that are standard technical tools; Sect. 2.2 gathers elementary facts on the power exponents δ , γ and

η associated with the branching mechanism ψ ; in Sect. 2.3 and in Sect. 2.4, we recall the definitions of—and various results on—the ψ -height process, the corresponding ψ -Lévy tree and the associated ψ -Lévy snake. In Sect. 3, we prove estimates on a specific subordinator (Sects. 3.1 and 3.3) and on functionals of the snake involving the hitting time of a given ball (Sects. 3.2 and 3.4). Section 4 is devoted to the proof of the two main theorems: we prove Theorem 1.2 first and Theorem 1.1 next.

2 Notations, definitions and preliminary results

2.1 Packing measures

In this section, we briefly recall basic results on packing measures on the Euclidian space \mathbb{R}^d that have been introduced by Taylor and Tricot in [39].

A *gauge function* is an increasing continuous function $g : (0, r_0) \rightarrow (0, \infty)$, where $r_0 \in (0, \infty)$, such that $\lim_{0+} g = 0$ and that satisfies a *doubling condition*: namely, there exists $C \in (0, \infty)$ such that

$$\forall r \in (0, r_0/2), \quad g(2r) \leq Cg(r). \tag{13}$$

Let $B \subset \mathbb{R}^d$ and let $\varepsilon \in (0, \infty)$. Recall that a (closed) ε -packing of B is a finite collection of pairwise disjoint closed ball $(\overline{B}(x_m, r_m))_{1 \leq m \leq n}$ whose centers x_m belong to B and whose radii r_m are not greater than ε . We then set

$$\mathcal{P}_g^{(\varepsilon)}(B) = \sup \left\{ \sum_{1 \leq m \leq n} g(r_m) ; (\overline{B}(x_m, r_m))_{1 \leq m \leq n}, \varepsilon\text{-packing of } B \right\} \tag{14}$$

and

$$\mathcal{P}_g^*(B) = \lim_{\varepsilon \rightarrow 0+} \mathcal{P}_g^{(\varepsilon)}(B) \in [0, \infty] \tag{15}$$

that is called the *g-packing pre-measure of B*. The *g-packing measure of B* is then given by

$$\mathcal{P}_g(B) = \inf \left\{ \sum_{n \geq 0} \mathcal{P}_g^*(B_n) ; B \subset \bigcup_{n \geq 0} B_n \right\}. \tag{16}$$

Remark 2.1 The definition (14) of $\mathcal{P}_g^{(\varepsilon)}$ that we adopt here is slightly different from the one introduced by Taylor and Tricot [39] who take the infimum of $\sum_{m=1}^n g(2r_m)$ over ε -packings with open balls. However, since g is increasing, continuous and since it satisfies a doubling condition (13), the resulting measure is quite close to Taylor and Tricot’s definition: the packing measures are equivalent (bounded above and below by constants) and their main results on packing measures immediately apply to the g -packing measures defined by (14), (15) and (16). Consequently, (12) in Theorem 1.2 also holds true with Taylor and Tricot’s definition of g -packing measures, only the constant $\kappa_{d,\psi}$ may differ. □

Next recall from [39, Lemma 5.1] that \mathcal{P}_g is a Borel-regular outer measure. Moreover, it is obvious from the definition that for any subset $B \subset \mathbb{R}^d$,

$$\mathcal{P}_g(B) \leq \mathcal{P}_g^*(B). \tag{17}$$

Next, if B is a \mathcal{P}_g -measurable set such that $\mathcal{P}_g(B) < \infty$, then for any $\varepsilon > 0$, there exists a closed subset $F_\varepsilon \subset B$ such that

$$\mathcal{P}_g(B) \leq \mathcal{P}_g(F_\varepsilon) + \varepsilon. \tag{18}$$

We recall here [39, Theorem 5.4] that is a standard comparison result for packing measures.

Theorem 2.1 [39, Theorem 5.4] *Let μ be a finite Borel measure on \mathbb{R}^d . Let B be a Borel subset of \mathbb{R}^d . Let g be a gauge function satisfying a doubling condition (13) with a constant $C > 0$. Then, the following holds true.*

- (i) *If $\liminf_{r \rightarrow 0} \frac{\mu(B(x,r))}{g(r)} > 1$ for any $x \in B$, then $\mu(B) \geq \mathcal{P}_g(B)$.*
- (ii) *If $\liminf_{r \rightarrow 0} \frac{\mu(B(x,r))}{g(r)} < 1$ for any $x \in B$, then $\mu(B) \leq C^2 \mathcal{P}_g(B)$.*

We also recall the following specific result due to Edgar in [19, Corollary 5.10].

Lemma 2.2 [19, Corollary 5.10] *Let μ be a finite Borel measure on \mathbb{R}^d . Let $\kappa \in (0, \infty)$ and let B be a Borel subset of \mathbb{R}^d such that*

$$\forall x \in B, \quad \liminf_{r \rightarrow 0^+} \frac{\mu(B(x,r))}{g(r)} = \kappa.$$

Then, $\mu(B) = \kappa \mathcal{P}_g(B)$.

Remark 2.2 The main purpose of Edgar’s article [19] is to deal with fractal measures in metric spaces with respect to possibly irregular gauge functions. Corollary 5.10 [19] (stated here as Lemma 2.2) holds true in this general setting if μ satisfies the *Strong Vitali Property* (see [19, p. 43] for a definition and a discussion of this topic). A result due to Besicovitch [2] ensures that any finite measure on \mathbb{R}^d enjoys the Strong Vitali Property. Therefore, Lemma 2.2 is an immediate consequence of Edgar’s Corollary 5.10 [19]. □

We finally recall the definition of the *packing dimension*: let $\alpha \in (0, \infty)$; we simply write \mathcal{P}_α instead of \mathcal{P}_g when $g(r) = r^\alpha, r \in (0, \infty)$. Let $B \subset \mathbb{R}^d$. Then, the *packing dimension of B* , denoted by $\dim_p(B)$, is the unique real number in $[0, d]$ such that

$$\mathcal{P}_\alpha(B) = \infty \text{ if } \alpha < \dim_p(B) \text{ and } \mathcal{P}_\alpha(B) = 0 \text{ if } \alpha > \dim_p(B). \tag{19}$$

2.2 Exponents

In this section we briefly recall from [12] several results relating power exponents associated with ψ to properties of the gauge function g introduced in (11). We first

recall that the branching mechanism ψ has the Lévy–Khintchine form (1); it is well-known that ψ' is the Laplace exponent of a subordinator, just like ψ^{-1} , the inverse function of ψ . Thus, $\varphi = \psi' \circ \psi^{-1}$ is also the Laplace exponent of a subordinator. Note that $\psi'(0) = \alpha$. As already mentioned, the inverse function of φ , denoted by φ^{-1} , is then defined from $[\alpha, \infty)$ to $[0, \infty)$. We also introduce the function $\tilde{\psi}(\lambda) = \psi(\lambda)/\lambda$ that is the Laplace exponent of a subordinator. Next, observe that $1/\varphi$ is the derivative of ψ^{-1} ; recall that ψ is convex and that $\psi', \tilde{\psi}, \psi^{-1}$ and φ are concave. In particular, it implies $\tilde{\psi}(2\lambda) \leq 2\tilde{\psi}(\lambda)$ and the following inequalities

$$\begin{aligned} \psi(2\lambda) &\leq 4\psi(\lambda), \quad \tilde{\psi}(\lambda) \leq \psi'(\lambda) \leq \frac{\psi(2\lambda) - \psi(\lambda)}{\lambda} \leq 4\tilde{\psi}(\lambda) \quad \text{and} \quad \frac{\lambda}{\psi^{-1}(\lambda)} \leq \varphi(\lambda) \\ &\leq \frac{4\lambda}{\psi^{-1}(\lambda)}. \end{aligned} \tag{20}$$

Let $\phi : [0, \infty) \rightarrow [0, \infty)$ be a continuous increasing function. We agree on $\sup \phi = 0$ and $\inf \phi = \infty$ and we define the following exponents that compare ϕ with power functions at infinity:

$$\gamma_\phi = \sup \left\{ c \in \mathbb{R}_+ : \lim_{\lambda \rightarrow \infty} \phi(\lambda)\lambda^{-c} = \infty \right\}, \quad \eta_\phi = \inf \left\{ c \in \mathbb{R}_+ : \lim_{\lambda \rightarrow \infty} \phi(\lambda)\lambda^{-c} = 0 \right\}. \tag{21}$$

Then, γ_ϕ (resp. η_ϕ) is the lower exponent (resp. the upper exponent) of ϕ at ∞ . We also introduce the following exponent

$$\delta_\phi = \sup \left\{ c \in \mathbb{R}_+ : \exists C \in (0, \infty) \text{ such that } C\phi(\mu)\mu^{-c} \leq \phi(\lambda)\lambda^{-c}, \quad 1 \leq \mu \leq \lambda \right\} \tag{22}$$

that plays a important role for the regularity of the gauge function. By (2) and (3)

$$\boldsymbol{\gamma} = \gamma_\psi, \quad \boldsymbol{\eta} = \eta_\psi, \quad \boldsymbol{\delta} = \delta_\psi.$$

It is easy to check that $1 \leq \boldsymbol{\delta} \leq \boldsymbol{\gamma} \leq \boldsymbol{\eta} \leq 2$. If ψ is regularly varying at ∞ , all these exponents coincide. In general, they are however distinct and we mention that there exist branching mechanisms ψ of the form (1) such that $\boldsymbol{\delta} = 1 < \boldsymbol{\gamma} = \boldsymbol{\eta}$: see Lemma 2.3 and Lemma 2.4 in [12] for more detail. As a direct consequence of (20) we have $\delta_{\tilde{\psi}} = \delta_{\psi'} = \boldsymbol{\delta} - 1$, $\gamma_{\tilde{\psi}} = \gamma_{\psi'} = \boldsymbol{\gamma} - 1$ and $\eta_{\tilde{\psi}} = \eta_{\psi'} = \boldsymbol{\eta} - 1$. Moreover, we get $\delta_\varphi = (\boldsymbol{\delta} - 1)/\boldsymbol{\delta}$, $\gamma_\varphi = (\boldsymbol{\gamma} - 1)/\boldsymbol{\gamma}$ and $\eta_\varphi = (\boldsymbol{\eta} - 1)/\boldsymbol{\eta}$.

Recall from (11) the definition of the function g . The arguments of the proof of Lemma 2.3 (i) in [12] can be immediately adapted to prove that $g : (0, r_0) \rightarrow (0, \infty)$ is an increasing continuous function such that $\lim_{0+} g = 0$ and such that it satisfies the following.

- (i) The function g satisfies the doubling condition (13) iff $\boldsymbol{\delta} > 1$.
- (ii) If ψ is regularly varying at ∞ with exponent $c > 1$, then $\boldsymbol{\delta} = \boldsymbol{\gamma} = \boldsymbol{\eta} = c$ and g is regularly varying at ∞ with exponent $c/(c - 1)$.

We shall further need the following bound that is a consequence of (20).

Lemma 2.3 *Let g the gauge function defined by (11). Let $c \in (0, \infty)$. Then there exists $r(c) \in (0, r_0)$ that only depends on c such that*

$$\forall r \in (0, r(c)), \quad g(r) \psi'^{-1}(c/r^2) \leq 4r^2.$$

Proof Take $r(c) \in (0, r_0)$ such that $\log \log 1/r(c) \geq 1 \vee \sqrt{c}$. Thus,

$$r \in (0, r(c)), \quad \psi'^{-1}(cr^{-2}) \leq \psi'^{-1}((r^{-1} \log \log r^{-1})^2).$$

Recall that $\varphi^{-1} = \psi \circ \psi'^{-1}$. By comparing $\tilde{\psi}$ and ψ' thanks to (20), we get for all $r \in (0, r(c))$

$$\begin{aligned} g(r) \psi'^{-1}(cr^{-2}) &= \frac{\psi'^{-1}(cr^{-2}) \log \log \frac{1}{r}}{\psi(\psi'^{-1}((\frac{1}{r} \log \log \frac{1}{r})^2))} \leq \frac{\psi'^{-1}((\frac{1}{r} \log \log \frac{1}{r})^2) \log \log \frac{1}{r}}{\psi(\psi'^{-1}((\frac{1}{r} \log \log \frac{1}{r})^2))} \\ &\leq \frac{\log \log \frac{1}{r}}{\tilde{\psi}(\psi'^{-1}((\frac{1}{r} \log \log \frac{1}{r})^2))} \leq \frac{4r^2}{\log \log \frac{1}{r}} \leq 4r^2, \end{aligned}$$

which is the desired result. □

2.3 Height processes and Lévy trees

In this section we recall the definition of height processes, that encode Lévy trees. The Lévy trees are the scaling limit of Galton–Watson trees and they are the genealogy of super-Brownian motion.

The height process. Recall that ψ stands for a branching mechanism of the form (1). We always assume that $\gamma > 1$. It is convenient to work on the canonical space $\mathbb{D}([0, \infty), \mathbb{R})$ of càdlàg paths equipped with Skorohod topology and the corresponding Borel sigma-field. We denote by $X = (X_t)_{t \geq 0}$ the canonical process and by \mathbb{P} the distribution of the spectrally positive Lévy processes starting from 0 and whose Laplace exponent is ψ . Namely,

$$\forall t, \lambda \in [0, \infty), \quad \mathbb{E}[\exp(-\lambda X_t)] = \exp(t\psi(\lambda)).$$

Note that the specific form (1) of ψ implies that X_t is integrable and that $\mathbb{E}[X_t] = -\alpha t$. This easily entails that X does not drift to ∞ . Conversely, if a spectrally positive Lévy process does not drift to ∞ , then its Laplace exponent is necessarily of the form (1). We shall assume that $\gamma > 1$ which easily implies that either $\beta > 0$ (and $\gamma = 2$) or $\int_{(0,1)} r\pi(dr) = \infty$. It entails that \mathbb{P} -a.s. X has unbounded variation paths: see Bertoin [1, Chapter VII, Corollary 5 (iii)].

Note that $\gamma > 1$ entails $\int_0^\infty d\lambda/\psi(\lambda) < \infty$ and, as shown by Le Gall and Le Jan [30] (see also [13, Chapter 1]), there exists a continuous process $H = (H_t)_{t \geq 0}$ such that for any $t \in [0, \infty)$, the following limit holds true in \mathbb{P} -probability:

$$H_t = \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \int_0^t \mathbf{1}_{\{I_t^s < X_s < I_t^{s+\varepsilon}\}} \, ds, \tag{23}$$

where I_t^s stands for $\inf_{s \leq r \leq t} X_r$. The process $H = (H_t)_{t \geq 0}$ is called the ψ -height process; it turns out to encode the genealogy of super-Brownian motion with branching mechanism ψ as explained below. We shall need the following result that is proved in [13].

Lemma 2.4 [13, Theorem 1.4.4] *Assume that $\gamma > 1$. Then for every $c \in (0, \frac{\gamma-1}{\gamma})$, H is \mathbb{P} -a.s. locally c -Hölder continuous.*

Excursions of the height process. We denote by I , the infimum process of X :

$$\forall t \in \mathbb{R}_+, \quad I_t = \inf_{0 \leq r \leq t} X_r.$$

When ψ is of the form $\psi(\lambda) = \beta\lambda^2$, X is distributed as a Brownian motion and (23) easily implies that H is proportional to $X - I$, which is distributed as a reflected Brownian motion. In the general cases, H is neither a Markov process nor a martingale. However it is possible to develop an excursion theory for H as follows.

Since X has unbounded variation sample paths, basic results of fluctuation theory (see Bertoin [1, Sections VI.1 and VII.1]) entail that $X - I$ is a strong Markov process in $[0, \infty)$ and that 0 is regular for $(0, \infty)$ and recurrent with respect to this Markov process. Moreover, $-I$ is a local time at 0 for $X - I$ (see Bertoin [1, Theorem VII.1]). We denote by N the corresponding excursion measure of $X - I$ above 0. More precisely, we denote by (a_j, b_j) , $j \in \mathcal{I}$, the excursion intervals of $X - I$ above 0 and we define the corresponding excursions by $X^j = X_{(a_j+\cdot) \wedge b_j} - I_{a_j}$, $j \in \mathcal{I}$. Then, $\sum_{j \in \mathcal{I}} \delta_{(-I_{a_j}, X^j)}$ is a Poisson point measure on $[0, \infty) \times \mathbb{D}([0, \infty), \mathbb{R})$ with intensity $dx \, N(dx)$.

Next observe that under \mathbb{P} , the value of H_t only depends on the excursion of $X - I$ straddling time t and we easily check that

$$\bigcup_{j \in \mathcal{I}} (a_j, b_j) = \{t \geq 0 : H_t > 0\}.$$

This allows to define the height process under N as a certain measurable function $H(X)$ of X . We denote by $C(\mathbb{R}_+, \mathbb{R})$ the space of the continuous functions from $[0, \infty)$ to \mathbb{R} equipped with the topology of the uniform convergence on every compact subsets of $[0, \infty)$; by convenience, we shall slightly abuse notation by denoting by $H = (H_t)_{t \geq 0}$ the *canonical process on $C(\mathbb{R}_+, \mathbb{R})$* and by denoting by $N(dH)$ the “distribution” of the height process $H(X)$ associated with X under the excursion measure $N(dX)$. Then we derive from the previous results the following Poisson decomposition of the height process $H(X)$ associated with X under \mathbb{P} : for any $j \in \mathcal{I}$, set $H^j = H_{(a_j+\cdot) \wedge b_j}$; then, under \mathbb{P} , the point measure

$$\sum_{j \in \mathcal{I}} \delta_{(-I_{a_j}, H^j)} \tag{24}$$

is distributed as a Poisson point measure on $[0, \infty) \times C(\mathbb{R}_+, \mathbb{R})$ with intensity $dx \, N(dH)$. For more details, we refer to [13, Chapter 1].

We denote by σ the *duration* of X under its excursion measure N (with an obvious definition). It is easy to check that H and X under N have the same duration and that the following holds true.

$$N\text{-a.e. } \sigma < \infty, \quad H_0 = H_\sigma = 0 \quad \text{and} \quad H_t > 0 \iff t \in (0, \sigma).$$

Basic results of fluctuation theory (see Bertoin [1, Chapter VII]) also entail:

$$\forall \lambda \in (0, \infty) \quad N \left[1 - e^{-\lambda\sigma} \right] = \psi^{-1}(\lambda). \tag{25}$$

Local times of the height process. We recall from [13, Chapter 1, Section 1.3], the following result: there exists a jointly measurable process $(L_s^a)_{a,s \in [0, \infty)}$ such that \mathbb{P} -a.s. for any $a \in [0, \infty)$, $s \mapsto L_s^a$ is continuous, non-decreasing and such that

$$\forall t, a \geq 0, \quad \lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[\sup_{0 \leq s \leq t} \left| \frac{1}{\varepsilon} \int_0^s dr \mathbf{1}_{\{a-\varepsilon < H_r \leq a\}} - L_s^a \right| \right] = 0. \tag{26}$$

The process $(L_s^a)_{s \in [0, \infty)}$ is called the *a-local time of H*. Recall that I stands for the infimum process of X . One can show (see [14, Lemma 1.3.2]) that for fixed t , $L_t^0 = -I_t$. Moreover, one can observe that the support of the random Stieltjes measure dL^a is contained in the closed set $\{t \geq 0 : H_t = a\}$.

A general version of the *Ray–Knight theorem for H* asserts the following. For any $x \geq 0$, set $T_x = \inf\{t \geq 0 : X_t = -x\}$; then, *the process $(L_{T_x}^a ; a \geq 0)$ is distributed as a continuous-states branching process CSBP with branching mechanism ψ and initial state x* (see Le Gall and Le Jan [30, Theorem 4.2], and also [13, Theorem 1.4.1]).

It is possible to define the local times of H under the excursion measure N as follows. For any $b > 0$, let us set $v(b) = N(\sup_{t \in [0, \sigma]} H_t > b)$. Since H is continuous, the Poisson decomposition (24) implies that $v(b) < \infty$, for any $b > 0$. It is moreover clear that v is non-increasing and that $\lim_{\infty} v = 0$. Then, for every $a \in (0, \infty)$, we define a continuous increasing process $(L_t^a)_{t \in [0, \infty)}$, such that for every $b \in (0, \infty)$ and for any $t \in [0, \infty)$, one has

$$\lim_{\varepsilon \rightarrow 0} N \left[\mathbf{1}_{\{\sup H > b\}} \sup_{0 \leq s \leq t \wedge \sigma} \left| \frac{1}{\varepsilon} \int_0^s dr \mathbf{1}_{\{a-\varepsilon < H_r \leq a\}} - L_s^a \right| \right] = 0. \tag{27}$$

We refer to [13, Section 1.3] for more details. Note that N -a.e. $L_t^a = L_\sigma^a$ for all $t \geq \sigma$. The process $(L_t^a)_{t \in [0, \infty)}$ is the *a-local time of the excursion of the height process*.

Lévy trees. We briefly explain how the height process H under its excursion measure N can be viewed as the contour process of a tree called the Lévy tree. Recall that σ is the duration of H under N . For any $s, t \in [0, \sigma]$, we set

$$d(s, t) = H_t + H_s - 2 \inf_{u \in [s \wedge t, s \vee t]} H_u. \tag{28}$$

The quantity $d(s, t)$ represents the distance between the points corresponding to s and t in the Lévy tree. Indeed d is obviously symmetric in s and t and we easily check that d satisfies the triangle inequality. Two real numbers $s, t \in [0, \sigma]$ correspond to the same point in the Lévy tree iff $d(s, t) = 0$, which is denoted by $s \sim t$. Observe that \sim is an equivalence relation. The Lévy tree is then given by the quotient set

$$\mathcal{T} = [0, \sigma] / \sim .$$

Then, d induces a true metric on \mathcal{T} that we keep denoting d . Denote by $p : [0, \sigma] \rightarrow \mathcal{T}$ the *canonical projection*. Since H is continuous N -a.e., so is p , which implies that (\mathcal{T}, d) is a random compact connected metric space. More specifically, (\mathcal{T}, d) is N -a.e a compact *real-tree*, namely a compact metric space such that any two points are connected by a unique self-avoiding path that is furthermore a geodesic: see [14] for more details on Lévy trees viewed as real-trees.

The *mass measure* of the Lévy tree \mathcal{T} , denoted by \mathbf{m} , is the pushforward measure of the Lebesgue measure ℓ on $[0, \sigma]$ via the canonical projection p . Namely, N -a.e. for all Borel measurable function $f : \mathcal{T} \rightarrow \mathbb{R}_+$,

$$\langle \mathbf{m}, f \rangle = \int_0^\sigma \ell(dt) f(p(t)).$$

One can show that N -a.e. the mass measure is diffuse; obviously its topological support is \mathcal{T} and $\mathbf{m}(\mathcal{T}) = \sigma$. We refer to [14] for more details.

Let $r \in (0, \infty)$ and let $t \in [0, \sigma]$. Let $B(p(t), r)$ denote the open ball in (\mathcal{T}, d) with center $p(t)$ and radius r . Then the mass measure of $B(p(t), r)$ in (\mathcal{T}, d) is given by

$$\mathbf{a}(t, r) := \mathbf{m}(B(p(t), r)) = \int_0^\sigma \ell(ds) \mathbf{1}_{\{d(s,t) \leq r\}}. \tag{29}$$

We shall need the following result on the lower density of \mathbf{m} at typical points that is proved in [12, Theorem 1.2]. To that end, we set

$$\forall r \in (0, \alpha \wedge e^{-e}), \quad k(r) := \frac{\log \log \frac{1}{r}}{\varphi^{-1} \left(\frac{1}{r} \log \log \frac{1}{r} \right)}. \tag{30}$$

Lemma 2.5 [12, Theorem 1.2] *Let ψ be a branching mechanism of the form (1). Let σ be the duration of the height process H under its excursion measure N . Let k be as in (30). Assume that $\delta > 1$. Then, there exists a constant $c_1 \in (0, \infty)$ that only depends on ψ such that*

$$N\text{-a.e. for } \ell\text{-almost all } t \in [0, \sigma] \quad \liminf_{r \rightarrow 0} \frac{\mathbf{a}(t, r)}{k(r)} \geq c_1 ,$$

where $\mathbf{a}(t, r)$ is defined by (29) for all $r \in (0, \infty)$ and for all $t \in [0, \sigma]$.

The exploration process. As already mentioned, the height process is not a Markov process. To explore in a Markovian way the Lévy tree, Le Gall and Le Jan in [30]

introduce a measure valued process $\rho = (\rho_t)_{t \geq 0}$ that is called *the exploration process* and that is defined as follows: denote by $M_f(\mathbb{R}_+)$ the set of finite measures on $[0, \infty)$ equipped with the *total variation distance*. Recall that X under \mathbb{P} is a spectrally positive Lévy process starting from 0 whose Laplace exponent is ψ that satisfies $\gamma > 1$. We denote by \mathcal{F}_t^X the sigma-field generated by $X_{\cdot \wedge t}$ augmented with the \mathbb{P} -negligible events. Recall that for all $s, t \in [0, \infty)$ such that $s \leq t$, I_t^s stands for $\inf_{\mu \in [s, t]} X_\mu$. Then, for all $t \in [0, \infty)$, the following definition makes sense under \mathbb{P} or N :

$$\rho_t(dr) = \beta \mathbf{1}_{[0, H_t]}(r) dr + \sum_{\substack{0 < s \leq t \\ X_{s-} < I_t^s}} (I_t^s - X_{s-}) \delta_{H_s}(dr). \tag{31}$$

Note that the $M_f(\mathbb{R}_+)$ -valued process ρ is $(\mathcal{F}_t^X)_{t \geq 0}$ -adapted. The height process H can be deduced from ρ as follows: for any $\mu \in M_f(\mathbb{R}_+)$, we denote by $\text{Supp}(\mu)$ its topological support. We can show that

$$\mathbb{P}\text{-a.s. (or } N\text{-a.e.) } \forall t \in [0, \infty), \quad \text{Supp}(\rho_t) = [0, H_t].$$

As proved in [13], the exploration process ρ admits a càdlàg modification under \mathbb{P} and N . By [13, Proposition 1.2.3], under N , ρ is a càdlàg strong Markov process with respect to $(\mathcal{F}_{t+}^X)_{t \geq 0}$.

2.4 The Lévy Snake

The ψ -Lévy snake is a generalization of Le Gall’s Brownian snake that greatly facilitates the study of super processes: it provides a Markovian parametrisation of the genealogy and the spatial positions of the underlying continuous population that gives rise to the super process. We recall from [13, Chapter 4], the following definition of the ψ -Lévy snake. To that end, recall that $\xi = (\xi_t)_{t \geq 0}$ is a \mathbb{R}^d -valued continuous process defined on (Ω, \mathcal{F}) and that for any $x \in \mathbb{R}^d$, \mathbf{P}_x is a probability measure on (Ω, \mathcal{F}) such that under \mathbf{P}_x , ξ is distributed as a standard d -dimensional Brownian motion starting from x .

Snake with a deterministic Hölder-regular lifetime process. We denote \mathcal{W} the set of *continuous stopped paths*, namely the set of pairs (w, ζ_w) where $\zeta_w \in [0, \infty)$ and $w : [0, \zeta_w] \rightarrow \mathbb{R}^d$ is a *continuous* function. Here ζ_w is the *lifetime* of w . We shall slightly abuse notation by simply denoting w instead of (w, ζ_w) in the sequel. The set \mathcal{W} is equipped with the metric \mathbf{d} that is defined for all $w, w' \in \mathcal{W}$ by

$$\mathbf{d}(w, w') = |\zeta_w - \zeta_{w'}| + \sup_{t \geq 0} \|w(t \wedge \zeta_w) - w'(t \wedge \zeta_{w'})\|.$$

Here $\|\cdot\|$ stands for the Euclidian norm on \mathbb{R}^d . It can be shown that $(\mathcal{W}, \mathbf{d})$ is a Polish space.

To define the finite dimensional marginal distributions of the snake, we first need to introduce its transition kernels. Let $w \in \mathcal{W}$, let $a \in [0, \zeta_w]$ and let $b \in [a, \infty)$. We plainly define a law $R_{a,b}(w, dw')$ on \mathcal{W} by requiring the following.

- (i) $R_{a,b}(w, dw')$ -a.s. $w'(t) = w(t), \forall t \in [0, a]$.
- (ii) $R_{a,b}(w, dw')$ -a.s. $\zeta_{w'} = b$.
- (iii) The law of $(w'(a+t))_{0 \leq t \leq b-a}$ under $R_{a,b}(w, dw')$ is the law of $(\xi_t)_{0 \leq t \leq b-a}$ under $\mathbf{P}_{w(a)}$.

In particular, $R_{0,b}(w, dw')$ is the law of $(\xi_t)_{0 \leq t \leq b}$ under $\mathbf{P}_{w(0)}$.

We denote by $(W_s)_{s \geq 0}$ the canonical process on the space $\mathcal{W}^{\mathbb{R}_+}$ of the \mathcal{W} -valued functions on \mathbb{R}_+ equipped with the product sigma-field. We next fix $x \in \mathbb{R}^d$. We slightly abuse notation by denoting x the stopped path with null lifetime starting from x (and therefore ending at x). Let $h \in C(\mathbb{R}_+, \mathbb{R}_+)$ such that $h(0) = 0$. We call h the *lifetime process*. For all $s, s' \in \mathbb{R}_+$ such that $s' \geq s$, we use the notation $b_h(s, s') = \inf_{s \leq r \leq s'} h(r)$. By the Kolmogorov Extension Theorem there is a unique probability measure Q_x^h on $\mathcal{W}^{\mathbb{R}_+}$ such that for all $0 = s_0 < s_1 < \dots < s_n$,

$$\begin{aligned}
 &Q_x^h [W_{s_0} \in A_0, \dots, W_{s_n} \in A_n] \\
 &= \mathbf{1}_{A_0}(x) \int_{A_1 \times \dots \times A_n} R_{b_h(s_0, s_1), h(s_1)}(x, dw_1) \dots R_{b_h(s_{n-1}, s_n), h(s_n)}(w_{n-1}, dw_n).
 \end{aligned}
 \tag{32}$$

Note that $(h, x) \mapsto Q_x^h$ is measurable and for all $t \in \mathbb{R}_+$, Q_x^h -a.s. $\zeta_{W_t} = h(t)$.

We next discuss the regularity of the process W under Q_x^h . To that end, we assume the following.

The lifetime process h is locally Hölder continuous with exponent $r \in (0, 1]$.

Fix $p \in (1, \infty)$ and $t_0 \in (0, \infty)$. The last inequality of the proof of Proposition 4.4.1 [13, p. 120], entails that there exists a constant C that only depends on t_0 , on p and on the Hölder constant of h on $[0, t_0]$, such that

$$\forall s, t \in [0, t_0], \quad Q_x^h [\mathbf{d}(W_s, W_t)^p] \leq C |t - s|^{pr/2}.
 \tag{33}$$

If $p > 2/r$, then the Kolmogorov continuity criterion applies and asserts that there exists a continuous modification of the process W . We slightly abuse notation by keeping notation Q_x^h for law on $C(\mathbb{R}_+, \mathcal{W})$ of such a modification; likewise, we also keep denoting by $(W_t)_{t \geq 0}$ the canonical process on $C(\mathbb{R}_+, \mathcal{W})$. We then call Q_x^h the law of the *snake with lifetime process h and starting from x* . Working on $C(\mathbb{R}_+, \mathcal{W})$, we see that Q_x^h -a.s. for all $t \in \mathbb{R}_+$, $\zeta_{W_t} = h(t)$ and we next set

$$\widehat{W}_t = W_t(h(t)).
 \tag{34}$$

The process \widehat{W} is called the *snake's endpoint process* that is Q_x^h -a.s. continuous. From (32), we easily get that under Q_x^h , the endpoint process is a Gaussian process whose covariance is characterized by

$$\forall s, t \in \mathbb{R}_+, \quad Q_x^h \left(\|\widehat{W}_t - \widehat{W}_s\|^2 \right) = h(t) + h(s) - 2 \inf_{s \wedge t \leq u \leq s \vee t} h(u).
 \tag{35}$$

Moreover since (33) holds for any $p > 1$, the Kolmogorov criterion implies that for any $q \in (0, r/2)$, \widehat{W} is Q_x^h -a.s. locally q -Hölder continuous.

The definition of Lévy snake. The Lévy snake is the snake whose lifetime process is the height process H introduced in Sect. 2.3. Recall that we assume that $\gamma > 1$. By Lemma 2.4, H is \mathbb{P} -a.s. (or N -a.e.) Hölder regular and the previous construction of the snake applies. We then define the excursion measure of the ψ -Brownian snake starting from $x \in \mathbb{R}^d$ by

$$\mathbb{N}_x = \int_{C(\mathbb{R}_+, \mathbb{R})} N(H \in dh) Q_x^h. \tag{36}$$

Then H is the lifetime process of W . Namely \mathbb{N}_x -a.e. for all $t \in [0, \sigma]$, $\zeta_{W_t} = H_t$ and thus, $\widehat{W}_t = W_t(H_t)$. Moreover, under \mathbb{N}_x , the conditional law of W given H is Q_x^H : we refer to [13, Chapter 4], for more details. Lemma 2.4 and the results discussed right after (35) entail the following lemma.

Lemma 2.6 *Assume that $\gamma > 1$. Then, for any $q \in (0, \frac{\gamma-1}{2\gamma})$, \widehat{W} is \mathbb{N}_x -a.e. locally q -Hölder.*

We need to introduce notations concerning the snake. We next fix $x \in \mathbb{R}^d$ and we work under the excursion measure \mathbb{N}_x of the ψ -Brownian snake. We first note that the range of the endpoint process \widehat{W} is a connected compact subset of \mathbb{R}^d . We shall use the following notation

$$\mathcal{R}_W = \{\widehat{W}_t; t \in [0, \sigma]\}. \tag{37}$$

Recall that for any $a \in (0, \infty)$, $(L_s^a)_{s \geq 0}$ stands for the local time of H at level a . We then denote by $\mathcal{Z}_a(W)$ the random measure on \mathbb{R}^d defined by

$$\langle \mathcal{Z}_a(W), f \rangle = \int_0^\sigma dL_s^a f(\widehat{W}_s),$$

for any Borel measurable $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$. We also set $\mathcal{Z}_0(W) = 0$, the null measure. Recall that $M_f(\mathbb{R}^d)$ stands for the space of finite Borel measures on \mathbb{R}^d equipped with the topology of weak convergence. We can prove that under \mathbb{N}_x , the $M_f(\mathbb{R}^d)$ -valued process $(\mathcal{Z}_a)_{a \geq 0}$ has a càdlàg modification that is denoted in the same way by convenience. The *occupation measure of the snake* \mathcal{M}_W is then defined by

$$\langle \mathcal{M}_W, f \rangle = \int_0^\infty \langle \mathcal{Z}_a(W), f \rangle da = \int_0^\sigma f(\widehat{W}_s) ds, \tag{38}$$

for any Borel measurable $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$. We then recall [13, Theorem 4.2.1] that connects the ψ -Lévy snake to super Brownian motions.

Theorem 2.7 [13, Theorem 4.2.1] *We keep notation from above. We assume that $\gamma > 1$. Let $\mu \in M_f(\mathbb{R}^d)$ and let*

$$\sum_{i \in \mathcal{I}} \delta_{(x_i, W_i)}$$

be a Poisson point measure on $\mathbb{R}^d \times \mathcal{W}$ with intensity $\mu(dx)\mathbb{N}_x(dW)$. We next set the following:

$$Z_0 = \mu \quad \text{and} \quad Z_a = \sum_{i \in \mathcal{J}} \mathcal{Z}_a(W^i), \quad a \in (0, \infty).$$

Then, the process $(Z_a)_{a \geq 0}$ is a càdlàg ψ -super Brownian motion starting from μ (as defined in Sect. 1). Moreover, if \mathbf{R} and \mathbf{M} are defined in terms of Z by (6) and (5), then,

$$\mathbf{R} \cup \{x_j; j \in \mathcal{J}\} = \bigcup_{j \in \mathcal{J}} \mathcal{R}_{W^j} \quad \text{and} \quad \mathbf{M} = \sum_{j \in \mathcal{J}} \mathcal{M}_{W^j}. \tag{39}$$

The last point (39) is not part of [13, Theorem 4.2.1] but it is an easy consequence of that result. To simplify notation, when there is no risk of confusion we shall simply write

$$\mathcal{Z}_a := \mathcal{Z}_a(W), \quad \mathcal{R} := \mathcal{R}_W \quad \text{and} \quad \mathcal{M} := \mathcal{M}_W.$$

Consequences of the Markov property. As the height process, the ψ -Lévy snake $(W_t)_{t \geq 0}$ defined above is not a Markov process. However $\overline{W} := (\rho_t, W_t)_{t \geq 0}$ is a strong Markov process under \mathbb{N}_x : see [13, Theorem 4.1.2] for more details. Instead of fully discussing the Markov property of \overline{W} , we only state here the various results we need, that are consequences of the strong Markov property.

Denote by $(\mathcal{F}_t^W)_{t \geq 0}$ the filtration generated by $(\overline{W}_t)_{t \geq 0}$. As a consequence of the strong Markov property for \overline{W} (see [13, Theorem 4.1.2]) and a specific decomposition of the snake into excursions above the infimum of its lifetime (see [13, Lemma 4.2.4]), we get the following result that is used in Sect. 3.4.

Lemma 2.8 [13, Theorem 4.1.2 and Lemma 4.2.4] *Let T be a $(\mathcal{F}_{t+}^W)_{t \geq 0}$ -stopping time. Let Y be a nonnegative \mathcal{F}_{T+}^W -measurable random variable. Let $G : \mathbb{R}^d \rightarrow \mathbb{R}_+$ be Borel measurable. Then,*

$$\begin{aligned} & \mathbb{N}_0 \left[\mathbf{1}_{\{0 < T < \infty\}} Y \exp \left(- \int_T^\sigma G(\widehat{W}_s) ds \right) \right] \\ &= \mathbb{N}_0 \left[\mathbf{1}_{\{0 < T < \infty\}} Y \exp \left(- \int_{[0, H_T]} \rho_T(dh) \mathbb{N}_{W_T(h)} \left(1 - e^{-\int_0^\sigma G(\widehat{W}_s) ds} \right) \right) \right]. \end{aligned}$$

We shall apply the strong Markov property of \overline{W} at specific hitting times of the snake. More precisely, let us introduce several notations. Let $r \in (0, \infty)$.

(1) The first hitting time of W in the closed ball $\overline{B}(0, r)$ in \mathbb{R}^d is denoted by

$$\tau_r := \inf \{ t \in \mathbb{R}_+ : \widehat{W}_t \in \overline{B}(0, r) \}, \tag{40}$$

with the convention $\inf \emptyset = \infty$. We also introduce the following function

$$\forall x \in \overline{B}(0, r)^c, \quad u_r(x) := \mathbb{N}_x(\tau_r < \infty). \tag{41}$$

Since $t \mapsto \widehat{W}_t$ is continuous, we also get

$$\forall x \in \overline{B}(0, r)^c, \quad u_r(x) = \mathbb{N}_x(\mathcal{R} \cap \overline{B}(0, r) \neq \emptyset). \tag{42}$$

(2) From [13, p. 121 and p. 131], we know that $u_r(x) \in (0, \infty)$, for all $r \in (0, \infty)$ and all $x \in \overline{B}(0, r)^c$, and that u_r is moreover radial. We then denote by \tilde{u}_r the function from (r, ∞) to $(0, \infty)$ such that

$$\forall x \in \overline{B}(0, r)^c, \quad \tilde{u}_r(\|x\|) = u_r(x).$$

(3) Let $r' \in (0, \infty)$. For all stopped path $w \in \mathcal{W}$, we next set

$$T_{r'}(w) = \inf \{s \in [0, \zeta_w] : w(s) \in \overline{B}(0, r')\}, \tag{43}$$

with the convention $\inf \emptyset = \infty$.

(4) We then define a function $\varpi : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$ by

$$\forall \lambda_1, \lambda_2 \in \mathbb{R}_+, \quad \varpi(\lambda_1, \lambda_2) = \begin{cases} (\psi(\lambda_1) - \psi(\lambda_2)) / (\lambda_1 - \lambda_2) & \text{if } \lambda_1 \neq \lambda_2, \\ \psi'(\lambda_1) & \text{if } \lambda_1 = \lambda_2. \end{cases} \tag{44}$$

Recall that $\xi = (\xi_t)_{t \geq 0}$ is a \mathbb{R}^d -valued continuous process defined on (Ω, \mathcal{F}) and that for any $x \in \mathbb{R}^d$, \mathbf{P}_x is a probability measure on (Ω, \mathcal{F}) such that under \mathbf{P}_x , ξ is distributed as a standard d -dimensional Brownian motion starting from x . The following lemma is a specific application of [13, Theorem 4.6.2] that we use in the proof of Lemma 3.15 in Sect. 3.4.

Lemma 2.9 [13, Theorem 4.6.2] *Let $x \in \mathbb{R}^d$. Let $r, r' \in (0, \infty)$ be such that $r' > r$ and $x \in \overline{B}(0, r')^c$. We keep the previous notation. Let $F, G : \mathcal{W} \rightarrow \mathbb{R}_+$ be Borel-measurable. Then,*

$$\begin{aligned} & \mathbb{N}_x \left[\mathbf{1}_{\{\tau_r < \infty\}} F((W_{\tau_r}(s))_{0 \leq s \leq T_{r'}(W_{\tau_r})}) \right. \\ & \quad \times \exp \left(- \int_{[0, T_{r'}(W_{\tau_r})]} \rho_{\tau_r}(dh) G((W_{\tau_r}(s))_{0 \leq s \leq h}) \right) \Big] \\ & = \tilde{u}_r(r') \mathbf{E}_x \left[\mathbf{1}_{\{T_{r'}(\xi) < \infty\}} F((\xi_s)_{0 \leq s \leq T_{r'}(\xi)}) \right. \\ & \quad \times \exp \left(- \int_{[0, T_{r'}(\xi)]} dh \varpi(u_r(\xi_h), G((\xi_s)_{0 \leq s \leq h})) \right) \Big]. \end{aligned}$$

Palm formula. We next recall a decomposition of the occupation measure of the snake according to a point that is randomly chosen. We refer to this decomposition as to the Palm formula. We introduce the following notation

$$\forall \lambda \in \mathbb{R}_+, \quad \psi^*(\lambda) = \psi(\lambda) - \alpha\lambda, \tag{45}$$

that is clearly the Laplace exponent of a spectrally Lévy process. We then fix $x \in \mathbb{R}^d$. Let $U = (U_a)_{a \geq 0}$ be a subordinator defined on $(\Omega, \mathcal{F}, \mathbf{P}_x)$ that is assumed to be independent of ξ and whose Laplace exponent is

$$\forall \lambda \in \mathbb{R}_+, \quad \tilde{\psi}^*(\lambda) := \frac{\psi(\lambda)}{\lambda} - \alpha.$$

For any $a \in \mathbb{R}_+$, we denote by $R_a(db)$ the random measure $\mathbf{1}_{[0,a]}(b)dU_b$. We first recall from [13, see (106), p. 105], that for any measurable function $F : M_f(\mathbb{R}_+) \times \mathcal{W} \rightarrow \mathbb{R}_+$, the following holds true:

$$\mathbb{N}_x \left(\int_0^\sigma ds F(\rho_s, W_s) \right) = \int_0^\infty da e^{-\alpha a} \mathbf{E}_x [F(R_a, (\xi_s)_{0 \leq s \leq a})]. \tag{46}$$

We next provide a Palm decomposition for the occupation measure \mathcal{M} of the snake that is used to estimate its lower local density at “typical” points. To that end, we need to introduce the following auxiliary random variables. Let $(V_t)_{t \geq 0}$ be a subordinator defined on $(\Omega, \mathcal{F}, \mathbf{P}_0)$ that is independent of ξ and whose Laplace exponent is $\psi^{*'}(\lambda) := \psi'(\lambda) - \alpha$. We then introduce the following point measure on $[0, \infty) \times C(\mathbb{R}_+, \mathcal{W})$:

$$\mathcal{N}^*(dt dW) = \sum_{j \in \mathcal{J}^*} \delta_{(t_j, W^j)}, \tag{47}$$

such that under \mathbf{P}_0 and conditionally given (ξ, V) , \mathcal{N}^* is distributed as a Poisson point process with intensity $dV_t \mathbb{N}_{\xi_t}(dW)$.

For all $j \in \mathcal{J}^*$, we denote by \mathcal{M}_j the occupation measure of the snake W^j . Then for all $a \in \mathbb{R}_+$, we define the following random measure on \mathbb{R}^d :

$$\mathcal{M}_a^* = \sum_{j \in \mathcal{J}^*} \mathbf{1}_{[0,a]}(t_j) \mathcal{M}_j. \tag{48}$$

Note that \mathcal{M}_a^* is \mathbf{P}_0 -a.s. a random finite Borel measure on \mathbb{R}^d . Informally \mathcal{M}_a^* is the sum of the occupation measure of the snakes grafted at a rate given by V on the *spatial spine* ξ between time 0 and a . As a by-product of Formula (113) p.113 in [13], we get the following Palm decomposition of \mathcal{M} under \mathbb{N}_0 . Let us mention that in the case of stable branching mechanisms, this is a consequence of the Palm decomposition of historical processes given in Proposition 4.1.5 page 68, in the monograph [8] by Dawson and Perkins.

Proposition 2.10 [13, (113)] *Let $F : \mathbb{R}^d \times M_f(\mathbb{R}^d) \rightarrow \mathbb{R}_+$ be measurable. Then,*

$$\mathbb{N}_0 \left[\int_{\mathbb{R}^d} \mathcal{M}(dy) F(y, \mathcal{M}) \right] = \int_0^\infty da e^{-\alpha a} \mathbf{E}_0 [F(\xi_a, \mathcal{M}_a^*)]. \tag{49}$$

We shall mostly use the Palm formula under the more specific form: for any measurable functional $G : \mathbb{D}(\mathbb{R}_+, \mathbb{R}) \rightarrow \mathbb{R}_+$, we get

$$\mathbb{N}_0 \left[\int_{\mathbb{R}^d} \mathcal{M}(dy) G \left((\mathcal{M}(B(y, r)))_{r \geq 0} \right) \right] = \int_0^\infty da e^{-\alpha a} \mathbf{E}_0 \left[G \left((\mathcal{M}_a^*(B(0, r)))_{r \geq 0} \right) \right]. \tag{50}$$

3 Estimates

3.1 Tail of a subordinator

Recall from (45) that $\psi^*(\lambda) = \psi(\lambda) - \alpha\lambda$; ψ^* is the Laplace transform of a spectrally positive Lévy process. Therefore, $\psi^{*'}$ is the Laplace exponent of a subordinator that is conservative (for $\psi^{*'}(0) = 0$). By subordination, $\sqrt{\psi^{*'} \circ \psi^{-1}}$ is also the Laplace exponent of a conservative subordinator. The main idea of the proof of Theorem 4.1 consists in comparing the occupation measure of a typical ball with a subordinator whose Laplace exponent is $\sqrt{\psi^{*'} \circ \psi^{-1}}$. To that end, we first need the following result.

Lemma 3.1 *Assume that $\delta > 1$. Recall from (11) the definition of the gauge function $g : (0, r_0) \rightarrow (0, \infty)$. Let $(S_r)_{r \in [0, \infty)}$ be a subordinator defined on the auxiliary probability space $(\Omega, \mathcal{F}, \mathbf{P}_0)$. We assume that the Laplace exponent of S is $\sqrt{\psi^{*'} \circ \psi^{-1}}$. Let $\rho_n \in (0, r_0)$, $n \in \mathbb{N}$, be such that*

$$\rho_{n+1} \leq e^{-n} \rho_n \quad \text{and} \quad \sup_{n \geq 0} n^{-2} \log 1/\rho_n < \infty. \tag{51}$$

Then, $\sum_{n \geq 0} \mathbf{P}_0(S_{\rho_n} \leq g(4\rho_n)) = \infty$. Moreover, we get

$$\mathbf{P}_0\text{-a.s.} \quad \limsup_{n \rightarrow \infty} \frac{S_{\rho_{n+1}}}{g(4\rho_n)} \leq 1.$$

Proof To simplify notation, we set $\Phi = \sqrt{\psi^{*'} \circ \psi^{-1}}$. Thus, $\mathbf{E}_0[\exp(-\lambda S_r)] = \exp(-r\Phi(\lambda))$. Denote by Φ^{-1} the inverse function of Φ . For any $r \in (0, e^{-1})$, we set

$$g_*(r) = \frac{\log \log \frac{1}{r}}{\Phi^{-1}\left(\frac{1}{r} \log \log \frac{1}{r}\right)}.$$

An easy computation implies that $\Phi^{-1}(y) = \varphi^{-1}(y^2 + \alpha)$. Since $\alpha \in [0, \infty)$, we easily get $g_*(r) \leq g(r)$, $r \in (0, r_0)$. For any $n \in \mathbb{N}$, we then set $\lambda_n = \Phi^{-1}((4\rho_n)^{-1} \log \log 1/4\rho_n)$. Observe that

$$\lambda_n g(4\rho_n) \geq \lambda_n g_*(4\rho_n) = \log \log(1/4\rho_n).$$

Next note that for all $a, x \in [0, \infty)$, $(1 - e^{-a})\mathbf{1}_{\{0 \leq x \leq a\}} \geq e^{-x} - e^{-a}$, which easily entails

$$\mathbf{P}_0(S_{\rho_n} \leq g(4\rho_n)) \geq \frac{\exp(-\rho_n \Phi(\lambda_n)) - \exp(-\lambda_n g(4\rho_n))}{1 - \exp(-\lambda_n g(4\rho_n))} \underset{n \rightarrow \infty}{\sim} (\log 1/(4\rho_n))^{-\frac{1}{4}}.$$

By the second assumption in (51), $\sum_{n \geq 0} (\log 1/(4\rho_n))^{-\frac{1}{4}} = \infty$, which proves the first point of the lemma.

Let us prove the second point. By a standard Markov inequality, we get

$$\mathbf{P}_0(S_{\rho_{n+1}} \geq g_*(4\rho_n)) \leq \frac{1 - \exp(-\rho_{n+1} \Phi(\lambda_n))}{1 - \exp(-\lambda_n g_*(4\rho_n))} \leq \frac{\rho_{n+1} \Phi(\lambda_n)}{1 - \exp(-\lambda_n g_*(4\rho_n))}.$$

First observe that $1 - \exp(-\lambda_n g_*(4\rho_n)) = 1 - (\log 1/4\rho_n)^{-1} \rightarrow 1$, as $n \rightarrow \infty$. By (51), there exists a constant $c_2 \in (0, \infty)$ such that

$$\rho_{n+1} \Phi(\lambda_n) = \frac{\rho_{n+1}}{4\rho_n} \log \log 1/(4\rho_n) \leq c_2 e^{-n} \log n,$$

Thus, $\sum_{n \geq 0} \mathbf{P}_0(S_{\rho_{n+1}} \geq g(4\rho_n)) \leq \sum_{n \geq 0} \mathbf{P}_0(S_{\rho_{n+1}} \geq g_*(4\rho_n)) < \infty$, which completes the proof by the Borel–Cantelli Lemma. \square

3.2 Estimates on hitting probabilities for the snake

As already mentioned in (7), the total range \mathbf{R} of a ψ -super Brownian motion is bounded if the starting measure μ has compact support and if

$$\int_1^\infty \frac{db}{\sqrt{\int_1^b \psi(a) da}} < \infty. \tag{52}$$

Observe that if $\delta > 1$, then $\gamma > 1$ and (52) holds true, which allows to define the following function

$$\forall v \in (0, \infty), \quad I(v) = \int_v^\infty \frac{db}{\sqrt{\int_v^b \psi(a) da}} = \int_0^\infty \frac{db}{\sqrt{\int_0^b \psi(v+a) da}}. \tag{53}$$

This function is clearly decreasing and continuous and it plays a role in the proof of an upper bound of the hitting probabilities of the ψ -Lévy snake. We first need the following elementary lemma.

Lemma 3.2 *Assume that $\delta > 1$. Then, there exists $c_3 \in (0, \infty)$ that only depends on ψ such that for all $v \in (1, \infty)$ and all $r \in (0, \infty)$ satisfying $r \leq I(v)$, we have*

$$v \leq \psi'^{-1}(4c_3 r^{-2}).$$

Proof By an elementary change of variable, we get

$$I(v) = \int_1^\infty \frac{\sqrt{v} db}{\sqrt{\int_1^b \psi(va) da}}.$$

Fix $c \in (1, \delta)$. By the definition (3) of δ , there exists $C \in (0, \infty)$ such that $\psi(va) \geq C\psi(v)a^c$, for any $a, v \in (1, \infty)$. Let $v \in (1, \infty)$ and $r \in (0, \infty)$ be such that $r \leq I(v)$. Then,

$$r \leq I(v) \leq C^{-1/2}(v/\psi(v))^{1/2} \int_1^\infty \frac{db}{\sqrt{\int_1^b a^c da}} =: (c_3 v/\psi(v))^{1/2},$$

which implies the desired result since $\psi'(v) \leq 4\psi(v)/v$ by (20). □

Recall from (37) the definition of the range \mathcal{R} of the snake. Recall from (36) the notation \mathbb{N}_x for the excursion measure of the snake starting from x . Let $r \in (0, \infty)$. Recall that $B(0, r)$ stands for the open ball in \mathbb{R}^d with radius r and center 0. We set for all $x \in B(0, r)$

$$v_r(x) = \mathbb{N}_x(\mathcal{R} \cap B(0, r)^c \neq \emptyset). \tag{54}$$

From [13, p. 121 and p. 131], we know that

$$\forall x \in B(0, r), \quad v_r(x) \in (0, \infty) \quad \text{and} \quad \lim_{\|x\| \rightarrow r-} v_r(x) = \infty.$$

Moreover, v_r is C^2 on $B(0, r)$ and it satisfies $\frac{1}{2} \Delta v_r = \psi(v_r)$. As an easy consequence of Brownian motion isotropy, v_r is a radial function: namely, $v_r(x)$ only depends on $\|x\|$. Therefore, one can derive estimates on v_r by studying the associated ordinary differential equation corresponding to the radial function, as done by Keller in [25, p. 507 inequality (25)], who proves the following:

$$\forall r \in (0, \infty), \quad \frac{2}{\sqrt{d}}r \leq I(v_r(0)) \leq 2r. \tag{55}$$

We use this bound as follows. For any $r \in (0, \infty)$ and any $x \in B(0, r)^c$, recall that

$$u_r(x) = \mathbb{N}_x(\mathcal{R} \cap \overline{B}(0, r) \neq \emptyset). \tag{56}$$

From [13, p. 121 and p. 131], we know that $u_r(x) \in (0, \infty)$ for all $x \in B(0, r)^c$, that

$$u_r \text{ is radial, } \lim_{\|x\| \rightarrow \infty} u_r(x) = 0 \quad \text{and} \quad \lim_{\|x\| \rightarrow r+} u_r(x) = \infty. \tag{57}$$

Recall that we denote by \tilde{u}_r the radial function yielded by u_r , namely:

$$\forall x \in \overline{B}(0, r)^c, \quad \tilde{u}_r(\|x\|) = u_r(x). \tag{58}$$

Moreover, u_r is C^2 in $B(0, r)^c$ and it satisfies

$$\frac{1}{2} \Delta u_r = \psi(u_r) \quad \text{in } B(0, r)^c. \tag{59}$$

We shall use several times the following upper bound of u_r .

Lemma 3.3 *Assume that $\delta > 1$ and that $d \geq 3$. Let $\varrho \in (0, \infty)$. Then there exist $r_1, c_4(\varrho) \in (0, \infty)$, that only depend on ψ, d and ϱ , such that*

$$\forall r \in (0, r_1), \forall x \in B(0, (1 + \varrho)r)^c, \quad u_r(x) \leq ((1 + \varrho)r/\|x\|)^{d-2} \psi'^{-1}(c_4(\varrho)r^{-2}).$$

Proof Let $y \in \mathbb{R}^d$ be such that $\|y\| = (1 + \varrho)r$. First note that

$$\mathbb{N}_y(\mathcal{R} \cap \overline{B}(0, r) \neq \emptyset) \leq \mathbb{N}_y(\mathcal{R} \cap B(y, \varrho r)^c \neq \emptyset).$$

By translation invariance of Brownian motion, the right member of the previous inequality does not depend on y and we get $u_r(y) \leq v_{\varrho r}(0)$, where $v_{\varrho r}$ is defined by (54). For any $x \in B(0, (1 + \varrho)r)^c$, we next set $w(x) = u_r(x) - v_{\varrho r}(0)((1 + \varrho)r/\|x\|)^{d-2}$ that is clearly subharmonic. The previous upper bound implies that $w(y) \leq 0$, if $\|y\| = (1 + \varrho)r$. By (57), $\lim_{\|x\| \rightarrow \infty} w(x) = 0$ and the maximum principle implies that $w \leq 0$ on $B(0, (1 + \varrho)r)^c$. Namely,

$$\forall r \in (0, \infty), \quad \forall x \in B(0, (1 + \varrho)r)^c, \quad u_r(x) \leq v_{\varrho r}(0) ((1 + \varrho)r/\|x\|)^{d-2}. \tag{60}$$

Since $\delta > 1$, (52) is satisfied and the function I given by (53) is well-defined; we easily check that $I(v) \rightarrow 0$ iff $v \rightarrow \infty$. Then, (55) implies that $\lim_{r \rightarrow 0} v_{\varrho r}(0) = \infty$, so we can find $r_1 \in (0, \infty)$ such that for all $r \in (0, r_1)$, $v_{\varrho r}(0) \geq 1$ and by the left member of (55) we also have $2r\varrho/\sqrt{d} \leq I(v_{\varrho r}(0))$. Thus, Lemma 3.2 applies and asserts that $v_{\varrho r}(0) \leq \psi'^{-1}(c_4(\varrho)r^{-2})$, where $c_4(\varrho) := c_3d\varrho^{-2}$, which completes the proof thanks to (60). \square

We use the previous lemma to get an upper bound of the expectation of a specific additive functional of the Brownian motion that involves u_r . More precisely, for any $r \in (0, \infty)$, we set

$$q_r = \psi'^{-1}(c_4(1)r^{-2}), \tag{61}$$

where $c_4(1)$ is the constant appearing in Lemma 3.3 with $\varrho = 1$. Note that for all $0 < r < (c_4(1)/\psi'(1))^{1/2}$, we have $q_r \geq 1$. We then define

$$\forall r \in \left(0, (c_4(1)/\psi'(1))^{1/2}\right), \quad J(r) = r^2 q_r^{\frac{2}{d-2}} \int_1^{q_r} \psi'(v) v^{-\frac{d}{d-2}} dv. \tag{62}$$

Recall that $\xi = (\xi_t)_{t \geq 0}$ stands for a standard Brownian motion starting from 0 that is defined on $(\Omega, \mathcal{F}, \mathbf{P}_0)$. We next prove the following lemma.

Lemma 3.4 Assume that $\delta > 1$ and that $d \geq 3$. Let $a \in (0, \infty)$. Then, there exist $c_5, c_6, r_2 \in (0, \infty)$ that only depend on ψ, d and a , such that

$$\forall r \in (0, r_2) \quad \mathbf{E}_0 \left[\int_0^{2a} ds \mathbf{1}_{\{\|\xi_s\| \geq 2r\}} \psi'(u_r(\xi_s)) \right] \leq c_5 + c_6 J(r).$$

Proof To simplify notation, we denote by L the expectation in the left member of the previous inequality. Recall from (58) the notation \tilde{u}_r for the radial function yielded by u_r . By Fubini and easy changes of variable, we have the following.

$$\begin{aligned} L &= \int_{B(0,2r)^c} dx \int_0^{2a} ds (2\pi s)^{-d/2} e^{-\|x\|^2/2s} \psi'(\tilde{u}_r(\|x\|)) \\ &= c_7 \int_{2r}^\infty dy y^{d-1} \psi'(\tilde{u}_r(y)) \int_0^{2a} ds s^{-d/2} e^{-y^2/2s} \\ &= c_8 \int_{2r}^\infty dy y f(y) \psi'(\tilde{u}_r(y)), \end{aligned}$$

where for any $y \in (0, \infty)$, we have set $f(y) = \int_{y^2/(4a)}^\infty du u^{d/2-2} e^{-u}$ and where c_7 and c_8 are constants that only depend on d . Since we assume that $d \geq 3$, $f(0)$ is well-defined and finite, and it is easy to check that $\int_0^\infty y f(y) dy < \infty$.

We next use Lemma 3.3 with $\varrho = 1$ to get for all $r \in (0, r_1)$ and all $y \in (2r, \infty)$ that $\tilde{u}_r(y) \leq (2r/y)^{d-2} q_r$. We then set $\alpha_r = 2r q_r^{1/(d-2)}$. Thus, $\tilde{u}_r(y) \leq (\alpha_r/y)^{d-2}$, for all $r \in (0, r_1)$ and all $y \in (2r, \infty)$. We next set

$$r_2 := r_1 \wedge (c_4(1)/\psi'(1))^{1/2}.$$

Observe that for any $r \in (0, r_2)$, $q_r \geq 1$, which implies that $\alpha_r \geq 2r$. Next, observe that for all $r \in (0, r_2)$ and all $y \in (\alpha_r, \infty)$, $(\alpha_r/y)^{d-2} \leq 1$. Thus, $\psi'(\tilde{u}_r(y)) \leq \psi'(1)$. It implies

$$\begin{aligned} L &\leq c_8 \psi'(1) \int_{\alpha_r}^\infty y f(y) dy + c_8 \int_{2r}^{\alpha_r} y f(y) \psi'((\alpha_r/y)^{d-2}) dy \\ &\leq c_5 + c_8 f(0) \int_{2r}^{\alpha_r} y \psi'((\alpha_r/y)^{d-2}) dy, \end{aligned} \tag{63}$$

where $c_5 := c_8 \psi'(1) \int_0^\infty y f(y) dy$. By using the change of variable $v = (\alpha_r/y)^{d-2}$ we get

$$\begin{aligned} &c_8 f(0) \int_{2r}^{\alpha_r} y \psi'((\alpha_r/y)^{d-2}) dy \\ &= \frac{1}{d-2} c_8 f(0) \alpha_r^2 \int_1^{(\alpha_r/2r)^{d-2}} \psi'(v) v^{-\frac{d}{d-2}} dv = c_6 J(r), \end{aligned}$$

where we have set $c_6 := \frac{4}{d-2} c_8 f(0)$. Then, the desired result follows from (63). \square

When d is greater than $\frac{2\delta}{\delta-1}$, the function J is bounded for all small values of r as proved in the following lemma.

Lemma 3.5 *Assume that $\delta > 1$ and that $d > \frac{2\delta}{\delta-1}$. Then, there exists a constant $c_9 \in (0, \infty)$ that depends on d and ψ such that $J(r) \leq c_9$ for all $r \in (0, r_2)$.*

Proof Observe that $\frac{2}{d-2} < \delta - 1$. Recall that $\delta_{\psi'} = \delta - 1$. Let us fix $u \in (\frac{2}{d-2}, \delta - 1)$. By the definition (3) of the exponent $\delta_{\psi'}$, there exists $K \in (0, \infty)$ depending on ψ and u such that $\forall 1 \leq \lambda \leq \mu, \psi'(\lambda) \leq K \psi'(\mu)(\lambda/\mu)^u$. Recall from (61) that $\psi'(q_r) = c_4(1)r^{-2}$, where $c_4(1)$ is the constant appearing in Lemma 3.3, with $\varrho = 1$. Then, we get the following.

$$\begin{aligned} J(r) &= r^2 q_r^{\frac{2}{d-2}} \int_1^{q_r} \psi'(v) v^{-\frac{d}{d-2}} dv \leq r^2 q_r^{\frac{2}{d-2}} \psi'(q_r) \int_1^{q_r} (v/q_r)^u v^{-\frac{d}{d-2}} dv \\ &\leq c_4(1) q_r^{\frac{2}{d-2}-u} \int_1^{q_r} v^{u-\frac{2}{d-2}-1} dv \\ &\leq c_4(1) q_r^{\frac{2}{d-2}-u} \int_0^{q_r} v^{u-\frac{2}{d-2}-1} dv = \frac{c_4(1)}{u-\frac{2}{d-2}}, \end{aligned}$$

which implies the desired result with $c_9 := \frac{c_4(1)}{u-\frac{2}{d-2}}$. □

When $d \in (\frac{2\gamma}{\gamma-1}, \frac{2\delta}{\delta-1}]$, we are only able to prove that $\liminf_{r \rightarrow 0} J(r) < \infty$. More precisely, we prove the following lemma.

Lemma 3.6 *Assume that $\gamma > 1$ and that $d > \frac{2\gamma}{\gamma-1}$. Recall that $c_4(1)$ appears in Lemma 3.3 with $\varrho = 1$. Then, there exists a decreasing function*

$$\theta \in (\psi'(1), \infty) \mapsto r_\theta \in \left(0, (c_4(1)/\psi'(1))^{\frac{1}{2}}\right)$$

such that $\lim_{\theta \rightarrow \infty} r_\theta = 0$ and such that there exists $c_{10} \in (0, \infty)$, that only depends on d and ψ , and that satisfies

$$\forall \theta \in (\psi'(1), \infty), \quad J(r_\theta) \leq c_{10}. \tag{64}$$

Moreover, for any $\theta', \theta \in (\psi'(1), \infty)$ such that $\theta' \geq \theta$, we also have

$$r_{\theta'}/r_\theta \leq (\theta/\theta')^{1/2} \quad \text{and} \quad r_\theta \geq c_{11} \theta^{-c_{12}}, \tag{65}$$

where $c_{11}, c_{12} \in (0, \infty)$ only depend on d and ψ .

Proof Note that $\frac{2}{d-2} < \gamma - 1 = \gamma_{\psi'}$. Let us fix $c \in (\frac{2}{d-2}, \gamma_{\psi'})$. Thus, $\lambda^{-c} \psi'(\lambda) \rightarrow \infty$ as $\lambda \rightarrow \infty$. This allows to define for all $\theta \in (\psi'(1), \infty)$

$$r_\theta = (c_4/\psi'(\lambda_\theta))^{\frac{1}{2}} \quad \text{where} \quad \lambda_\theta = \inf \{ \lambda \in [1, \infty) : \lambda^{-c} \psi'(\lambda) = \theta \}. \tag{66}$$

Note that if $\theta > \psi'(1)$, then $r_\theta < (c_4(1)/\psi'(1))^{\frac{1}{2}}$ and $J(r_\theta)$ is well-defined by (62). Clearly, $\theta \mapsto \lambda_\theta$ increases to ∞ as $\theta \rightarrow \infty$. Consequently $\theta \mapsto r_\theta$ decreases to 0 as $\theta \rightarrow \infty$. Recall from (61) and (62) the definitions of q_r and $J(r)$ and note that $q_{r_\theta} = \lambda_\theta$. By definition, $\psi'(v) \leq \theta v^c$, for any $v \in [1, \lambda_\theta]$, which implies

$$\begin{aligned} J(r_\theta) &= r_\theta^2 \lambda_\theta^{\frac{2}{d-2}} \int_1^{\lambda_\theta} \psi'(v) v^{-\frac{d}{d-2}} dv \leq r_\theta^2 \lambda_\theta^{\frac{2}{d-2}} \theta \int_1^{\lambda_\theta} v^{c-\frac{2}{d-2}-1} dv \\ &\leq \frac{1}{c-\frac{2}{d-2}} r_\theta^2 \lambda_\theta^{\frac{2}{d-2}} \theta \lambda_\theta^{c-\frac{2}{d-2}} = c_{10} \frac{\theta \lambda_\theta^c}{\psi'(\lambda_\theta)} = c_{10}, \end{aligned}$$

where $c_{10} = c_4(1)/(c - \frac{2}{d-2})$ and since $\psi'(\lambda_\theta) = \theta \lambda_\theta^c$, by definition. Next, observe that

$$\theta r_\theta^2 = \frac{c_4(1)\theta}{\psi'(\lambda_\theta)} = c_4(1)\lambda_\theta^{-c}.$$

Thus, $\theta \mapsto \theta r_\theta^2$ decreases, which proves the first inequality in (65). To prove the second inequality, we fix $\varepsilon \in (0, \infty)$ such that $c + \varepsilon < \gamma_{\psi'}$. By definition of $\gamma_{\psi'}$, there exists $K \in (0, \infty)$ such that $\lambda^{-c} \psi'(\lambda) \geq K \lambda^\varepsilon$, for any $\lambda \in [1, \infty)$. It entails that $\theta = \lambda_\theta^{-c} \psi'(\lambda_\theta) \geq K \lambda_\theta^\varepsilon$. Thus,

$$r_\theta = (c_4(1)/\psi'(\lambda_\theta))^{\frac{1}{2}} = (c_4(1)/(\theta \lambda_\theta^c))^{\frac{1}{2}} \geq \left(c_4(1)K^{\frac{c}{\varepsilon}}\right)^{\frac{1}{2}} \theta^{-\frac{1}{2}(1+\frac{c}{\varepsilon})},$$

which implies the desired result with $c_{11} = (c_4(1)K^{\frac{c}{\varepsilon}})^{\frac{1}{2}}$ and $c_{12} = \frac{1}{2}(1 + \frac{c}{\varepsilon})$. □

By combing the previous lemmas we obtain the following result.

Lemma 3.7 *Assume that $\delta > 1$ and that $d > \frac{2\gamma}{\gamma-1}$. Then $d > 4$. Let $a \in (0, \infty)$. For any $n \in \mathbb{N}$, set $\rho_n = r_{e^{n^2}}$, where $(r_\theta)_{\theta \in [\psi'(1), \infty)}$ is defined as in Lemma 3.6. Then, the sequence $(\rho_n)_{n \geq 0}$ satisfies (51) in Lemma 3.1. Moreover, there exists a constant $c_{13} \in (0, \infty)$, that only depends on d, ψ and a , such that for all sufficiently large $n \in \mathbb{N}$,*

$$\mathbf{E}_0 \left[\int_0^{2a} ds \mathbf{1}_{\{\|\xi_s\| \geq 2\rho_n\}} \psi'(u_{\rho_n}(\xi_s)) \right] \leq c_{13}.$$

Proof Note that $\gamma \leq 2$, which easily entails that $d > 4$. By (65) in Lemma 3.6,

$$\begin{aligned} \rho_{n+1}/\rho_n &\leq \left(e^{n^2}/e^{(n+1)^2} \right)^{1/2} = e^{-n-\frac{1}{2}} \leq e^{-n} \quad \text{and} \\ n^{-2} \log 1/\rho_n &\leq c_{12} - n^{-2} \log c_{11}, \end{aligned}$$

which proves that $(\rho_n)_{n \geq 0}$ satisfies (51) in Lemma 3.1. Moreover, for all $n \in \mathbb{N}$ such that $\rho_n \in (0, r_2)$, Lemma 3.4 and (64) in Lemma 3.6 imply

$$\mathbf{E}_0 \left[\int_0^{2a} ds \mathbf{1}_{\{\|\xi_s\| \geq 2\rho_n\}} \psi'(u_{\rho_n}(\xi_s)) \right] \leq c_5 + c_6 J(r_{e_n^2}) \leq c_5 + c_6 c_{10} =: c_{13},$$

which completes the proof of the lemma. □

3.3 The spine and the associated subordinator

Recall from Sect. 2.4 the Palm formula for the occupation measure of the snake. Recall that $\xi = (\xi_t)_{t \geq 0}$ is d -dimensional Brownian motion starting from 0 that is defined on $(\Omega, \mathcal{F}, \mathbf{P}_0)$. Recall that $(V_t)_{t \geq 0}$ is a subordinator defined on $(\Omega, \mathcal{F}, \mathbf{P}_0)$ that is independent of ξ and whose Laplace exponent is $\psi^{*\prime}(\lambda) = \psi'(\lambda) - \alpha$. Recall from (47) that under \mathbf{P}_0 , conditionally given (ξ, V) , $\mathcal{N}^*(dt dW) = \sum_{j \in \mathcal{J}^*} \delta_{(t_j, W^j)}$ is a Poisson point process on $[0, \infty) \times C(\mathbb{R}_+, \mathcal{W})$ with intensity $dV_t \mathbb{N}_{\xi_t}(dW)$. Then recall from (48) that for all $a \in \mathbb{R}_+$, we have set $\mathcal{M}_a^* = \sum_{j \in \mathcal{J}^*} \mathbf{1}_{[0, a]}(t_j) \mathcal{M}_j$ where for all $j \in \mathcal{J}^*$, \mathcal{M}_j stands for the occupation measure of the snake W^j as defined in (38). For any $a \in \mathbb{R}_+$, we next set

$$T_a := \langle \mathcal{M}_a^*, \mathbf{1} \rangle = \sum_{j \in \mathcal{J}^*} \mathbf{1}_{[0, a]}(t_j) \sigma_j, \tag{67}$$

where σ_j is the total duration of the excursion of the snake W^j (and therefore, the total mass of \mathcal{M}_j). By construction of the snake excursion measure,

$$\mathbb{N}_x [1 - e^{-\lambda \sigma}] = N [1 - e^{-\lambda \sigma}] = \psi^{-1}(\lambda). \tag{68}$$

We shall assume throughout the paper that $d \geq 4$. We then introduce the following two last exit times: for all $r \in \mathbb{R}_+$, we set

$$\vartheta(r) = \sup\{t \geq 0 : \|\xi_t\| \leq r\} \tag{69}$$

$$\gamma(r) = \sup \left\{ t \geq 0 : \sqrt{(\xi_t^{(1)})^2 + (\xi_t^{(2)})^2 + (\xi_t^{(3)})^2} \leq r \right\} \tag{70}$$

where $(\xi_t^{(i)})_{t \geq 0}$ stands for the i -th coordinate of ξ . We then recall the two basic facts on the processes γ and ϑ .

- (a) The increments of $(\vartheta(r))_{r \geq 0}$ are independent. Moreover, $(\|\xi_{\vartheta(r)+t}\|)_{t \geq 0}$ is independent of the two processes $(\vartheta(r'))_{0 \leq r' \leq r}$ and $(\|\xi_{t \wedge \vartheta(r)}\|)_{t \geq 0}$.
- (b) The process $(\gamma(r))_{r \geq 0}$ has independent and stationary increments: it is a subordinator with Laplace exponent $\lambda \mapsto \sqrt{2\lambda}$.

The first point is proved in Gettoor [22]. The second is a celebrated result of Pitman [36].

We next define the following random variable

$$\forall t \geq s \geq 0, \quad N_r(s, t) = \#\{j \in \mathcal{J}^* : s < t_j < t \text{ and } \mathcal{R}_j \cap \overline{B}(0, r) \neq \emptyset\}, \tag{71}$$

that counts the number of snakes that are grafted on the spatial spine ξ between times s and t and that hit the ball $\bar{B}(0, r)$.

Lemma 3.8 *Assume that $d \geq 4$. We keep the previous notation. Then, the following holds true.*

(i) *For all real numbers $r > r' > \rho > \rho' > 0$ and all $a \in (0, \infty)$, the random variables*

$$T_{\vartheta(2\rho)} - T_{\vartheta(2\rho')}, \quad T_{\vartheta(2r)} - T_{\vartheta(2r')} \quad \text{and} \quad N_r(\vartheta(2r), a + \vartheta(2r))$$

are independent.

(ii) *The process $(T_{\gamma(r)})_{r \geq 0}$ is a subordinator with Laplace exponent $\sqrt{2 \psi^{*'} \circ \psi^{-1}}$.*

Remark 3.1 We take the opportunity to mention that it is stated incorrectly in [10, (60)], that $T_{\gamma(2\rho)} - T_{\gamma(2\rho')}$, $T_{\gamma(2r)} - T_{\gamma(2r')}$ and $N(\vartheta(2r), a)$ are independent. More precisely, the statement (49) in [10] is incorrect. We provide here a correct statement and a correct proof. □

Proof Let us prove (i). Let $\lambda, \mu, \theta \in \mathbb{R}_+$. To simplify notation we set

$$Y = \exp \left(-\lambda(T_{\vartheta(2r)} - T_{\vartheta(2r')}) - \mu(T_{\vartheta(2\rho)} - T_{\vartheta(2\rho')}) - \theta N_r(\vartheta(2r), a + \vartheta(2r)) \right).$$

Then recall (68) and recall from (58) the notation $\tilde{u}_r(\|x\|) = \mathbb{N}_x(\mathcal{R} \cap \bar{B}(0, r) \neq \emptyset)$. Then, basic results on Poisson processes imply

$$\begin{aligned} \mathbf{E}_0 [Y | (\xi, V)] &= \exp \left(-\psi^{-1}(\lambda)(V_{\vartheta(2r)} - V_{\vartheta(2r')}) - \psi^{-1}(\mu)(V_{\vartheta(2\rho)} - V_{\vartheta(2\rho')}) \right. \\ &\quad \left. - (1 - e^{-\theta}) \int_{(0,a)} dV_t \tilde{u}_r(\|\xi_{\vartheta(2r)+t}\|) \right). \end{aligned}$$

Since V is a subordinator with Laplace exponent ψ^{*} , we then get

$$\begin{aligned} \mathbf{E}_0 [Y | \xi] &= \exp \left(-(\vartheta(2r) - \vartheta(2r'))\psi^{*'}(\psi^{-1}(\lambda)) \right. \\ &\quad - (\vartheta(2\rho) - \vartheta(2\rho'))\psi^{*'}(\psi^{-1}(\mu)) \\ &\quad \left. - \int_0^a dt \psi^{*'} \left((1 - e^{-\theta})\tilde{u}_r(\|\xi_{\vartheta(2r)+t}\|) \right) \right). \end{aligned}$$

The above mentioned property (a) for last exit times then implies that

$$\vartheta(2r) - \vartheta(2r'), \quad \vartheta(2\rho) - \vartheta(2\rho') \quad \text{and} \quad \int_0^a dt \psi^{*'} \left((1 - e^{-\theta})\tilde{u}_r(\|\xi_{\vartheta(2r)+t}\|) \right)$$

are independent which easily implies (i).

The second point is proved in a similar way: let $0 = r_0 < r_1 < \dots < r_n$ and let $\lambda_1, \dots, \lambda_n \in \mathbb{R}_+$. We set

$$Y' = \exp \left(- \sum_{1 \leq k \leq n} \lambda_k (T_{\gamma(r_k)} - T_{\gamma(r_{k-1})}) \right)$$

Thus,

$$\begin{aligned} \mathbf{E}_0 [Y'] &= \mathbf{E}_0 \left[\exp \left(- \sum_{1 \leq k \leq n} \psi^{-1}(\lambda_k) (V_{\gamma(r_k)} - V_{\gamma(r_{k-1})}) \right) \right] \\ &= \mathbf{E}_0 \left[\exp \left(- \sum_{1 \leq k \leq n} \psi^{*'}(\psi^{-1}(\lambda_k)) (\gamma(r_k) - \gamma(r_{k-1})) \right) \right] \\ &= \exp \left(- \sum_{1 \leq k \leq n} (r_k - r_{k-1}) \sqrt{2\psi^{*'}(\psi^{-1}(\lambda_k))} \right). \end{aligned}$$

Indeed, the first equality comes from basic results on Poisson point measures, the second equality comes from the fact that V is a subordinator with Laplace exponent $\psi^{*'}$ and the last equality is a consequence of the above mentioned Property (b) of the last exit times $\gamma(r)$. This completes the proof of (ii). \square

Recall from (71) the notation $N_r(s, t)$ that counts the snakes that are grafted on the spatial spine ξ between times s and t and that hit the ball $\overline{B}(0, r)$. We state the following lemma that actually means, in some sense, that in supercritical dimensions, there is no snake W^j grafted far away that hits $\overline{B}(0, r)$.

Lemma 3.9 Assume that $\gamma > 1$ and that $d > \frac{2\gamma}{\gamma-1}$. Then, for all $t > s > 0$,

$$\forall t > s > 0, \quad \lim_{r \rightarrow 0} \mathbf{P}_0 (N_r(s, t) = 0) = 1.$$

Proof Recall from (56) the definition of u_r and from (58) that of \tilde{u}_r . By the definition (71), conditionally given (ξ, V) , $N_r(s, t)$ is a Poisson random variable with parameter $\int_s^t dV_w u_r(\xi_w)$. Thus

$$\mathbf{P}_0(N_r(s, t) = 0) = \mathbf{E}_0 \left[\exp \left(- \int_s^t dw \psi^{*'}(u_r(\xi_w)) \right) \right]. \tag{72}$$

Next, note that $d > \frac{2\gamma}{\gamma-1}$ implies that $d - 2 > \frac{2}{\gamma-1}$. Then, there exists $b \in (0, 1)$ and $a \in (0, \gamma - 1)$ such that

$$(d - 2)(1 - b) > \frac{2}{a}. \tag{73}$$

By the definition (69) of the last exit time process ϑ ,

$$\mathbf{E}_0 \left[\exp \left(- \int_s^t dw \psi^{*'}(u_r(\xi_w)) \right) \right] \geq \exp \left(-t \psi^{*'}(\tilde{u}_r(r^b)) \right) \mathbf{P}_0 \left(s > \vartheta(r^b) \right). \tag{74}$$

Clearly $\lim_{r \rightarrow 0} \mathbf{P}_0(s > \vartheta(r^b)) = 1$. Thus, we only have to choose b such that $\lim_{r \rightarrow 0} \tilde{u}_r(r^b) = 0$. To that end, we apply Lemma 3.3 with $\varrho = 1$: for all $r \in (0, r_1 \wedge 2^{-\frac{1}{1-b}})$, we get

$$\tilde{u}_r(r^b) \leq (2r/r^b)^{d-2} \psi'^{-1}(c_4(1)r^{-2}) = 2^{d-2} r^{(d-2)(1-b)} \psi'^{-1}(c_4(1)r^{-2}). \tag{75}$$

Since $a \in (0, \boldsymbol{\gamma} - 1)$ and since $\gamma_{\psi'} = \boldsymbol{\gamma} - 1$, the definition (21) of $\gamma_{\psi'}$ entails that for all sufficiently large λ , $\psi'(\lambda) \geq \lambda^a$ and thus $\psi'^{-1}(\lambda) \leq \lambda^{1/a}$. Then (75) implies that there exists a constant $c \in (0, \infty)$ such that $\tilde{u}_r(r^b) \leq c r^{(d-2)(1-b)-2/a}$ for all sufficiently small r . By (73), this entails $\lim_{r \rightarrow 0} \tilde{u}_r(r^b) = 0$, which completes the proof by (72) and (74). □

We next prove a similar estimate for $N_r(\vartheta(2r), a + \vartheta(2r))$.

Lemma 3.10 *Assume that $\delta > 1$ and that $d > \frac{2\delta}{\delta-1}$. Then, there exist two constants $c_{14}, r_3 \in (0, \infty)$ that only depends on d, ψ and a , such that*

$$\forall r \in (0, r_3) \quad \mathbf{P}_0(N_r(\vartheta(2r), a + \vartheta(2r)) = 0) \geq c_{14}.$$

Proof Recall from (56) the definition of u_r and from (58) that of \tilde{u}_r . By the definition (71), conditionally given (ξ, V) , $N_r(\vartheta(2r), a + \vartheta(2r))$ is a Poisson random variable with parameter $\int_{\vartheta(2r)}^{a+\vartheta(2r)} dV_t u_r(\xi_t)$. Thus

$$\mathbf{P}_0(N_r(\vartheta(2r), a + \vartheta(2r)) = 0) = \mathbf{E}_0 \left[\exp \left(- \int_{\vartheta(2r)}^{a+\vartheta(2r)} dt \psi^{*'}(u_r(\xi_t)) \right) \right]. \tag{76}$$

Next note that on $\{\vartheta(2r) \leq a\}$, $a + \vartheta(2r) \leq 2a$ and that $t \geq \vartheta(2r)$ implies that $\|\xi_t\| \geq 2r$. Thus, by (76)

$$\begin{aligned} & \mathbf{P}_0(N_r(\vartheta(2r), a + \vartheta(2r)) = 0) \\ & \geq \mathbf{E}_0 \left[\mathbf{1}_{\{\vartheta(2r) \leq a\}} \exp \left(- \int_0^{2a} dt \mathbf{1}_{\{\|\xi_t\| \geq 2r\}} \psi^{*'}(u_r(\xi_t)) \right) \right] \\ & \geq \mathbf{E}_0 \left[\exp \left(- \int_0^{2a} dt \mathbf{1}_{\{\|\xi_t\| \geq 2r\}} \psi^{*'}(u_r(\xi_t)) \right) \right] - \mathbf{P}_0(\vartheta(2r) > a) \\ & \geq \exp \left(-\mathbf{E}_0 \left[\int_0^{2a} dt \mathbf{1}_{\{\|\xi_t\| \geq 2r\}} \psi^{*'}(u_r(\xi_t)) \right] \right) - \mathbf{P}_0(\vartheta(2r) > a), \end{aligned} \tag{77}$$

where we use Jensen inequality in the last line. By Lemmas 3.4 and 3.5, for any $r \in (0, r_2)$,

$$\exp\left(-\mathbf{E}_0\left[\int_0^{2a} dt \mathbf{1}_{\{\|\xi_t\|\geq 2r\}} \psi^{*'}(u_r(\xi_t))\right]\right) \geq \exp(-c_5 - c_6c_9).$$

Since $\mathbf{P}_0(\vartheta(2r) > a) \rightarrow 0$ as $r \rightarrow 0$, there exists $r_3 \in (0, r_2)$ such that $\mathbf{P}_0(\vartheta(2r_3) > a) \leq \frac{1}{2}e^{-c_5 - c_6c_9}$. Thus by (77) and the previous inequality, for any $r \in (0, r_3)$,

$$\mathbf{P}_0(N_r(\vartheta(2r), a + \vartheta(2r)) = 0) \geq \frac{1}{2} \exp(-c_5 - c_6c_9) =: c_{14},$$

which completes the proof of lemma. □

Under the less restrictive condition $d > \frac{2\gamma}{\gamma-1}$, we get a similar lower bound but only for the family of radii $(\rho_n)_{n \geq 1}$ introduced in Lemma 3.7.

Lemma 3.11 *Assume that $\delta > 1$ and that $d > \frac{2\gamma}{\gamma-1}$. Let $(\rho_n)_{n \geq 1}$ be the sequence introduced in Lemma 3.7. Then, there exists a constant $c_{15} \in (0, \infty)$, that only depend on d, ψ and a , such that*

$$\forall n \geq 1 \quad \mathbf{P}_0(N_{\rho_n}(\vartheta(2\rho_n), a + \vartheta(2\rho_n)) = 0) \geq c_{15}.$$

Proof The lower bound (77) applies for $r = \rho_n$. Then, Lemma 3.7, entails that for all sufficiently large n ,

$$\exp\left(-\mathbf{E}_0\left[\int_0^{2a} dt \mathbf{1}_{\{\|\xi_t\|\geq 2\rho_n\}} \psi^{*'}(u_{\rho_n}(\xi_t))\right]\right) \geq \exp(-c_{13}),$$

and we completes the proof arguing as in the proof of Lemma 3.10 with $c_{15} := \frac{1}{2}e^{-c_{13}}$. □

We then shall need the following result that is used in the proof of Lemma 4.4.

Lemma 3.12 *Assume that $\delta > 1$ and that $d \geq 4$. Let $(\rho_n)_{n \geq 1}$ be the sequence of radii introduced in Lemma 3.7. Then,*

$$(i) : \sum_{n \geq 1} \mathbf{P}_0(T_{\gamma(2\rho_n)} \leq g(8\rho_n)) = \infty \quad \text{and} \quad (ii) : \mathbf{P}_0\text{-a.s.} \limsup_{n \rightarrow \infty} \frac{T_{\gamma(2\rho_{n+1})}}{g(8\rho_n)} < \infty.$$

Proof Lemma 3.8 shows that $T_{\gamma(\cdot)}$ is a subordinator with Laplace exponent $\sqrt{2\psi^{*'} \circ \psi^{-1}}$. Then, Lemma 3.7 asserts that the sequence $(\rho_n)_{n \geq 0}$ satisfies the conditions (51) in Lemma 3.1, which immediately entails (i) and (ii). □

We end this section with the following result that is close to the previous one and that is used in the proof of Theorem 1.1.

Lemma 3.13 Assume that $\gamma > 1$. Let $u \in (0, \frac{2\gamma}{\gamma-1})$. Then, there exists a decreasing sequence $(s_n)_{n \geq 1}$ that tends to 0, that only depends on ψ and u and that satisfies the following.

$$(i) : \sum_{n \geq 1} \mathbf{P}_0 (T_{\gamma(2s_n)} \leq s_n^u) = \infty \quad \text{and} \quad (ii) : \mathbf{P}_0\text{-a.s.} \limsup_{n \rightarrow \infty} \frac{T_{\gamma(2s_{n+1})}}{s_n^u} < \infty.$$

Proof Let $u' \in (u, \frac{2\gamma}{\gamma-1})$. We set $\varphi^* = \psi^{*'} \circ \psi^{-1}$, where $\psi^*(\lambda) = \psi(\lambda) - \alpha$. Recall from (21) that $\gamma_\varphi = \gamma_{\varphi^*} = \frac{\gamma-1}{\gamma}$. We fix $a \in (0, \frac{\gamma-1}{\gamma})$. By the definition (21) of γ_{φ^*} , there exists $\lambda_0 \in (0, \infty)$ such that $\varphi^*(\lambda) \geq \lambda^a$ for any $\lambda \in [\lambda_0, \infty)$. Next observe that $2/u' > \frac{\gamma-1}{\gamma} = \gamma_{\varphi^*}$. As an easy consequence of definition (21) of γ_{φ^*} , we get $\liminf_{\lambda \rightarrow \infty} \varphi^*(\lambda)\lambda^{-2/u'} = 0$. Consequently, there exists an increasing sequence $(\lambda_n)_{n \geq 0}$ such that

$$\forall n \in \mathbb{N}, \quad 2^n \leq \lambda_n \quad \text{and} \quad \lambda_n^a \leq \varphi^*(\lambda_n) \leq \lambda_n^{2/u'}. \tag{78}$$

We next fix $\varepsilon \in (0, \infty)$ such that $(1 + \varepsilon)u/u' < 1$, which is possible since $u' > u$. Then, we set

$$\forall n \in \mathbb{N}, \quad s_n = \varphi^*(\lambda_n)^{-\frac{1+\varepsilon}{2}}. \tag{79}$$

Since, by Lemma 3.8, $(T_{\gamma(r)})_{r \geq 0}$ is a subordinator with Laplace exponent $\sqrt{2\varphi^*}$, a Markov inequality entails

$$\mathbf{P}_0 (T_{\gamma(2s_n)} > s_n^u) \leq \frac{1 - \exp(-2\sqrt{2}s_n\sqrt{\varphi^*(\lambda_n)})}{1 - \exp(-\lambda_n s_n^u)} \leq 2\sqrt{2} \frac{s_n\sqrt{\varphi^*(\lambda_n)}}{1 - \exp(-\lambda_n s_n^u)}. \tag{80}$$

By (79) and the last inequality of (78), we get

$$\lambda_n s_n^u = \lambda_n \varphi^*(\lambda_n)^{-(1+\varepsilon)u/2} \geq \lambda_n^{1-(1+\varepsilon)u/u'} \xrightarrow{n \rightarrow \infty} \infty. \tag{81}$$

Moreover, the first two inequalities in (78) and (79) imply

$$s_n\sqrt{\varphi^*(\lambda_n)} = \varphi^*(\lambda_n)^{-\varepsilon/2} \leq \lambda_n^{-a\varepsilon/2} \leq 2^{-na\varepsilon/2}. \tag{82}$$

Then (80), (81) and (82) imply that there exists $c \in (0, \infty)$ such that

$$\mathbf{P}_0 (T_{\gamma(2s_n)} > s_n^u) \leq c2^{-na\varepsilon/2}.$$

It immediately implies (i). Point (ii) follows from $\mathbf{P}_0 (T_{\gamma(2s_{n+1})} > s_n^u) \leq \mathbf{P}_0 (T_{\gamma(2s_n)} > s_n^u)$ and from the Borel–Cantelli lemma. □

3.4 Estimates for bad points

Recall from (36) the definition of the excursion measure \mathbb{N}_0 of the ψ -Lévy snake W . Recall that the lifetime process of W is the height process H . Namely, \mathbb{N}_0 -a.e. for all $t \in \mathbb{R}_+$, $\zeta_{W_t} = H_t$. Therefore, the duration σ of W under \mathbb{N}_0 is the duration of the excursion of H . Recall that \widehat{W} is the endpoint process of the snake. Recall from (38) that \mathcal{M} stands for the occupation measure of the snake, namely the random measure on \mathbb{R}^d that is the image via \widehat{W} of the Lebesgue measure on $[0, \sigma]$. Recall from (37) the definition of \mathcal{R} , the range of the snake.

Let $\lambda, r \in (0, \infty)$. Note that $\langle \mathcal{M} \rangle = \sigma$, therefore we get

$$\mathbb{N}_0 \left[1 - e^{-\lambda \mathcal{M}(B(0,r))} \right] \leq \mathbb{N}_0 \left[1 - e^{-\lambda \sigma} \right] = N \left[1 - e^{-\lambda \sigma} \right] = \psi^{-1}(\lambda), \tag{83}$$

where the last two equalities come from (25) and (68). The next lemma states a lower bound of the same kind.

Lemma 3.14 *There exists $c_{16} \in (0, 1)$, that only depends on d , such that*

$$\forall \mu, r \in (0, \infty), \quad \frac{r^2 \mu}{\psi^{-1}(\mu)} \geq 1 \implies \mathbb{N}_0 \left[1 - e^{-\mu \mathcal{M}(B(0,r))} \right] \geq c_{16} \psi^{-1}(\mu). \tag{84}$$

Proof To simplify, set $q(\mu, r) := \mathbb{N}_0 \left[1 - e^{-\mu \mathcal{M}(B(0,r))} \right]$. Note that $\mathcal{M}(B(0, r)) = \int_0^\sigma dt \mathbf{1}_{\{\|\widehat{W}_t\| < r\}}$. An easy argument combined with Fubini first entails the following:

$$q(\mu, r) = \mu \int_0^\infty dt \mathbb{N}_0 \left[\mathbf{1}_{\{t \leq \sigma; \|\widehat{W}_t\| < r\}} e^{-\mu \int_t^\sigma ds \mathbf{1}_{\{\|\widehat{W}_s\| < r\}}} \right]. \tag{85}$$

We next apply Lemma 2.8 at the (deterministic) time t (recall that this lemma is a specific form of the Markov property for $\overline{W} = (\rho, W)$). Then, we get for all $t \in \mathbb{R}_+$

$$\begin{aligned} & \mathbb{N}_0 \left[\mathbf{1}_{\{t \leq \sigma; \|\widehat{W}_t\| < r\}} e^{-\mu \int_t^\sigma ds \mathbf{1}_{\{\|\widehat{W}_s\| < r\}}} \right] \\ &= \mathbb{N}_0 \left[\mathbf{1}_{\{t \leq \sigma; \|\widehat{W}_t\| < r\}} \exp \left(- \int_{[0, H_t]} \rho_t(dh) \mathbb{N}_{W_t(h)} \left[1 - e^{-\mu \mathcal{M}(B(0,r))} \right] \right) \right]. \end{aligned} \tag{86}$$

From (83), we get that for all $t, h \geq 0$, $\mathbb{N}_{W_t(h)} \left[1 - e^{-\mu \mathcal{M}(B(0,r))} \right] \leq \psi^{-1}(\mu)$. Then,

$$\begin{aligned} & \mathbb{N}_0 \left[\mathbf{1}_{\{t \leq \sigma; \|\widehat{W}_t\| < r\}} \exp \left(- \int_{[0, H_t]} \rho_t(dh) \mathbb{N}_{W_t(h)} \left[1 - e^{-\mu \mathcal{M}(B(0,r))} \right] \right) \right] \\ & \geq \mathbb{N}_0 \left[\mathbf{1}_{\{t \leq \sigma; \|\widehat{W}_t\| < r\}} e^{-\psi^{-1}(\mu) \int_{[0, H_t]} \rho_t(dh)} \right] \end{aligned} \tag{87}$$

Then by (85), (86), (87) and Fubini we get

$$q(\mu, r) \geq \mu \mathbb{N}_0 \left[\int_0^\sigma dt \mathbf{1}_{\{\|\widehat{W}_t\| < r\}} e^{-\psi^{-1}(\mu) \int_{[0, H_t]} \rho_t(dh)} \right]. \tag{88}$$

We next apply (46) to the right member of the previous inequality: to that end, recall that $U = (U_a)_{a \geq 0}$ stands for a subordinator defined on $(\Omega, \mathcal{F}, \mathbf{P}_0)$ that is independent of ξ and whose Laplace exponent is $\tilde{\psi}^*(\lambda) = \psi(\lambda)/\lambda - \alpha$. We then apply (46) to the right member of (88) and we get

$$\begin{aligned} & \mu \mathbb{N}_0 \left[\int_0^\sigma dt \mathbf{1}_{\{\|\widehat{W}_t\| < r\}} e^{-\psi^{-1}(\mu) \int_{[0, H_t]} \rho_t(dh)} \right] \\ &= \mu \int_0^\infty da e^{-\alpha a} \mathbf{E}_0 \left[\mathbf{1}_{\{\|\xi_a\| < r\}} e^{-\psi^{-1}(\mu) U_a} \right] \\ &= \mu \int_0^\infty da e^{-a\mu/\psi^{-1}(\mu)} \mathbf{P}_0(\|\xi_a\| < r), \end{aligned}$$

since ξ and U are independent. Thus, (88) and a simple change of variable using the scaling property of Brownian motion, entail

$$q(\mu, r) \geq \psi^{-1}(\mu) \int_0^\infty \mathbf{P}_0 \left(\|\xi_c\| \leq r\sqrt{\mu/\psi^{-1}(\mu)} \right) e^{-c} dc. \tag{89}$$

If $r^2\mu/\psi^{-1}(\mu) \geq 1$, then (89) implies (84) with $c_{16} := \int_0^\infty \mathbf{P}_0(\|\xi_c\| \leq 1) e^{-c} dc$. Clearly, c_{16} only depends on d and $c_{16} \in (0, 1)$. \square

Before stating the next lemma, we recall a result about the first exit time from a ball for a d -dimensional Brownian motion. First, we set

$$\chi_{d,r} := \inf \{t \in \mathbb{R}_+ : \|\xi_t\| = r\}. \tag{90}$$

In dimension 1, we easily prove that

$$\forall r, \lambda \in \mathbb{R}_+, \quad \mathbf{E}_0 [\exp(-\lambda \chi_{1,r})] = \left(\cosh(r\sqrt{2\lambda}) \right)^{-1} \leq 2 \exp(-r\sqrt{2\lambda}). \tag{91}$$

(Indeed, note that $t \mapsto \exp(\sqrt{2\lambda}\xi_t - \lambda t)$ is a martingale: see e.g. Revuz and Yor [37, Chapter II, (3.7)]). In dimension d , observe that

$$\mathbf{P}_0\text{-a.s. } \chi_{d,r} \geq \min_{1 \leq j \leq d} \inf \left\{ t \in \mathbb{R}_+ : \sqrt{d} |\xi_t^{(j)}| = r \right\},$$

where $\xi_t^{(j)}$ stands for the j -th coordinate of ξ_t . The previous inequality combined with (91) then entails

$$\forall \lambda, r \in \mathbb{R}_+, \quad \mathbf{E}_0 [\exp(-\lambda \chi_{d,r}(\xi))] \leq 2d \exp(-r\sqrt{2\lambda/d}). \tag{92}$$

Lemma 3.15 *Assume that $\delta > 1$ and that $d \geq 3$. Recall that r_1 and $c_4(\frac{1}{4})$ are the constants appearing in Lemma 3.3. with $\varrho = \frac{1}{4}$. Then, there exist $c_{17}, c_{18} \in (0, \infty)$, that only depend on d and ψ , such that*

$$\begin{aligned} &\forall \mu \in (0, \infty), \forall r \in (0, r_1) \text{ such that } \frac{r^2 \mu}{\psi^{-1}(\mu)} \geq 16, \quad \forall x \in \mathbb{R}^d \setminus \overline{B}(0, 2r), \\ &\mathbb{N}_x \left[\mathbf{1}_{\{\mathcal{R} \cap \overline{B}(0,r) \neq \emptyset\}} e^{-\mu \mathcal{M}(B(0,2r))} \right] \\ &\leq c_{17} (r/\|x\|)^{d-2} \psi'^{-1} \left(c_4 \left(\frac{1}{4} \right) r^{-2} \right) \exp \left(-c_{18} r \sqrt{\varphi(\mu)} \right). \end{aligned}$$

Proof We fix $\mu \in (0, \infty)$ and $x \in \mathbb{R}^d \setminus \overline{B}(0, 2r)$. To simplify notation we set

$$p(x, \mu, r) = \mathbb{N}_x \left[\mathbf{1}_{\{\mathcal{R} \cap \overline{B}(0,r) \neq \emptyset\}} e^{-\mu \mathcal{M}(B(0,2r))} \right].$$

Recall from (40) the definition of τ_r that is the hitting time in $\overline{B}(0, r)$ of the snake W . First recall that $\{\mathcal{R} \cap \overline{B}(0, r) \neq \emptyset\} = \{\tau_r < \infty\}$ and observe that

$$\mathbb{N}_x\text{-a.e. on the event } \{\mathcal{R} \cap \overline{B}(0, r) \neq \emptyset\}, \quad \int_{\tau_r}^{\sigma} \mathbf{1}_{\{\|\widehat{w}_s\| < 2r\}} ds \leq \mathcal{M}(B(0, 2r)).$$

Thus, we get

$$p(x, \mu, r) \leq \mathbb{N}_x \left[\mathbf{1}_{\{\tau_r < \infty\}} e^{-\mu \int_{\tau_r}^{\sigma} \mathbf{1}_{\{\|\widehat{w}_s\| < 2r\}} ds} \right]. \tag{93}$$

We next apply Lemma 2.8 to the right member of (93) and to the stopping time τ_r ; we get

$$p(x, \mu, r) \leq \mathbb{N}_x \left[\mathbf{1}_{\{\tau_r < \infty\}} \exp \left(- \int_{[0, H_{\tau_r}] } \rho_{\tau_r}(dh) \mathbb{N}_{W_{\tau_r}(h)} \left[1 - e^{-\mu \mathcal{M}(B(0,2r))} \right] \right) \right]. \tag{94}$$

Let $w \in \mathcal{W}$ be a continuous stopped path starting from $x \in \mathbb{R}^d \setminus \overline{B}(0, 2r)$; we denote by ζ_w its lifetime. We define $T_1(w), T_2(w)$ and $T_3(w)$ by

$$\begin{aligned} T_1(w) &= \inf \{t \in [0, \zeta_w] : \|w(t)\| \leq 3r/2\} \\ T_2(w) &= \inf \{t \in [0, \zeta_w - T_1(w)] : \|w(t + T_1(w)) - w(T_1(w))\| > r/4\} \\ T_3(w) &= \inf \{t \in [0, \zeta_w] : \|w(t)\| \leq 5r/4\} \end{aligned} \tag{95}$$

with the convention that $\inf \emptyset = \infty$. Observe that if $T_3(w) < \infty$, then $T_1(w) + T_2(w) \leq T_3(w)$. Moreover, since $x \in \overline{B}(0, 2r)^c$, \mathbb{N}_x -a.e. on the event $\{\tau_r < \infty\}$, we have $T_1(W_{\tau_r}) + T_2(W_{\tau_r}) \leq T_3(W_{\tau_r}) < \tau_r < \infty$ and for any $t \in [T_1(W_{\tau_r}), T_1(W_{\tau_r}) +$

$T_2(W_{\tau_r})$], the following inequality holds true:

$$\begin{aligned} \mathbb{N}_{W_{\tau_r}(t)} \left[1 - e^{-\mu \mathcal{M}(B(0,2r))} \right] &\geq \mathbb{N}_{W_{\tau_r}(t)} \left[1 - e^{-\mu \mathcal{M}(B(W_{\tau_r}(t),r/4))} \right] \\ &= \mathbb{N}_0 \left[1 - e^{-\mu \mathcal{M}(B(0,r/4))} \right] =: \Lambda_{\mu,r}, \end{aligned}$$

the last equality being a consequence of the invariance of the snake under translation; here $\Lambda_{\mu,r}$ only depends on μ and r . To simplify notation we set $T_1 = T_1(W_{\tau_r})$ and $T_2 = T_2(W_{\tau_r})$. An elementary inequality combined with (94) entails

$$\begin{aligned} p(x, \mu, r) &\leq \mathbb{N}_x \left[\mathbf{1}_{\{\tau_r < \infty\}} \exp \left(- \int_{[T_1, T_1+T_2]} \rho_{\tau_r}(dh) \mathbb{N}_{W_{\tau_r}(h)} \left[1 - e^{-\mu \mathcal{M}(B(0,2r))} \right] \right) \right] \\ &\leq \mathbb{N}_x \left[\mathbf{1}_{\{\tau_r < \infty\}} \exp \left(- \Lambda_{\mu,r} \int_{[T_1, T_1+T_2]} \rho_{\tau_r}(dh) \right) \right]. \end{aligned} \tag{96}$$

Recall from (95) the definition of T_3 . Recall from (44) the definition of the function ϖ . We next apply Lemma 2.9 with $r' = \frac{5}{4}r$ to the right member of (96). Then we get

$$\begin{aligned} &\mathbb{N}_x \left[\mathbf{1}_{\{\tau_r < \infty\}} \exp \left(- \Lambda_{\mu,r} \int_{[T_1, T_1+T_2]} \rho_{\tau_r}(dh) \right) \right] \\ &= \tilde{u}_r(5r/4) \mathbf{E}_x \left[\mathbf{1}_{\{T_3(\xi) < \infty\}} \exp \left(- \int_0^{T_3(\xi)} dt \varpi \left(u_r(\xi_t), \Lambda_{\mu,r} \mathbf{1}_{[T_1(\xi), T_1(\xi)+T_2(\xi)]}(t) \right) \right) \right] \\ &\leq \tilde{u}_r(5r/4) \mathbf{E}_x \left[\mathbf{1}_{\{T_1(\xi) < \infty\}} \exp \left(- \int_{T_1(\xi)}^{T_1(\xi)+T_2(\xi)} dt \varpi \left(u_r(\xi_t), \Lambda_{\mu,r} \right) \right) \right]. \end{aligned} \tag{97}$$

Here recall that ξ under \mathbf{P}_x is distributed as a standard d -dimensional Brownian motion starting from x . The convexity of ψ provides the following lower bounds for $\varpi(u_r(\xi_t), \Lambda_{\mu,r})$:

- if $u_r(\xi_t) \geq \Lambda_{\mu,r}$, then $\varpi(u_r(\xi_t), \Lambda_{\mu,r}) \geq \psi'(\Lambda_{\mu,r})$;
- if $u_r(\xi_t) < \Lambda_{\mu,r}$, then $\varpi(u_r(\xi_t), \Lambda_{\mu,r}) \geq \psi(\Lambda_{\mu,r})/\Lambda_{\mu,r} \geq \frac{1}{4}\psi'(\Lambda_{\mu,r})$, by (20) for the last inequality.

These inequalities combined with (96) and (97), entail

$$p(x, \mu, r) \leq \tilde{u}_r(5r/4) \mathbf{E}_x \left[\mathbf{1}_{\{T_1(\xi) < \infty\}} \exp \left(- \frac{1}{4} \psi'(\Lambda_{\mu,r}) T_2(\xi) \right) \right]. \tag{98}$$

We now assume that

$$\frac{r^2 \mu}{16 \psi^{-1}(\mu)} \geq 1.$$

Recall that $\Lambda_{\mu,r} = \mathbb{N}_0[1 - e^{-\mu \mathcal{M}(B(0,r/4))}]$. By Lemma 3.14, $\Lambda_{\mu,r} \geq c_{16} \psi^{-1}(\mu)$. We next use the concavity of ψ' and the fact that $c_{16} \in (0, 1)$, to get

$$\psi'(\Lambda_{\mu,r}) \geq \psi'(c_{16} \psi^{-1}(\mu)) \geq c_{16} \psi'(\psi^{-1}(\mu)) = c_{16} \varphi(\mu). \tag{99}$$

Recall that r_1 and $c_4(\frac{1}{4})$ are the constants appearing in Lemma 3.3 (with $\varrho = \frac{1}{4}$). We assume that $r \in (0, r_1)$. Then, by Lemma 3.3, $\tilde{u}_r(5r/4) \leq \psi'^{-1}(c_4(\frac{1}{4})r^{-2})$. Thus, by (98) and (99) we get

$$p(x, \mu, r) \leq \psi'^{-1} \left(c_4 \left(\frac{1}{4} \right) r^{-2} \right) \mathbf{E}_x \left[\mathbf{1}_{\{T_1(\xi) < \infty\}} \exp \left(-\frac{1}{4} c_{16} \varphi(\mu) T_2(\xi) \right) \right]. \tag{100}$$

Recall from (95) the definition of $T_1(\xi)$ and $T_2(\xi)$. By the Markov property at time $T_1(\xi)$, we get

$$\begin{aligned} & \mathbf{E}_x \left[\mathbf{1}_{\{T_1(\xi) < \infty\}} \exp \left(-\frac{1}{4} c_{16} \varphi(\mu) T_2(\xi) \right) \right] \\ &= \mathbf{P}_x(T_1(\xi) < \infty) \mathbf{E}_0 \left[\exp \left(-\frac{1}{4} c_{16} \varphi(\mu) \chi_{d,r/4}(\xi) \right) \right]. \end{aligned}$$

Since $d \geq 3$, we get $\mathbf{P}_x(T_1(\xi) < \infty) = (3r/2\|x\|)^{d-2}$. Moreover by (92), we get

$$\mathbf{E}_0 \left[\exp \left(-\frac{1}{4} c_{16} \varphi(\mu) \chi_{d,r/4}(\xi) \right) \right] \leq 2d \exp \left(-\frac{1}{8} r \sqrt{2c_{16} \varphi(\mu)/d} \right).$$

Then we set $c_{17} = 2d(3/2)^{d-2}$ and $c_{18} = \sqrt{c_{16}/(32d)}$ and we get

$$\mathbf{E}_x \left[\mathbf{1}_{\{T_1(\xi) < \infty\}} \exp \left(-\frac{1}{4} c_{16} \varphi(\mu) T_2(\xi) \right) \right] \leq c_{17} (r/\|x\|)^{d-2} \exp \left(-c_{18} r \sqrt{\varphi(\mu)} \right),$$

which implies the desired result by (100). □

Recall from (11) the definition of the gauge function g :

$$g(r) = \frac{\log \log \frac{1}{r}}{\varphi^{-1} \left(\left(\frac{1}{r} \log \log \frac{1}{r} \right)^2 \right)}.$$

Lemma 3.16 *Assume that $\delta > 1$ and that $d \geq 3$. Recall that c_{17} is the constant appearing in Lemma 3.15 and that $c_4(\frac{1}{4})$ is the constant appearing in Lemma 3.3 with $\varrho = \frac{1}{4}$. There exists $c_{19}, c_{20}, r_4, \kappa_0 \in (0, \infty)$, that only depend d and ψ , such that for all $r \in (0, r_4)$, for all $\kappa \in (0, \kappa_0)$ and for all $x \in \mathbb{R}^d \setminus \overline{B}(0, 2r)$,*

$$\begin{aligned} & \mathbb{N}_x \left(\mathcal{R} \cap \overline{B}(0, r) \neq \emptyset ; \mathcal{M}(B(0, 2r)) \leq \kappa g(r) \right) \\ & \leq c_{17} \psi'^{-1} \left(c_4 r^{-2} \right) (r/\|x\|)^{d-2} (\log 1/r)^{-c_{19} \kappa^{-c_{20}}}. \end{aligned}$$

Proof Let $\kappa, \mu \in (0, \infty)$. Let $r \in (0, r_1)$ where r_1 is the constant appearing in Lemma 3.3. Assume that $\frac{r^2 \mu}{\psi^{-1}(\mu)} \geq 16$. A Markov inequality, combined with Lemma 3.15 entails:

$$\begin{aligned} \mathbb{N}_x (\mathcal{R} \cap \overline{B}(0, r) \neq \emptyset; \mathcal{M}(B(0, 2r))) &\leq \kappa g(r) \\ &\leq e^{\kappa \mu g(r)} \mathbb{N}_x \left[\mathbf{1}_{\{\mathcal{R} \cap \overline{B}(0, r) \neq \emptyset\}} e^{-\mu \mathcal{M}(B(0, 2r))} \right] \\ &\leq c_{17} (r/\|x\|)^{d-2} \psi'^{-1} \left(c_4 r^{-2} \right) \exp(F(\mu, r, \kappa)), \end{aligned} \tag{101}$$

where we have set

$$F(\mu, r, \kappa) := \kappa \mu g(r) - c_{18} r \sqrt{\varphi(\mu)}.$$

For any $q \in [1, \infty)$ and any $r \in (0, r_0)$, we set

$$\mu_{r,q} = \varphi^{-1} \left(q \left(\frac{1}{r} \log \log \frac{1}{r} \right)^2 \right).$$

We first get an estimate for $\mu_{q,r}$. To that end, we fix a such that $1/a \in (0, \delta_\varphi)$. By the definition (22) of δ_φ , there exists $C \in (0, \infty)$ such that for any $p, \lambda \in [1, \infty)$, $Cp^{1/a}\varphi(\lambda) \leq \varphi(p\lambda)$. Thus, for any $z \geq \varphi(1)$ and any $q \geq C$, we get $\varphi^{-1}(qz) \leq (q/C)^a \varphi^{-1}(z)$. This easily entails that there exists $r_5 \in (0, r_0 \wedge r_1)$ and $c_{21} \in (0, \infty)$ such that

$$\forall r \in (0, r_5), \forall q \in [1, \infty), \quad \mu_{r,1} \leq \mu_{r,q} \leq c_{21} q^a \mu_{r,1}. \tag{102}$$

Recall that $\tilde{\psi}(\lambda) = \psi(\lambda)/\lambda$. We next observe that for all $r \in (0, e^{-1})$ and all $q \in [1, \infty)$, (20) implies

$$\begin{aligned} r^2 \frac{\mu_{r,q}}{\psi^{-1}(\mu_{r,q})} &= r^2 \tilde{\psi}(\psi^{-1}(\mu_{r,q})) \\ &\geq \frac{1}{4} r^2 \psi'(\psi^{-1}(\mu_{r,q})) \geq \frac{1}{4} r^2 \psi'(\psi^{-1}(\mu_{r,1})) = \frac{1}{4} (\log \log 1/r)^2. \end{aligned}$$

Then we set $r_4 = \exp(-e^8) \wedge r_5$ and we get

$$\forall r \in (0, r_4), \forall q \in [1, \infty), \quad \frac{r^2 \mu_{r,q}}{\psi^{-1}(\mu_{r,q})} \geq 16. \tag{103}$$

Let $r \in (0, r_4)$, $q \in [1, \infty)$ and $\kappa \in (0, \infty)$. Observe that $g(r) = (\log \log 1/r)/\mu_{r,1}$ and that $r\sqrt{\varphi(\mu_{r,q})} = \sqrt{q} \log \log 1/r$. Thus, we get

$$F(\mu_{r,q}, r, \kappa) = \left(\kappa \frac{\mu_{r,q}}{\mu_{r,1}} - c_{18} \sqrt{q} \right) \log \log \frac{1}{r} \leq \sqrt{q} \left(c_{21} \kappa q^{a-\frac{1}{2}} - c_{18} \right) \log \log \frac{1}{r}.$$

Since $\delta_\varphi \leq 1$, we get $a > 1$ and thus $a - \frac{1}{2} > \frac{1}{2}$. We then set $\kappa_0 = c_{18}/(2c_{21})$ and for all $\kappa \in (0, \kappa_0)$ we also set $q_\kappa = (\kappa_0/\kappa)^{\frac{1}{a-\frac{1}{2}}}$. Then $q_\kappa \geq 1$ and

$$\sqrt{q_\kappa} \left(c_{21} \kappa q_\kappa^{a-\frac{1}{2}} - c_{18} \right) = -\frac{1}{2} c_{18} (\kappa_0/\kappa)^{\frac{1}{2a-1}}.$$

We then set $c_{19} = \frac{1}{2}c_{18}\kappa_0^{\frac{1}{2a-1}}$ and $c_{20} = \frac{1}{2a-1}$. Then

$$F(\mu_{r,q\kappa}, r, \kappa) \leq -c_{19}\kappa^{-c_{20}} \log \log \frac{1}{r}$$

and we complete the proof thanks to (103) that allows to apply (101) for any $r \in (0, r_4)$. □

4 Proof of the results

Recall from (38) the definition of the total occupation measure of the snake \mathcal{M} . We prove in Sect. 4.1 the following results on the lower density of \mathcal{M} .

Theorem 4.1 *Let ψ be a branching mechanism the form (1). Let g be defined by (11). Assume that $\delta > 1$ and that $d > \frac{2\gamma}{\gamma-1}$. Let \mathcal{M} be the total occupation of the ψ -Lévy snake under the excursion measure \mathbb{N}_0 . Then, there exists a constant $\kappa_{d,\psi} \in (0, \infty)$, that only depends on d and ψ , such that*

$$\mathbb{N}_0\text{-a.e. for } \mathcal{M}\text{-almost all } x, \quad \liminf_{r \rightarrow 0_+} \frac{\mathcal{M}(B(x, r))}{g(r)} = \kappa_{d,\psi}. \tag{104}$$

This result is then used to prove the following theorem in Sect. 4.2.

Theorem 4.2 *Let ψ be a branching mechanism the form (1). Let g be defined by (11). Assume that $\delta > 1$ and that $d > \frac{2\gamma}{\gamma-1}$. Let \mathcal{M} be the total occupation of the ψ -Lévy snake under the excursion measure \mathbb{N}_0 . Then, there exists a constant $\kappa_{d,\psi} \in (0, \infty)$, that only depends on d and ψ , such that*

$$\mathbb{N}_0\text{-a.e. for any Borel set } B, \quad \mathcal{M}(B) = \kappa_{d,\psi} \mathcal{P}_g(B \cap \mathcal{R}).$$

4.1 Proof of Theorem 4.1

Recall from Sect. 2.4, that $(\xi_t)_{t \geq 0}$ and $(V_t)_{t \geq 0}$ are two processes on $(\Omega, \mathcal{F}, \mathbf{P}_0)$ that are independent: ξ is a standard d -dimensional Brownian motion starting from the origin 0 and V is a subordinator whose Laplace exponent is $\psi^{*'}(\lambda) = \psi'(\lambda) - \alpha$. Recall from (47) that under \mathbf{P}_0 , conditionally given (ξ, V) , $\mathcal{N}^*(dt dW) = \sum_{j \in \mathcal{J}^*} \delta_{(t_j, W^j)}$ is a Poisson point process on $[0, \infty) \times C(\mathbb{R}_+, \mathcal{W})$ with intensity $dV_t \mathbb{N}_{\xi_t}(dW)$. Recall from (48) that for all $a \in \mathbb{R}_+$, we have set $\mathcal{M}_a^* = \sum_{j \in \mathcal{J}^*} \mathbf{1}_{[0,a]}(t_j) \mathcal{M}_j$ where for all $j \in \mathcal{J}^*$, \mathcal{M}_j stands for the occupation measure of the snake W^j as defined in (38). Also recall from (71) the definition of the following random variables:

$$\forall t \geq s \geq 0 \quad N_r(s, t) = \#\{j \in \mathcal{J}^* : s < t_j < t \text{ and } \mathcal{R}_j \cap \overline{B}(0, r) \neq \emptyset\}.$$

Namely $N_r(s, t)$ counts the number of snakes that are grafted on the spatial spine ξ between times s and t and that hit the ball $\overline{B}(0, r)$. We first prove the following lemma that is a consequence of the Blumenthal 0-1 law.

Lemma 4.3 Assume that $\delta > 1$ and that $d > \frac{2\gamma}{\gamma-1}$. Recall from (11) the definition of the gauge function g . There exists a constant $\kappa_{d,\psi} \in [0, \infty]$ that only depends on d and ψ , such that

$$\forall a \in (0, \infty), \quad \mathbf{P}_0\text{-a.s.} \quad \liminf_{r \rightarrow 0+} \frac{\mathcal{M}_a^*(B(0, r))}{g(r)} = \kappa_{d,\psi}. \tag{105}$$

Proof Let $a \in (0, \infty)$ and let $s \in (0, a)$. Observe that if $N_r(s, a) = 0$, then $\mathcal{M}_a^*(B(0, r)) = \mathcal{M}_s^*(B(0, r))$. By Lemma 3.9, there exists a deterministic sequence $(r_n)_{n \geq 0}$ decreasing to 0 such that

$$\sum_{n \geq 0} \mathbf{P}_0(N_{r_n}(s, a) \neq 0) < \infty.$$

By the Borel Cantelli Lemma, \mathbf{P}_0 -a.s. for all sufficiently large n , $N_{r_n}(s, a) = 0$. Since $r \mapsto N_r(s, a)$ is non-decreasing, we get that \mathbf{P}_0 -a.s. for all sufficiently small r , $N_r(s, a) = 0$. Consequently,

$$\forall s \in (0, a), \quad \mathbf{P}_0\text{-a.s.} \quad \liminf_{r \rightarrow 0+} \frac{\mathcal{M}_a^*(B(0, r))}{g(r)} = \liminf_{r \rightarrow 0+} \frac{\mathcal{M}_s^*(B(0, r))}{g(r)}. \tag{106}$$

Let \mathcal{G}_s be the sigma-field generated by $\mathbf{1}_{[0,s]}(t) \mathcal{N}^*(dt dW)$ and completed by the \mathbf{P}_0 -negligible sets. Using properties of Poisson random measures and the Blumenthal zero-one law for ξ , we easily check that $\mathcal{G}_{0+} := \bigcap_{s>0} \mathcal{G}_s$ is \mathbf{P}_0 -trivial: namely, for all $A \in \mathcal{G}_{0+}$, either $\mathbf{P}_0(A) = 0$ or $\mathbf{P}_0(A) = 1$. Then observe that (106) implies that the random variable $\liminf_{r \rightarrow 0} \mathcal{M}_a^*(B(0, r))/g(r)$ is \mathcal{G}_{0+} -measurable. It is therefore \mathbf{P}_0 -a.s. equal to a deterministic constant $\kappa_{d,\psi} \in [0, \infty]$ that does not depends on a . \square

Remark 4.1 We point out that Lemma 4.3 holds true for all gauge functions g . \square

By (50), the previous lemma entails that

$$\mathbf{N}_0\text{-a.e. for } \mathcal{M}\text{-almost all } x, \quad \liminf_{r \rightarrow 0+} \frac{\mathcal{M}(B(x, r))}{g(r)} = \kappa_{d,\psi} \in [0, \infty]. \tag{107}$$

Now, we need to prove that $0 < \kappa_{d,\psi} < \infty$, which is done in two steps.

Lemma 4.4 Assume that $\delta > 1$ and that $d > \frac{2\gamma}{\gamma-1}$. Then, $\kappa_{d,\psi} < \infty$.

Proof Fix $a \in (0, \infty)$ and recall from (67) the definition of T_a that is the sum of the durations of the snakes that are grafted on the spine $(\xi_s)_{0 \leq s \leq a}$. Recall from (71) the definition of $N_r(s, t)$ and from (69) the definition of the last exit time $\vartheta(r)$. Observe that if $\vartheta(2r) \geq a$, then $\mathcal{M}_a^*(B(0, r)) \leq T_a \leq T_{\vartheta(2r)}$. Next note that if $\vartheta(2r) < a$ and if $N_r(\vartheta(2r), a + \vartheta(2r)) = 0$, then, $\mathcal{M}_a^*(B(0, r)) \leq T_{\vartheta(2r)}$. Therefore,

$$\mathbf{P}_0\text{-a.s. on } \{N_r(\vartheta(2r), a + \vartheta(2r)) = 0\}, \quad \mathcal{M}_a^*(B(0, r)) \leq T_{\vartheta(2r)}.$$

Then, for all $r > r' \geq 0$, and all $A > 0$,

$$\begin{aligned} & \mathbf{P}_0\text{-a.s. on } \{N_r(\vartheta(2r), a + \vartheta(2r)) = 0\} \cap \{T_{\vartheta(2r)} - T_{\vartheta(2r')} \leq A\}, \\ & \mathcal{M}_a^*(B(0, r)) \leq A + T_{\vartheta(2r')}. \end{aligned} \tag{108}$$

Recall from (70) the definition of $\gamma(r)$ and observe that $\vartheta(r) \leq \gamma(r)$, which implies

$$\mathbf{P}_0\text{-a.s. } \forall r \in (0, \infty), \quad T_{\vartheta(r)} \leq T_{\gamma(r)}.$$

Thus, by (108), for all $r > r' \geq 0$, and all $A > 0$,

$$\begin{aligned} & \mathbf{P}_0\text{-a.s. on } \{N_r(\vartheta(2r), a + \vartheta(2r)) = 0\} \cap \{T_{\vartheta(2r)} - T_{\vartheta(2r')} \leq A\}, \\ & \mathcal{M}_a^*(B(0, r)) \leq A + T_{\gamma(2r')}, \end{aligned} \tag{109}$$

which allows us to bound $\mathcal{M}_a^*(B(0, r))$ by the subordinator $T_{\gamma(\cdot)}$ studied in section 3.3.

Recall Lemma 3.7, where the sequence $(\rho_n)_{n \geq 1}$ is introduced; recall Lemma 3.11 and Lemma 3.12 that provide a control respectively on $\mathbf{P}_0(N_{\rho_n}(\vartheta(2\rho_n), a + \vartheta(2\rho_n)) = 0)$ and on $T_{\gamma(2\rho_n)}$. For any $n \geq 1$, we next introduce the following random variable:

$$Y_n = \mathbf{1}_{\{N_{\rho_n}(\vartheta(2\rho_n), a + \vartheta(2\rho_n)) = 0\} \cap \{T_{\vartheta(2\rho_n)} - T_{\vartheta(2\rho_{n+1})} \leq g(8\rho_n)\}}. \tag{110}$$

By (109), we get

$$\forall n \geq 1, \quad \mathbf{P}_0\text{-a.s. on } \{Y_n = 1\}, \quad \mathcal{M}_a^*(B(0, \rho_n)) \leq g(8\rho_n) + T_{\gamma(2\rho_{n+1})}.$$

We then define the following event:

$$E = \left\{ \sum_{n \geq 1} Y_n = \infty \right\}.$$

We claim that

$$\mathbf{P}_0(E) > 0. \tag{111}$$

If (111) holds true, then the proof of the lemma is completed as follows: the previous arguments first entail

$$\mathbf{P}_0\text{-a.s. on } E, \quad \liminf_{n \rightarrow \infty} \frac{\mathcal{M}_a^*(B(0, \rho_n))}{g(8\rho_n)} \leq 1 + \limsup_{n \rightarrow \infty} \frac{T_{\gamma(2\rho_{n+1})}}{g(8\rho_n)}. \tag{112}$$

Recall that Assumption $\delta > 1$ implies that g satisfies a C -doubling condition (13), which entails $g(8r) \leq C^3 g(r)$. Then by (112),

$$\mathbf{P}_0\text{-a.s. on } E, \quad \kappa_{d,\psi} = \liminf_{r \rightarrow 0^+} \frac{\mathcal{M}_a^*(B(0, r))}{g(r)} \leq C^3 \left(1 + \limsup_{n \rightarrow \infty} \frac{T_{\gamma(2\rho_{n+1})}}{g(8\rho_n)} \right). \tag{113}$$

By Lemma 3.12 (ii) the right member of (113) is \mathbf{P}_0 -a.s. finite, which completes the proof of the lemma. It only remains to prove our claim (111).

Proof of (111) We use a second moment method. By the independence property of Lemma 3.8 (i), we first get

$$\mathbf{E}_0[Y_n] = \mathbf{P}_0(N_{\rho_n}(\vartheta(2\rho_n), a + \vartheta(2\rho_n)) = 0) \mathbf{P}_0(T_{\vartheta(2\rho_n)} - T_{\vartheta(2\rho_{n+1})} \leq g(8\rho_n)).$$

Then, the lower bound of Lemma 3.11 and the fact that $T_{\vartheta(r)} \leq T_{\gamma(r)}$ entail

$$\begin{aligned} \mathbf{E}_0[Y_n] &\geq c_{15} \mathbf{P}_0(T_{\vartheta(2\rho_n)} - T_{\vartheta(2\rho_{n+1})} \leq g(8\rho_n)) \\ &\geq c_{15} \mathbf{P}_0(T_{\vartheta(2\rho_n)} \leq g(8\rho_n)) \\ &\geq c_{15} \mathbf{P}_0(T_{\gamma(2\rho_n)} \leq g(8\rho_n)). \end{aligned} \tag{114}$$

So by Lemma 3.12 (i), we get

$$\sum_{n \geq 1} \mathbf{E}_0[Y_n] = \infty. \tag{115}$$

Besides, for $n > m \geq 1$, by the independence property of Lemma 3.8 (i)

$$\begin{aligned} \mathbf{E}_0[Y_n Y_m] &\leq \mathbf{P}_0(T_{\vartheta(2\rho_n)} - T_{\vartheta(2\rho_{n+1})} \leq g(8\rho_n); T_{\vartheta(2\rho_m)} - T_{\vartheta(2\rho_{m+1})} \leq g(8\rho_m); \\ &\quad N_{\rho_m}(\vartheta(2\rho_m), a + \vartheta(2\rho_m)) = 0) \\ &\leq \mathbf{P}_0(T_{\vartheta(2\rho_n)} - T_{\vartheta(2\rho_{n+1})} \leq g(8\rho_n)) \mathbf{E}_0[Y_m] \\ &\leq \frac{1}{c_{15}} \mathbf{E}_0[Y_n] \mathbf{E}_0[Y_m], \end{aligned} \tag{116}$$

where the last inequality follows from (114). Therefore, if we denote $L_n = \sum_{1 \leq k \leq n} Y_k$, then (115) and (116) entail

$$\limsup_{n \rightarrow \infty} \frac{\mathbf{E}[L_n^2]}{\mathbf{E}[L_n]^2} \leq \frac{1}{c_{15}} < \infty.$$

and the desired result (111) follows from the Kochen–Stone Lemma. □

The following lemma completes the proof of Theorem 4.1. Its proof relies on a density result on the ψ -Lévy tree that is proved in [12] and that has been recalled here as Lemma 2.5.

Lemma 4.5 *Assume that $\delta > 1$ and that $d > \frac{2\gamma}{\gamma-1}$. Then $\kappa_{d,\psi} > 0$.*

Proof We work with W under \mathbb{N}_0 . The proof consists in lifting to \widehat{W} the estimate of Lemma 2.5 by using the fact that conditionally given H , \widehat{W} is a Gaussian process (see (35) in Sect. 2.4). More precisely, recall that the height process H is the lifetime process of the snake W and recall from (36) the definition of \mathbb{N}_0 that shows that

conditionally given H , the law of W is Q_0^H (see Sect. 2.4). Recall from (28) that d stands for the pseudo-distance in the Lévy tree given by

$$\forall s, t \in [0, \sigma], \quad d(s, t) = H_t + H_s - 2 \inf_{u \in [s \wedge t, s \vee t]} H_u.$$

Let $r, R \in (0, \infty)$ and let $t \in [0, \sigma]$. We set

$$\mathbf{a}(t, r) = \int_0^\sigma \mathbf{1}_{\{d_H(s,t) \leq r\}} ds \quad \text{and} \quad \mathbf{b}(t, r, R) = \int_0^\sigma \mathbf{1}_{\{d_H(s,t) \leq r\} \cap \{\|\widehat{W}_s - \widehat{W}_t\| \geq R\}} ds.$$

The quantity $\mathbf{a}(t, r)$ has been already introduced in Lemma 2.5. First note that

$$\forall t \in [0, \sigma], \quad \mathbf{a}(t, r) \leq \mathbf{b}(t, r, R) + \mathcal{M}(B(\widehat{W}_t, R)). \tag{117}$$

By (35), we then get

$$\begin{aligned} N(dH)\text{-a.e. } \forall t \in [0, \sigma], \quad & Q_0^H [\mathbf{b}(t, r, R)] \\ & \leq \mathbf{a}(t, r) \int_{\mathbb{R}^d \setminus B(0, R/\sqrt{r})} (2\pi)^{-d/2} e^{-\|x\|^2/2} dx. \end{aligned} \tag{118}$$

For any integer $n \geq 2$, we next set $R_n = 2^{-n}$ and $r_n = \frac{1}{4} R_n^2 (\log \log 1/R_n)^{-1}$. An elementary computation entails

$$\forall n \geq 2, \quad \int_{\mathbb{R}^d \setminus B(0, R_n/\sqrt{r_n})} (2\pi)^{-d/2} e^{-\|x\|^2/2} dx \leq c_{22} n^{-3/2},$$

where $c_{22} \in (0, \infty)$ only depends on d (note that the power $3/2$ is not optimal). By (118), we get

$$N(dH) \forall t \in [0, \sigma], \quad Q_0^H \left[\sum_{n \geq 2} \frac{\mathbf{b}(t, r_n, R_n)}{\mathbf{a}(t, r_n)} \right] < \infty.$$

Thus, $N(dH)$ -a.e. for all $t \in [0, \sigma]$, $Q_0^H (\limsup_{n \rightarrow \infty} \mathbf{b}(t, r_n, R_n) / \mathbf{a}(t, r_n) > 0) = 0$. Then, by Fubini,

$$N(dH)\text{-a.e.} \quad Q_0^H \left[\int_0^\sigma \mathbf{1}_{\left\{ \limsup_{n \rightarrow \infty} \frac{\mathbf{b}(t, r_n, R_n)}{\mathbf{a}(t, r_n)} > 0 \right\}} dt \right] = 0,$$

which implies that

$$\mathbb{N}_0\text{-a.e. for } \ell\text{-almost all } t \in [0, \sigma], \quad \lim_{n \rightarrow \infty} \frac{\mathbf{b}(t, r_n, R_n)}{\mathbf{a}(t, r_n)} = 0 \tag{119}$$

(ℓ stands here for the Lebesgue measure on the real line). Recall from (30), the notation $k(r)$:

$$\forall r \in (0, \alpha \wedge e^{-e}), \quad k(r) := \frac{\log \log \frac{1}{r}}{\varphi^{-1} \left(\frac{1}{r} \log \log \frac{1}{r} \right)}.$$

Then, (119) combined with (117), entails

$$\mathbb{N}_0\text{-a.e. for } \ell \text{ almost all } t \in [0, \sigma], \quad \liminf_{n \rightarrow \infty} \frac{\mathbf{a}(t, r_n)}{k(r_n)} \leq \liminf_{n \rightarrow \infty} \frac{\mathcal{M}(B(\widehat{W}_t, R_n))}{k(r_n)}.$$

The definition of $\delta_\varphi \in (0, 1)$ easily implies that there exists a constant $c_{23} \in (0, \infty)$ such that for all sufficiently large n , $c_{23}k(r_n) \geq g(R_n)$. Then, Lemma 2.5 entails that

$$\mathbb{N}_0\text{-a.e. for almost all } t \in [0, \sigma], \quad \liminf_{n \rightarrow \infty} \frac{\mathcal{M}(B(\widehat{W}_t, 2^{-n}))}{g(2^{-n})} \geq \frac{c_1}{c_{23}}.$$

An easy argument involving the doubling property (13) for g completes the proof thanks to (107). □

4.2 Proof of Theorem 4.2

We first introduce a specific decomposition of \mathbb{R}^d into dyadic cubes. We adopt the following notation: we denote by $\lfloor \cdot \rfloor$ the integer part application and we write \log_2 for the logarithm in base 2; we fix $d > \frac{2\gamma}{\gamma-1}$ and we set

$$p := \lfloor \log_2(4\sqrt{d}) \rfloor,$$

so that $2^p > 2\sqrt{d}$. To simplify notation, we set $\mathcal{D}_n = 2^{-n-p}\mathbb{Z}^d$, for any $n \geq 0$. For any $y = (y_1, \dots, y_d)$ in \mathcal{D}_n , we also set

$$D_n(y) = \prod_{j=1}^d \left[y_j - \frac{1}{2}2^{-n}; y_j + \frac{1}{2}2^{-n} \right) \quad \text{and}$$

$$D_n^\bullet(y) = \prod_{j=1}^d \left[y_j - \frac{1}{2}2^{-n-p}; y_j + \frac{1}{2}2^{-n-p} \right).$$

It is easy to check the following properties.

- Prop(1) If y, y' are distinct points in \mathcal{D}_n , then $D_n^\bullet(y) \cap D_n^\bullet(y') = \emptyset$.
- Prop(2) Let $y \in \mathcal{D}_n$. Then, we have

$$D_n^\bullet(y) \subset \overline{B} \left(y, \frac{1}{2}2^{-n-p}\sqrt{d} \right) \subset \overline{B}(y, 2^{-n-p}\sqrt{d}) \subset D_n(y).$$

For any $r < (2d)^{-1}$, we set $n(r) = \lfloor \log_2(r^{-1}(1 + 2^{-p})\sqrt{d}) \rfloor$, so that the following inequalities hold:

$$\frac{1}{2}(1 + 2^{-p})\sqrt{d} 2^{-n(r)} < r \leq (1 + 2^{-p})\sqrt{d} 2^{-n(r)}. \tag{120}$$

Next, for any $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ and for $j \in \{1, \dots, d\}$, we set

$$y_j = 2^{-n(r)-p} \left\lfloor x_j 2^{n(r)+p} + \frac{1}{2} \right\rfloor.$$

Therefore, $y = (y_1, \dots, y_d) \in \mathcal{D}_{n(r)}$ and we easily check the following:

- Prop(3) The point x belongs to $D_{n(r)}^\bullet(y)$ and $D_{n(r)}(y) \subset B(x, r)$.

We work under \mathbb{N}_0 . Recall Lemma 3.16: we fix

$$\kappa_1 \in (0, \kappa_0) \quad \text{such that} \quad c_{19}\kappa_1^{-c_{20}} > 2. \tag{121}$$

Since we assume $\delta > 1$, g satisfies the doubling condition (13), which implies that there exists $\kappa_2 \in (0, \infty)$, that only depends on d, ψ and κ_1 , such that for all sufficiently large n ,

$$\kappa_2 g(2^{-n}) \leq \kappa_1 g\left(\frac{1}{2}2^{-n-p}\sqrt{d}\right). \tag{122}$$

We then fix $A > 100$ and for any n such that $2^{-n} \leq 1/(2A)$, we set

$$U_n(A) = \sum_{\substack{y \in \mathcal{D}_n \\ 1/A \leq \|y\| \leq A}} g(\sqrt{d}(1 + 2^{-p})2^{-n}) \mathbf{1}_{\{\mathcal{M}(D_n(y)) \leq \kappa_2 g(2^{-n})\} \cap \{\mathcal{R} \cap D_n^\bullet(y) \neq \emptyset\}}.$$

We first prove the following lemma.

Lemma 4.6 *Assume that $\delta > 1$ (so that (122) holds) and that $d \geq 3$. Then, for all $A > 100$,*

$$\mathbb{N}_0\text{-a.e.} \quad \lim_{N \rightarrow \infty} \sum_{n \geq N} U_n(A) = 0. \tag{123}$$

Proof We fix n such that $2^{-n} \leq 1/(2A)$ and we fix $y \in \mathcal{D}_n$ such that $1/A \leq \|y\| \leq A$. By Prop(2) and (122), we get

$$\begin{aligned} & \mathbb{N}_0(\mathcal{M}(D_n(y)) \leq \kappa_2 g(2^{-n}); \mathcal{R} \cap D_n^\bullet(y) \neq \emptyset) \\ & \leq \mathbb{N}_0(\mathcal{M}(B(y, 2^{-n-p}\sqrt{d})) \leq \kappa_1 g\left(\frac{1}{2}2^{-n-p}\sqrt{d}\right); \\ & \quad \mathcal{R} \cap \bar{B}\left(y, \frac{1}{2}2^{-n-p}\sqrt{d}\right) \neq \emptyset) \\ & = \mathbb{N}_{-y}(\mathcal{M}(B(0, 2^{-n-p}\sqrt{d})) \leq \kappa_1 g\left(\frac{1}{2}2^{-n-p}\sqrt{d}\right); \\ & \quad \mathcal{R} \cap \bar{B}\left(0, \frac{1}{2}2^{-n-p}\sqrt{d}\right) \neq \emptyset), \end{aligned}$$

the last equality being an immediate consequence of the invariance of the snake under translation. We next apply Lemma 3.16 with $x = -y$ and $r = \frac{1}{2}2^{-n-p}\sqrt{d}$ and $\kappa = \kappa_1$ that satisfies (121); thus, there exists $c_{24}, c_{25} \in (0, \infty)$, that only depends on d and ψ , such that

$$\begin{aligned} \mathbb{N}_0(\mathcal{M}(D_n(y)) \leq \kappa_2 g(2^{-n}) ; \mathcal{R} \cap D_n^\bullet(y) \neq \emptyset) \\ \leq c_{24}(2^{-n-p})^{d-2} \|y\|^{2-d} n^{-2} \psi'^{-1}(c_{25} 2^{2n+2p}). \end{aligned}$$

By Lemma 2.3 and the doubling property (13) of g , there exists $c_{26} \in (0, \infty)$, that only depends on d and ψ , such that for all sufficiently large n ,

$$g(\sqrt{d}(1 + 2^{-p})2^{-n}) \psi'^{-1}(c_{25} 2^{2n+2p}) \leq c_{26} 2^{-2n-2p},$$

which entails the following:

$$\begin{aligned} g(\sqrt{d}(1 + 2^{-p})2^{-n}) \mathbb{N}_0(\mathcal{M}(D_n(y)) \leq \kappa_2 g(2^{-n}) ; \mathcal{R} \cap D_n^\bullet(y) \neq \emptyset) \\ \leq c_{27}(2^{-n-p})^d \|y\|^{2-d} n^{-2}, \end{aligned} \tag{124}$$

where $c_{27} = c_{24}c_{26}$. Elementary arguments entail the following inequalities:

$$\begin{aligned} \mathbb{N}_0(U_n(A)) &\leq c_{27} n^{-2} \sum_{\substack{y \in D_n \\ 1/A \leq \|y\| \leq A}} (2^{-n-p})^d \|y\|^{2-d} \\ &\leq c_{28} n^{-2} \int \mathbf{1}_{\{1/A \leq \|x\| \leq A\}} \|x\|^{2-d} dx \\ &\leq c_{29} n^{-2} \int_{1/A}^A \rho d\rho \leq c_{30} A^2 n^{-2}, \end{aligned}$$

where $c_{28}, c_{29}, c_{30} \in (0, \infty)$ only depend on d and ψ . Therefore, $\mathbb{N}_0(\sum_{n \geq N} U_n(A)) < \infty$, which easily implies the lemma. □

Recall from (16) the definition of the g -packing measure \mathcal{P}_g . We next prove the following lemma.

Lemma 4.7 *Assume that $\delta > 1$ and that $d > \frac{2\gamma}{\gamma-1}$. Then,*

$$\mathbb{N}_0\text{-a.e. } \mathcal{P}_g \left(\left\{ x \in \mathcal{R} : \liminf_{r \rightarrow 0^+} g(r)^{-1} \mathcal{M}(B(x, r)) \neq \kappa_{d,\psi} \right\} \right) = 0, \tag{125}$$

where $\kappa_{d,\psi}$ is the constant appearing in Theorem 4.1.

Proof We fix $A > 100$. By Theorem 4.1 and Lemma 4.6 there exists a Borel subset \mathcal{W}_A of \mathcal{W} such that $\mathbb{N}_0(\mathcal{W} \setminus \mathcal{W}_A) = 0$ and such that on \mathcal{W}_A , (104) and (123) hold true. We shall work deterministically on \mathcal{W}_A .

Let B be any Borel subset of $\{x \in \mathbb{R}^d : 1/A \leq \|x\| \leq A\}$. Let $\varepsilon \in (0, \infty)$ and let $\overline{B}(x_1, r_1), \dots, \overline{B}(x_k, r_k)$ be any closed ε -packing of $B \cap \mathcal{R}$. Namely, the later balls are disjoint, their centres belong to $B \cap \mathcal{R}$ and their radii are smaller than ε . Let $c_{31} \in (0, \infty)$ be a constant to be specified later. First observe that

$$\begin{aligned} \sum_{i=1}^k g(r_i) &= \sum_{i=1}^k g(r_i) \mathbf{1}_{\{\mathcal{M}(B(x_i, r_i)) > c_{31} g(r_i)\}} + \sum_{i=1}^k g(r_i) \mathbf{1}_{\{\mathcal{M}(B(x_i, r_i)) \leq c_{31} g(r_i)\}} \\ &\leq c_{31}^{-1} \mathcal{M}(B^{(\varepsilon)}) + \sum_{i=1}^k g(r_i) \mathbf{1}_{\{\mathcal{M}(B(x_i, r_i)) \leq c_{31} g(r_i)\}}, \end{aligned} \tag{126}$$

where we have set $B^{(\varepsilon)} = \{x \in \mathbb{R}^d : \text{dist}(x, B) \leq \varepsilon\}$. Next, fix $1 \leq i \leq k$; recall notation $n(r_i)$ from (120) and denote by y_i the point of $\mathcal{D}_{n(r_i)}$ corresponding to x_i such that Prop(3) holds true. Therefore, by (120), we have

$$\begin{aligned} \mathcal{M}(B(x_i, r_i)) \leq c_{31} g(r_i) \quad \text{and} \quad x_i \in B \cap \mathcal{R} &\implies \\ \mathcal{M}(D_{n(r_i)}(y_i)) \leq c_{31} g((1 + 2^{-p})\sqrt{d}2^{-n(r_i)}) \quad \text{and} \\ \mathcal{R} \cap D_{n(r_i)}^\bullet(y_i) \neq \emptyset. \end{aligned}$$

We use the doubling condition (13) to choose c_{31} such that $c_{31} g((1 + 2^{-p})\sqrt{d}2^{-n(r)}) \leq \kappa_2 g(2^{-n(r)})$ for all sufficiently small $r \in (0, 1)$. Thus, we get

$$\sum_{i=1}^k g(r_i) \mathbf{1}_{\{\mathcal{M}(B(x_i, r_i)) \leq c_{31} g(r_i)\}} \leq \sum_{n: 2^{-n} \leq c_{32} \varepsilon} U_n(A),$$

where $c_{32} = 2((1 + 2^{-p})\sqrt{d})^{-1}$. Since W belongs to \mathcal{W}_A where (123) holds, this inequality combined with (126) implies the following.

$$\mathcal{P}_g(B \cap \mathcal{R}) \leq \mathcal{P}_g^*(B \cap \mathcal{R}) \leq c_{31}^{-1} \mathcal{M}\left(\bigcap_{\varepsilon > 0} B^{(\varepsilon)}\right). \tag{127}$$

We next apply (127) with $B = B_A$ given by

$$B_A = \left\{ x \in \mathcal{R} : 1/A \leq \|x\| \leq A \text{ and } \liminf_{r \rightarrow 0^+} g(r)^{-1} \mathcal{M}(B(x, r)) \neq \kappa_d, \psi \right\}.$$

Therefore $\mathcal{P}_g(B_A) < \infty$. Suppose now that $\mathcal{P}_g(B_A) > 0$. Then, as a consequence of (18), there exists a closed subset F , with $F \subset B_A$, such that $\mathcal{P}_g(F) > 0$. Since F is closed then $F = \bigcap_{\varepsilon > 0} F^{(\varepsilon)}$; since F is a subset of B_A and since $W \in \mathcal{W}_A$ (where (104) holds true), we get $\mathcal{M}(F) \leq \mathcal{M}(B_A) = 0$ and by (127) applied to $B = F$, we obtain $\mathcal{P}_g(F) = 0$, which rises a contradiction. Thus, we have proved that \mathbb{N}_0 -a.e. $\mathcal{P}_g(B_A) = 0$, which easily entails the lemma by letting A go to ∞ , since $\mathcal{P}_g(\{0\}) = 0$. □

End proof of Theorem 4.2 By Theorem 4.1 and Lemma 4.7 there exists a Borel subset \mathcal{W}^* of \mathcal{W} such that $\mathbb{N}_0(\mathcal{W} \setminus \mathcal{W}^*) = 0$ and such that (104) and (125) hold true on \mathcal{W}^* . We fix $W \in \mathcal{W}^*$ and we set

$$\text{Good} = \left\{ x \in \mathcal{R} : \liminf_{r \rightarrow 0^+} g(r)^{-1} \mathcal{M}(B(x, r)) = \kappa_{d, \psi} \right\} \quad \text{and} \quad \text{Bad} = \mathcal{R} \setminus \text{Good}.$$

Let B be any Borel subset of \mathbb{R}^d . By (104) and (125), we have

$$\mathcal{M}(B \cap \text{Bad}) = \mathcal{P}_g(B \cap \mathcal{R} \cap \text{Bad}) = 0.$$

Then, we apply Lemma 2.2 to $\text{Good} \cap B$ and we get

$$\mathcal{M}(B \cap \text{Good}) = \kappa_{d, \psi} \mathcal{P}_g(B \cap \mathcal{R} \cap \text{Good}).$$

Therefore, on \mathcal{W}^* , for Borel subset B of \mathbb{R}^d , $\mathcal{M}(B) = \kappa_{d, \psi} \mathcal{P}_g(B \cap \mathcal{R})$, which completes the proof of Theorem 4.2. □

4.3 Proof of Theorem 1.2

We derive Theorem 1.2 from Theorem 4.2. To that end, we first need an upper bound of the upper box-counting dimension of \mathcal{R} under \mathbb{N}_x . Let us briefly recall the definition of the box-counting dimensions of a bounded subset $K \subset \mathbb{R}^d$: let $\varepsilon \in (0, \infty)$ and let $n_\varepsilon(K)$ stand for the minimal number of open balls of radius ε that are necessary to cover K . Then,

$$\underline{\dim}(K) = \liminf_{\varepsilon \rightarrow 0} \frac{\log n_\varepsilon(K)}{\log 1/\varepsilon} \quad \text{and} \quad \overline{\dim}(K) = \limsup_{\varepsilon \rightarrow 0} \frac{\log n_\varepsilon(K)}{\log 1/\varepsilon}. \tag{128}$$

Fix $x \in \mathbb{R}^d$ and recall Lemma 2.6 that asserts that for any $q \in (0, \frac{\gamma-1}{2\gamma})$, \mathbb{N}_x -a.e. $(\widehat{W}_s)_{s \in [0, \sigma]}$ is q -Hölder continuous. As already mentioned in Comment 1.2, it easily implies the following:

$$\mathbb{N}_x\text{-a.e.} \quad \overline{\dim}(\mathcal{R}) \leq \frac{2\gamma}{\gamma - 1}, \tag{129}$$

where \mathcal{R} is the range of the Lévy snake as defined (37). We next prove the following lemma.

Lemma 4.8 *Assume that $\gamma > 1$ and that $d > \frac{2\gamma}{\gamma-1}$. Let $x \in \mathbb{R}^d$. For any compact subset K such that $\overline{\dim}(K) \leq \frac{2\gamma}{\gamma-1}$, we have \mathbb{N}_x -a.e. $\mathcal{M}(K) = 0$.*

Proof Let us first assume that $x \notin K$ and set $k := \inf_{y \in K} \|x - y\| > 0$. For any $\varepsilon \in (0, k/2)$, denote by n_ε the minimal number of balls with radius ε that are necessary to

cover K , and denote by $B(x_1^\varepsilon, \varepsilon), \dots, B(x_{n_\varepsilon}^\varepsilon, \varepsilon)$ such balls. Then, (46) combined with standard estimates of d -dimensional Green function entail the following inequalities.

$$\begin{aligned} \mathbb{N}_x(\mathcal{M}(K)) &\leq \sum_{i=1}^{n_\varepsilon} \mathbb{N}_x(\mathcal{M}(B(x_i^\varepsilon, \varepsilon))) \\ &\leq \sum_{i=1}^{n_\varepsilon} \int_0^\infty e^{-\alpha a} \mathbf{P}_x(\xi_a \in B(x_i^\varepsilon, \varepsilon)) \, da \\ &\leq c_{33} \sum_{i=1}^{n_\varepsilon} \int_{B(x_i^\varepsilon, \varepsilon)} \|x - y\|^{2-d} \, dy \\ &\leq c_{34} k^{2-d} \varepsilon^d n_\varepsilon, \end{aligned}$$

where $c_{33}, c_{34} \in (0, \infty)$ only depend on d . Since $d > \frac{2\gamma}{\gamma-1} \geq \overline{\dim}(K)$, the previous inequality implies that $\mathbb{N}_x(\mathcal{M}(K)) = 0$ as $\varepsilon \rightarrow 0$.

Let us now consider the general case: for any $r > 0$, the previous case applies to the compact set $K \setminus B(x, r)$ and we get

$$\mathbb{N}_x\text{-a.e. } \mathcal{M}(K) = \mathcal{M}(K \cap B(x, r)) + \mathcal{M}(K \setminus B(x, r)) \leq \mathcal{M}(B(x, r)) ,$$

which implies the desired result as $r \rightarrow 0$ since \mathcal{M} is diffuse. □

End of the proof of Theorem 1.2 We use an argument due to Le Gall in [28, pp. 312–313]. Theorem 4.2 and Lemma 4.8 imply that for any compact set K such that $\overline{\dim}(K) \leq \frac{2\gamma}{\gamma-1}$, and for any $x \in \mathbb{R}^d$,

$$\mathbb{N}_x\text{-a.e. } \mathcal{P}_g(K \cap \mathcal{R}) = 0. \tag{130}$$

Recall the connection (39) in Theorem 2.7 between \mathbf{R}, \mathbf{M} and the excursions $W^j, j \in \mathcal{J}$, of the Brownian snake. An easy argument on Poisson point processes combined with (129) and (130) implies that almost surely $\mathcal{P}_g(\mathcal{R}_{W^j} \cap \mathcal{R}_{W^i}) = 0$ for any $i \neq j$ in \mathcal{J} . Then, (39) entails

$$\mathcal{P}_g(\cdot \cap \mathbf{R}) = \sum_{j \in \mathcal{J}} \mathcal{P}_g(\cdot \cap \mathcal{R}_{W^j}).$$

Theorem 2.7 and (39) thus imply

$$\kappa_{d,\psi} \mathcal{P}_g(\cdot \cap \mathbf{R}) = \sum_{j \in \mathcal{J}} \kappa_{d,\psi} \mathcal{P}_g(\cdot \cap \mathcal{R}_{W^j}) = \sum_{j \in \mathcal{J}} \mathcal{M}_{W^j} = \mathbf{M} ,$$

which is the desired result.

4.4 Dimension of the range of the ψ -SBM

We now prove Theorem 1.1. Recall from Sect. 2.4, that $(\xi_t)_{t \geq 0}$ and $(V_t)_{t \geq 0}$ are two processes on $(\Omega, \mathcal{F}, \mathbf{P}_0)$ that are independent: ξ is a standard d -dimensional Brownian motion starting from the origin 0 and V is a subordinator whose Laplace exponent is $\psi^{*'}(\lambda) = \psi'(\lambda) - \alpha$. Recall from (47) that under \mathbf{P}_0 , conditionally given (ξ, V) , $\mathcal{N}^*(dt dW) = \sum_{j \in \mathcal{J}^*} \delta_{(t_j, W^j)}$ is a Poisson point process on $[0, \infty) \times C(\mathbb{R}_+, \mathcal{W})$ with intensity $dV_t \mathbb{N}_{\xi_t}(dW)$. Recall from (48) that for all $a \in \mathbb{R}_+$, we have set $\mathcal{M}_a^* = \sum_{j \in \mathcal{J}^*} \mathbf{1}_{[0, a]}(t_j) \mathcal{M}_j$ where for all $j \in \mathcal{J}^*$, \mathcal{M}_j stands for the occupation measure of the snake W^j as defined in (38). Also recall from (71) the definition of the following random variables:

$$\forall t \geq s \geq 0 \quad N_r(s, t) = \#\{j \in \mathcal{J}^* : s < t_j < t \text{ and } \mathcal{R}_j \cap \bar{B}(0, r) \neq \emptyset\}.$$

Namely $N_r(s, t)$ counts the number of snakes that are grafted on the spatial spine ξ between times s and t and that hit the ball $\bar{B}(0, r)$.

Lemma 4.9 *Assume that $\delta > 1$ and that $d > \frac{2\delta}{\delta-1}$. Then, for all $a \in (0, \infty)$, and for all $u \in (0, \frac{2\gamma}{\gamma-1})$,*

$$\mathbf{P}_0\text{-a.s.} \quad \liminf_{r \rightarrow 0^+} r^{-u} \mathcal{M}_a^*(B(0, r)) < \infty. \tag{131}$$

Proof The proof is very similar to that of Lemma 4.4. We present only the main steps. Recall from Lemma 3.8 the properties of the processes $(T_{\vartheta(r)})_{r \geq 0}$ and $(T_{\gamma(r)})_{r \geq 0}$. Recall from (71) the definition of $N_r(s, t)$. Recall from Lemma 3.13 the definition of the sequence $(s_n)_{n \geq 0}$. Then, for all $n \geq 0$, we set

$$Y'_n = \mathbf{1}_{\{N_{s_n}(\vartheta(2s_n), a + \vartheta(2s_n)) = 0\} \cap \{T_{\vartheta(2s_n)} - T_{\vartheta(2s_{n+1})} \leq s_n''\}}. \tag{132}$$

Reasoning as in the proof of (109), we get

$$\mathbf{P}_0\text{-a.s. on } \{Y'_n = 1\}, \quad \mathcal{M}_a^*(B(0, s_n)) \leq s_n'' + T_{\gamma(2s_{n+1})}. \tag{133}$$

Thus, if we show that

$$\mathbf{P}_0 \left(\sum_{n \geq 1} Y'_n = \infty \right) > 0, \tag{134}$$

we get (131), by use of Lemma 3.13 (ii).

The inequality (134) is obtained using the Kochen–Stone Lemma, as in the proof of (111). Indeed, under the assumption $d > \frac{2\delta}{\delta-1}$, Lemma 3.10 entails that for all $n \in \mathbb{N}$, $\mathbf{P}_0(N_{s_n}(\vartheta(2s_n), a + \vartheta(2s_n)) = 0) \geq c_{14} > 0$; we then argue exactly as in the proof of Lemma 4.4 to obtain (134). We leave the details to the reader. \square

Proof of Theorem 1.1 Assume that $\delta > 1$ and that $d > \frac{2\delta}{\delta-1}$. Recall from (37) the definition of \mathcal{R} , the range of the Lévy snake. By Theorem 2.7 and by spatial invariance

of the snake, it is sufficient to prove

$$\mathbb{N}_0\text{-a.e.} \quad \dim_p(\mathcal{R}) = \overline{\dim}(\mathcal{R}) = \frac{2\gamma}{\gamma - 1}.$$

Recall that for every bounded subset $K \subset \mathbb{R}^d$, $\dim_p(A) \leq \overline{\dim}(A)$ (see e.g. Falconer [21]). By (129), it then only remains to prove

$$\mathbb{N}_0\text{-a.e.} \quad \dim_p(\mathcal{R}) \geq \frac{2\gamma}{\gamma - 1}. \quad (135)$$

Lemma 4.9 combined with (50) implies that for all $u \in (0, \frac{2\gamma}{\gamma-1})$,

$$\mathbb{N}_0\text{-a.e. for } \mathcal{M}\text{-almost all } x, \quad \liminf_{r \rightarrow 0^+} r^{-u} \mathcal{M}(B(x, r)) < \infty, \quad (136)$$

which implies that $\mathbb{N}_0\text{-a.e. } \dim_p(\mathcal{R}) \geq u$ by the comparison results stated here as Theorem 2.1. This entails (135) and the proof of Theorem 1.1 is completed. \square

Acknowledgements We warmly thank J.-F. Delmas, E. Perkins and an anonymous referee for several comments that improved a first version of this article.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

References

1. Bertoin, J.: Lévy Processes, Cambridge Tracts in Mathematics, vol. 121. Cambridge University Press, Cambridge (1996)
2. Besicovitch, A.S.: A general form of the covering principle and relative differentiation of additive functions. Proc. Camb. Philos. Soc. **41**, 103–110 (1945)
3. Bingham, N.H.: Continuous branching processes and spectral positivity. Stochastic Process. Appl. **4**, 217–242 (1976)
4. Blumenthal, R.M., Gettoor, R.K.: Sample functions of stochastic processes with stationary independent increments. J. Math. Mech. **10**, 493–516 (1961)
5. Bolthausen, E., Perkins, E., van der Vaart, A.: *Lectures on Probability Theory and Statistics*, Lecture Notes in Mathematics, vol. 1781. Springer, Berlin (2002) (lectures from the 29th Summer School on Probability Theory held in Saint-Flour, July 8–24, 1999, edited by Pierre Bernard)
6. Dawson, D.A., Hochberg, K.J.: The carrying dimension of a stochastic measure diffusion. Ann. Probab. **7**(4), 693–703 (1979)
7. Dawson, D.A., Iscoe, I., Perkins, E.A.: Super-Brownian motion: path properties and hitting probabilities. Probab. Theory Related Fields **83**(1–2), 135–205 (1989)
8. Dawson, D.A., Perkins, E.A.: Historical processes. Mem. Am. Math. Soc. **93**(454), iv+179 (1991)
9. Delmas, J.-F.: Some properties of the range of the super-Brownian motion. Probab. Theory Related Fields **114**(4), 505–547 (1999)
10. Duquesne, T.: The packing measure of the range of super-Brownian motion. Ann. Probab. **37**(6), 2431–2458 (2009)
11. Duquesne, T.: Packing and Hausdorff measures of stable trees. In: Lévy matters I, Lecture Notes in Mathematics, vol. 2001, pp. 93–136. Springer, Berlin (2010)
12. Duquesne, T.: The exact packing measure of Lévy trees. Stochastic Process. Appl. **122**, 968–1002 (2012)

13. Duquesne, T., Le Gall, J.-F.: Random trees, Lévy processes and spatial branching processes. *Astérisque* **281** (2002)
14. Duquesne, T., Le Gall, J.-F.: Probabilistic and fractal aspects of Lévy trees. *Probab. Theory Related Fields* **131**(4), 553–603 (2005)
15. Duquesne, T., Le Gall, J.-F.: The Hausdorff measure of stable trees. *ALEA Lat. Am. J Probab. Math. Stat.* **1**, 393–415 (2006)
16. Dynkin, E.B.: A probabilistic approach to one class of nonlinear differential equations. *Probab. Theory Related Fields* **89**(1), 89–115 (1991)
17. Dynkin, E.B.: *Diffusions, Superdiffusions and Partial Differential Equations*, American Mathematical Society Colloquium Publications, vol. 50. American Mathematical Society, Providence (2002)
18. Dynkin, E.B.: *Superdiffusions and Positive Solutions of Nonlinear Partial Differential Equations*, University Lecture Series, vol. 34. American Mathematical Society, Providence (2004) (Appendix A by J.-F. Le Gall and Appendix B by I. E. Verbitsky)
19. Edgar, G.A.: Centered Densities and Fractal Measures. *N. Y. J. Math.* **13**, 33–87 (2007, electronic)
20. Etheridge, A.M.: *An Introduction to Superprocesses*, University Lecture Series, vol. 20, vol. 20. American Mathematical Society, Providence (2000)
21. Falconer, K.: *Fractal Geometry*. Wiley, Chichester (1990). (Mathematical foundations and applications)
22. Gettoor, R.K.: The Brownian escape process. *Ann. Probab.* **7**(5), 864–867 (1979)
23. Hesse, M., Kyprianou, A.E.: The mass of super-Brownian motion upon exiting balls and Sheu’s compact support condition. *Stochastic Process. Appl.* **124**(6), 2003–2022 (2014)
24. Jiřina, M.: Stochastic branching processes with continuous state space. *Czech. Math. J.* **8**(83), 292–313 (1958)
25. Keller, J.B.: On solutions of $\Delta u = f(u)$. *Commun. Pure Appl. Math.* **10**, 503–510 (1957)
26. Lamperti, J.: Continuous state branching processes. *Bull. Am. Math. Soc.* **73**, 382–386 (1967)
27. Lamperti, J.: The limit of a sequence of branching processes. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **7**, 271–288 (1967)
28. Le Gall, J.-F.: The Hausdorff measure of the range of super-Brownian motion. In: *Perplexing Problems in Probability*, *Progr. Probab.*, vol. 44, pp. 285–314. Birkhäuser, Boston (1999)
29. Le Gall, J.-F.: *Spatial Branching Processes, Random Snakes and Partial Differential Equations*. Lectures in Mathematics ETH Zürich. Birkhäuser, Basel (1999)
30. Le Gall, J.-F., Le Jan, Y.: Branching processes in Lévy processes: the exploration process. *Ann. Probab.* **26**(1), 213–252 (1998)
31. Le Gall, J.-F., Perkins, E.A.: The Hausdorff measure of the support of two-dimensional super-Brownian motion. *Ann. Probab.* **23**(4), 1719–1747 (1995)
32. Le Gall, J.-F., Perkins, E.A., Taylor, S.J.: The packing measure of the support of super-Brownian motion. *Stochastic Process. Appl.* **59**(1), 1–20 (1995)
33. Li, Z.: *Measure-Valued Branching Markov Processes* (New York). Probability and its Applications. Springer, Heidelberg (2011)
34. Perkins, E.: A space-time property of a class of measure-valued branching diffusions. *Trans. Am. Math. Soc.* **305**(2), 743–795 (1988)
35. Perkins, E.: The Hausdorff measure of the closed support of super-Brownian motion. *Ann. Inst. H. Poincaré Probab. Stat.* **25**(2), 205–224 (1989)
36. Pitman, J.W.: One-dimensional Brownian motion and the three-dimensional Bessel process. *Adv. Appl. Probab.* **7**(3), 511–526 (1975)
37. Revuz, D., Yor, M.: *Continuous Martingales and Brownian Motion*. Grundlehren der mathematischen Wissenschaften. Springer, Berlin (2004)
38. Sheu, Y.-C.: Asymptotic behavior of superprocesses. *Stochastics Stochastics Rep.* **49**(3–4), 239–252 (1994)
39. Taylor, S.J., Tricot, C.: Packing measure, and its evaluation for a Brownian path. *Trans. Am. Math. Soc.* **288**(2), 679–699 (1985)